(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 29 March 2001 (29.03.2001)

(10) International Publication Number WO 01/21772 A2

(51) International Patent Classification7:

C12N 9/00

PCT/US00/25993 (21) International Application Number:

(22) International Filing Date: 21 September 2000 (21.09.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

21 September 1999 (21.09.1999) US 09/400,494 7 June 2000 (07.06.2000) US 60/210,072 28 July 2000 (28.07.2000) US 60/221,836 US 60/227,860 24 August 2000 (24.08.2000)

(71) Applicant: OMNIGENE BIOPRODUCTS [US/US]; 763(d) Concord Avenue, Cambridge, MA 02138 (US)...

(72) Inventors: YOCUM, Rogers, R.; 4 Orchard Lane, Lexington, MA 02420 (US). PATTERSON, Thomas, A.; 89 Church Street, North Attleboro, MA 02760 (US). HER-MANN, Theron; 18 Chilhowie Drive, Kinnelon, NJ 07405 (US). PERO, Janice, G.; 20 Solomon Pierce Road, Lexington, MA 02420 (US).

(74) Agents: HANLEY, Elizabeth, A. et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (national): AU, BR, CA, CN, CZ, HU, ID, IL, IN, JP, KR, LT, LV, MX, NO, PL, RU, SG, SK, TR, UA, ZA.

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published:

Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS

(57) Abstract: The present invention features methods of producing panto-compounds (e.g., pantothenate) using microorganisms in which the pantothenate biosynthetic pathway and/or the isoleucine-valine biosynthetic pathway and/or the coenzymeA biosynthetic pathway has been manipulated. Methods featuring ketopantoate reductase overexpressing microorganisms as well as aspartate α-decarboxylase overexpressing microorganisms are provided. Methods of producing panto-compounds in a precursor-independent manner and in high yield are described. Recombinant microorganisms, vectors, isolated nucleic acid molecules, genes and gene products useful in practicing the above methodologies are also provided. The present invention also features a previously unidentified microbial pantothenate kinase gene, coal, as well as methods of producing panto-compounds utilizing microorganisms having modified pantothenate kinase activity. Recombinant microorganisms, vectors, isolatedcoal nucleic acid molecules and purified CoaX proteins are featured. Also featured are methods for identifying pantothenate kinase modulators utilizing the recombinant microorganisms and/or purified CoaX proteins of the present invention.

012177262 1 1

10

15

20

25

30

35

METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS

Background of the Invention

Pantothenate, also known as pantothenic acid or vitamin B5, is a member of the B complex of vitamins and is a nutritional requirement for mammals, including livestock and humans (e.g., from food sources, as a water soluble vitamin supplement or as a feed additive). In cells, pantothenate is used primarily for the biosynthesis of coenzyme A (CoA) and acyl carrier protein (ACP). These coenzymes function in the metabolism of acyl moieties which form thioesters with the sulfhydryl group of the 4'-phosphopantetheine portion of these molecules. These coenzymes are essential in all cells, participating in over 100 different intermediary reactions in cellular metabolism.

The conventional means of synthesizing pantothenate (in particular, the bioactive D isomer) is via chemical synthesis from bulk chemicals, a process which is hampered by excessive substrate cost as well as the requirement for optical resolution of racemic intermediates (e.g., resolution of DL-pantolactone to obtain D-pantolactone for chemical condensation with β-alanine). Accordingly, researchers have recently looked to bacterial or microbial systems that produce enzymes useful in pantothenate biosynthesis processes (as bacteria are themselves capable of synthesizing pantothenate). In particular, bioconversion processes have been evaluated as a means of favoring production of the D isomer of pantothenic acid, e.g., using microorganisms which selectively hydrolyze a DL-pantothenic acid ester to D-pantothenic acid; microorganisms which selectively decompose L-pantolactone resulting in D-pantolactone alone; and microorganisms which selectively hydrolyze DL-pantolactone to D-pantoic acid.

There is still, however, significant need for improved pantothenate production processes, in particular, for processes requiring reduced quantities of substrates and/or less expensive substrates. To this end, methods of direct microbial synthesis have recently been examined as a means of improving D-pantothenate production. In microbes, pantothenate biosynthetis is a multistep pathway resulting in condensation of pantoate (derived from α -ketoisovalerate) and β -alanine to form D-pantothenate. The isoleucine-valine (ilv) pathway biosynthetic enzymes, acetohydroxyacid synthetase (the ilvBN or alsS gene product), acetohydroxyacid isomeroreductase (the ilvC gene product) and dihydroxyacid dehydratase (the ilvD gene product) catalyze the conversion of pyruvate to α -ketoisovalerate. The reactions are further catalyzed by the pantothenate (pan) pathway biosynthetic enzymes ketopantoate hydroxymethyltransferase (the panB gene product), ketopantoate reductase (the panE gene product), aspartate- α -

20

25

decarboxylase (the panD gene product) and pantothenate synthetase (the panC gene product).

The genes encoding the enzymes involved in the biosynthesis of pantothenic acid in Salmonella typhimurium and Escherichia coli have recently been identified and characterized (Frodyma and Downs (1998) J. Biol. Chem. 273:5572-5576 and Jackowski (1996) pp. 687-694, In Neidhardt et al (ed.) Escherichia coli and Salmonella: Cellular and Molecular Biology, 2nd ed. Am. Soc. Microbiol. Wash, D.C). In E. coli, for example, the biosynthesis of pantothenic acid consists of four key steps. The first reaction is catalyzed by the panB gene product, ketopantoate hydroxymethyltransferase, and uses the L-valine intermediate α -ketoisovalerate to generate ketopantoate, which is subsequently reduced to pantoate by the panE gene product, ketopantoate reductase. The panD gene product, aspartate- α -decarboxylase, generates β -alanine from aspartate. The panC gene product, pantothenate synthetase, subsequently ligates \$-alanine with pantoate to yield D-pantothenate.

The authors Dusch et al. described the identification of the Corynebacterium glutamicum panD gene and reported that expression of the C. glutamicum panD gene in E. coli yielded a strain producing pantothenate with a specific productivity of 140 ng of pantothenate per mg (dry weight) per hour. (Dusch et al. (1999) Appl. Environ. Microbiol. 65:1530-1539).

The authors Sahm and Eggeling have further identified the Corynebacterium glutamicum panB and pan C genes and have described a genetically engineered strain of C. glutamicum which overexpresses the panBC genes (Sahm and Eggeling (1999) Appl. Environ. Microbiol. 65:1973-1979). The engineered strain produces pantothenate, however, it was necessary to overexpress the genes responsible for α -ketoisovalerate production in the host organism in order that pantothenic acid production could be detected. Moreover, without the addition of \beta-alanine, no substantial amounts of pantothenate accumulated with the strain constructed.

Likewise, a method of producing D-pantothenic acid has been described that takes advantage of a sodium salicylate resistant mutant strain of E. coli which produces 30 D-pantothenic acid when cultured in the presence of β -alanine (U.S. Patent No. 5,518,906). Generation of E. coli strains resistant to α -ketoisovaleric acid and/or α ketobutyric acid, and/or α-aminobutyric acid, and/or β-hydroxyaspartic acid and/or Omethyl-threonine, in addition to salicylic acid, further increased pantothenic acid production. Moreover, transformation of a plasmid DNA carrying the panB, panC and panD genes into the salicylic acid resistant mutant strain resulted in increased pantothenate production, however, up to 20 g/L β-alanine or more was fed in the examples given. The panB-panC-panD genes are clustered on the E. coli chromosome.

20

25

30

35

Finally, a method of producing D-pantothenic acid has been described which utilizes a salicylic acid-resistant, α -ketoisovalerate-resistant, α -ketobutyrate-resistant, β -hydroxyaspartate-resistant, o-methylthreonine-resistent *E. coli* strain transformed with pantothenate biosynthesis gene-containing DNA fragments and/or branched amino acid biosynthesis gene-containing DNA fragments and cultured in the presence of β -alanine (U.S. Patent No. 5,932,457).

Pantothenate production in bacteria results from the condensation of pantoate and β-alanine and involves the pantothenate biosynthetic enzymes ketopantoate hydroxymethyltransferase (the panB gene product), ketopantoate reductase (the panE gene product), aspartate-α-decarboxylase (the panD gene product) and pantothenate synthetase (the panC gene product). Although pantothenate is biologically active as a vitamin, it is further metabolized in all cells to Coenzyme A (CoA) which participates as an acyl group carrier in the tricarboxylic acid (TCA) cycle, fatty acid metabolism and numerous other reactions of intermediary metabolism. The initial (and possibly ratecontrolling) step in the conversion of pantothenate to Coenzyme A (CoA) is phosphorylation of pantothenate by pantothenate kinase. A pantothenate kinase activity was first identified in Salmonella typhimurium by screening for temperature-sensitive mutants which synthesized CoA at permissive temperatures but excreted pantothenate at non-permissive temperatures. The mutations were mapped in the Salmonella chromosome and the genetic locus was designated coaA. The gene encodes the enzyme that catalyzes the first step in the biosynthesis of coenzyme A from pantothenate (Dunn and Snell (1979) J. Bacteriol. 140:805-808). Escherichia coli temperature sensitive mutants have also been isolated and characterized (Vallari and Rock (1987) J. Bacteriol. 169:5795-5800). These mutants (named coaA15(Ts)) are defective in the conversion of pantothenate to CoA and further exhibit a temperature-sensitive growth phenotype, indicating that pantothenate kinase activity is essential for growth. Moreover, it was noted that CoA inhibited pantothenate kinase activity to the same degree in the mutant as compared to the wild-type enzyme.

Feedback resistant *E. coli* mutants (named *coaA16(Fr)*) have also been isolated that posses a pantothenate kinase activity that is refractory to feedback inhibition by CoA (Vallari and Jackowski (1988) *J. Bacteriol.* 170:3961-3966). The mutation responsible for the reversion is, suprisingly, not genetically linked to the *coaA* gene by transduction. Additional data described therein support the view that the total cellular CoA content is controlled by both modulation of biosynthesis at the pantothenate kinase step and possibly by degradation of CoA to 4'-phosphopantetheine.

25

30

35

The wild-type E. coli coaA gene was cloned by functional complementation of E. coli temperature-sensitive mutants. The sequence of the wild-type gene was determined (Song and Jackowski (1992) J. Bacteriol. 174:6411-6417 and Flamm et al. (1988) Gene (Amst.) 74:555-558). Strains containing multiple copies of the coaA gene possessed 76-fold higher specific activity of pantothenate kinase, however, there was only a 2.7-fold increase in the steady state level of CoA (Song and Jackowski, supra). It has further been reported that the prokaryotic enzyme (encoded by coaA in E.coli and a variety of other microorganisms) is feedback inhibited by CoA both in vivo and in vitro with CoA being about five times more potent than acetyl-CoA in inhibiting the enzyme (Song and Jackowski, supra and Vallari et al., supra). Moreover, it has been reported that the panB gene product in E. coli is inhibited by CoA (Powers and Snell (1976) J. Biol. Chem. 251:3786-3793). These data further support the view that feedback inhibition of pantothenate kinase activity is a critical factor controlling intracellular CoA concentration.

Using standard search and alignment tools, coaA homologues have been identified in Hemophilus influenzae, Mycobacterium tuberculosis, Vibrio cholerae, Streptococcus pyogenes and Bacillus subtilis. By contrast, proteins with significant similarity could not be identified in eukaryotic cells including Saccharomyces cerevisiae or in mammalian expressed sequence tag (EST) databases. Using a genetic selection strategy, a cDNA encoding pantothenate kinase activity has recently been identified from Aspergillus nidulans (Calder et al. (1999) J. Biol. Chem. 274:2014-2020). The eukaryotic pantothenate kinase gene (panK) has distinct primary structure and unique regulatory properties that clearly distinguish it from its prokaryotic counterpart. A mammalian pantothenate kinase gene (mpanKla) has also been isolated which encodes a protein having homology to the A. nidulans PanK protein and to the predicted gene product of GenBankTM Accession Number 927798 identified in the S. cerevisiae genome (Rock et al. (2000) J. Biol. Chem. 275:1377-1383).

Summary of the Invention

The present invention is based, at least in part, on the discovery of key enzyme-encoding genes of the pantothenate biosynthetic pathway in *Bacillus subtilis*. In particular, the present inventors have identified the *panE* gene of *B. subtilis*. Overexpression or deregulation of the *panE* gene in *B. subtilis* results in enhanced production of the *panE* gene product, ketopantoate reductase, further resulting in increased production of pantothenate. Likewise, mutations in this gene reduce pantothenate production in *B. subtilis* >90%. The present inventors have further identified the presumptive *panBCD* operon in *B. subtilis*, overexpression or

10

15

20

25

30

35

deregulation of which results in increased pantothenate production. The present inventors have further demonstrated that overexpression or deregulation of the panD gene in B. subtilis (resulting in enhanced production of the panD gene product, aspartate- α -decarboxylase) further results in increased production of pantothenate, in particular, in combination with deregulation of genes encoding key enzymes of the isoleucine-valine (ilv) biosynthetic pathway.

Accordingly, the present invention features methods of producing pantothenate. as well as other compounds of the pantothenate biosynthetic pathway (e.g., ketopantoate, pantoate and β-alanine), termed "panto-compounds" herein, using microorganisms in which the pantothenate biosynthetic pathway and/or isoleucinevaline biosynthetic pathway has been manipulated such that pantothenate or other desired panto-compounds are produced. In one embodiment, the invention features a method of producing a panto-compound (e.g., pantothenate or pantoate) that involves culturing a microorganism which overexpresses the panE gene product, ketopantoate reductase, also referred to herein as a ketopantoate reductase-overexpressing or "KPAR-O" microorganism, under conditions such that the panto-compound (e.g., pantothenate or pantoate) is produced. In another embodiment, the present invention features a method of producing panto-compounds (e.g., pantothenate or pantoate) which includes culturing a microorganism which overexpresses at least one pantothenate biosynthetic enzyme (e.g., at least one of the panB, panC or panD gene products), preferably in a KPAR-O microorganism, under conditions such that the panto-compound (e.g., pantothenate or pantoate) is produced.

Yet another aspect of the invention features methods of producing panto-compounds which are independent of the need to feed precursors (e.g., β -alanine or aspartate and/or α -ketoisovalerate or valine). In one embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed that includes culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism having a deregulated isoleucine-valine (ilv) pathway under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed that includes culturing an A α D-O microorganism having a deregulated pantothenate (pan) pathway and a deregulated isoleucine-valine (ilv) pathway, under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of aspartate or β -alanine feed that includes culturing an A α D-O microorganism under conditions such that pantothenate is produced. In another embodiment, the invention such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of valine or α -ketoisovalerate feed that includes

10

25

30

35

culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced. In yet another embodiment, the invention features a high yield production method for producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly high yield (e.g., at a level greater than 10 g/L, 20 g/L, 30 g/L or 40g/L).

The methods of the present invention further feature microorganisms that overexpresses acetohydroxyacid synthetase or acetohydroxyacid isomeroreductase (e.g., microorganisms transformed with a vector that includes an ilvBNC nucleic acid sequence), microorganisms that overexpresses dihydroxyacid dehydratase (e.g., microorganisms transformed with a vector that includes an ilvD nucleic acid sequence), microorganisms that overexpresses aspartate-α-decarboxylase (e.g., microorganisms transformed with a vector that includes a panD nucleic acid sequence), microorganisms having a deregulated isoleucine-valine (ilv) biosynthetic pathway and microorganisms having a deregulated pantothenate biosynthetic pathway (e.g., microorganisms that overexpress any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate- α -decarboxylase, for example, microorganisms transformed with a vector comprising a panBCD nucleic acid sequence or a vector comprising a panE1 nucleic acid sequence). In one embodiment, the recombinant 20 microorganism is Gram positive (e.g., microorganisms belonging to the genus Bacillus, Cornyebacterium, Lactobacillus, Lactococci or Streptomyces). In another embodiment, the recombinant microorganism is Gram negative. Particularly preferred is a Bacillus recombinant microorganism (e.g., Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus subtilis, Bacillus pumilus, Bacillus halodurans, and the like). Recombinant vectors that contain the genes encoding Bacillus pantothenate and/or isoleucine-valine biosynthetic enzymes (e.g., B. subtilis pantothenate and/or isoleucine-valine biosynthetic enzymes) are also described.

Also featured are methods of producing β-alanine that include culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that β -alanine is produced and methods of producing β -alanine that involve contacting a composition comprising aspartate with an isolated Bacillus aspartate-adecarboxylase enzyme under conditions such that \beta-alanine is produced.

The production methods of the present invention further can include recovering the panto-compound (e.g., pantothenate or pantoate).

The present invention further features recombinant microorganisms (e.g., AaD-O microorganisms, microorganisms having a deregulated isoleucine-valine (ilv) pathway, microorganisms overexpressing at least one of ketopantoate

30

35

hydroxymethyltransferase (the panB gene product), pantothenate synthetase (the panC gene product), aspartate-α-decarboxylase (the panD gene product), ketopantoate reductase (the panE1 gene product) and microorganisms having a deregulated panBCD operon. Also featured are panB, panC, panD, panE, ilvB, ilvN, alsS, ilvC, and/or ilvD nucleic acid molecules, as well as vectors including such nucleic acid molecules and gene products encoded by such nucleic acid molecules.

The methodology of the present invention further includes, for example in addition to overexpressing at least one pantothenate biosynthetic enzyme, deleting or mutating a second pantothenate biosynthetic enzyme, said second pantothenate biosynthetic enzyme preferably being downstream of the desired product in the 10 pantothenate biosynthetic pathway. For example, mutating panC, in addition to overexpressing the panE gene product, results in even further enhanced or increased production of pantoate. Accordingly, in one embodiment, the invention features a method of producing pantoate which includes culturing a microorganism which overexpresses the panE gene product and which has a deletion in the panC gene. In another embodiment, the invention features a method of producing pantoate which includes culturing a microorganism which overexpresses the panE gene product and/or panB gene product and which has a deletion in the panC gene. Other exemplary embodiments include a method of producing ketopantoate which includes culturing a microorganism which overexpresses the panB gene product and which has a deletion in the panE gene and a method of producing β-alanine which includes culturing a microorganism which overexpresses the panD gene product and which has a deletion in the panC gene. Also included are methods of producing panto-compounds which include overexpressing at least one valine biosynthetic enzyme in a microorganism 25 which has at least one pantothenate biosynthetic enzyme deleted.

The present invention is also based at least in part, on the identification and characterization of a previously unidentified microbial pantothenate kinase gene, coaX. CoaX was first identified in Bacillus subtilis and corresponds to an open reading frame in a portion of the chromosomal DNA that includes the 5' end of the ftsH gene, and all of the ftsH gene, and pabB genes. The present inventors have demonstrated that the ftsH open reading frame encodes a novel pantothenate kinase activity, the gene being unrelated by homology to any previously known pantothenate kinase gene. The gene has been renamed ftsH as it encodes the enzyme which catalyzes the first step in the pathway from pantothenate to ftsH coaA.

Accordingly, the present invention features new and improved methods of producing pantothenate and other key compounds of the pantothenate biosynthetic pathway (e.g., panto-compounds) utilizing microorganisms having modified

10

15

20

25

30

pantothenate kinase activity. In particular, the present invention features recombinant microorganisms that contain the coaX gene or that contain a mutant coaX gene, having reduced pantothenate kinase activity. In one embodiment, the invention features such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway. In another embodiment, the invention features such recombinant microorganisms further having a deregulated isoleucine-valine (ilv) pathway. In a preferred embodiment, the microorganisms belong to the genus Bacillus (e.g., B. subtilis).

The present invention also features recombinant microorganisms (e.g., microorganisms belonging to the genus Bacillus, for example, B. subtilis) that contain the coaA gene or that contain a mutant coaA gene, optionally including a coaX-gene or mutant thereof, having reduced pantothenate kinase activity. In one embodiment, the invention features such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway or having a deregulated isoleucine-valine (ilv) pathway.

Also featured are vectors that contain isolated coaX or coaA genes as well as mutant coaX and/or coaA genes. Isolated nucleic acid molecules that contain isolated coaX genes or mutant coaX genes are featured in addition to isolated CoaX proteins and mutant CoaX proteins.

The nucleic acids, vectors and recombinant microorganisms described above are particularly useful in the methodologies of the present invention. In particular, the invention features methods of enhancing panto-compound production (e.g., ketopantoate, pantoate and or pantothenate production) that include culturing a recombinant microorganism having a mutant coaX gene under conditions such that panto-compound production is enhanced. In one embodiment, the recombinant microorganism further includes a mutant coaA gene. In another embodiment, the recombinant microorganism further includes a mutant avtA and/or mutant ilvE gene and/or mutant ansB gene and/or mutant alsD gene. Also featured are methods for identifying pantothenate modulators utilizing the recombinant microorganisms and purified CoaX proteins of the present invention.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1 is a schematic representation of the pantothenate biosynthetic pathway. Figure 2 is a schematic representation of the plasmid pAN240, containing

sequences ligated upstream of the $P_{26}panBCD$ cassette, equivalent to the integrated version in strain PA221.

Figure 3A is a schematic representation of the plasmid pAN004, containing the panBCD operon expressed from P_{26} and RBS1.

Figure 3B is a schematic representation of the plasmid pAN006, containing the panBCD operon expressed from P_{26} and RBS2.

10 Figure 4 is a schematic representation of the plasmid pAN236, containing an integratable and amplifiable P_{26} -RBS2-panE1 expression cassette.

Figure 5 is a schematic representation of the construction of plasmid pAN423.

Figure 6 is a schematic representation of the construction of plasmids pAN426 and pAN427.

Figure 7 is a schematic representation of the construction of plasmids pAN428 and pAN429.

Figure 8 is a schematic representation of the construction of plasmid pAN431.

Figure 9 is a schematic representation of the construction of plasmid pAN441.

Figure 10 is a schematic representation of the construction of plasmid pAN440.

Figure 11 is a schematic representation of the plasmid pAN251 designed to integrate a single copy of a P_{26} -panE1 cassette at the panE1 locus by double crossover.

Figure 12 is a schematic representation of the plasmid pAN267 designed to integrate a single copy of a P_{26} -ilvBNC cassette at the amyE locus.

Figure 13 is a schematic representation of the plasmid pAN257, a clone of Bacillus subtilis ilvD in a low copy vector.

Figure 14 is a schematic representation of the plasmid pAN263, designed to integrate a single copy of a P_{26} -ilvD cassette at the ilvD locus.

Figure 15 is a schematic representation of the plasmid pAN261, designed to disrupt the Bacillus subtilis ilvD gene with the cat gene.

Figure 16 is a schematic representation of the Coenzyme A biosynthetic pathway in E. coli.

Figure 17 is a schematic representation of the structure of pAN296, a plasmid designed to delete most of the B. subtilis coaA gene and substitute a chloramphenicol resistance gene.

Figure 18 is a schematic representation of the structure of the Bacillus subtilis genome in the region of the coaA gene. The scale is in base pairs and the significant open reading frames are shown by open arrows.

20

Figure 19 is a schematic representation of the plasmid pAN281, a plasmid for expressing Bacillus subtilis coaA after integration at the bpr locus.

Figure 20A-B depicts a multiple sequence alignment (MSA) of the amino acid sequences encoded by six known or predicted microbial coaA genes. SEQ ID NOs:4-6 and 1-3 correspond to the amino acid sequences of Mycobacterium leprae (SwissProtTM Accession No. Q9X795), Mycobacterium tuberculosis (SwissProtTM Accession No. O53440), Streptomyces coelicolor (SwissProtTM Accession No. O86799), Haemophilus influenzae (SwissProtTM Accession No. P44793), Escherichia coli SwissProtTM Accession No. P15044) and Bacillus subtilis (SwissProtTM Accession No. P54556), respectively. The alignment was generated using ClustalW MSA software at the GenomeNet CLUSTALW Server at the Institute for Chemical Research, Kyoto University. The following parameters were used: Pairwise Alignment, K-tuple (word) size = 1, Window size = 5, Gap Penalty = 3, Number of Top Diagonals = 5, Scoring Method = Percent; Multiple Alignment, Gap Open Penalty = 10, Gap Extension Penalty = 0.0, Weight Transition = No, Hydrophilic residues = Gly, Pro, Ser, Asn, Asp, Gln, Glu, Arg and Lys, Hydrophobic Gaps = Yes; and Scoring Matrix = BLOSUM.

Figure 21 is a schematic representation of the structure of the Bacillus subtilis genome in the region of the coaX (yacB) gene. The scale is in base pairs, the significant open reading frames are shown by open arrows and certain predicted restriction fragments are indicated by thick bars.

Figure 22 is a schematic representation of the structure of pAN341 and pAN342, two independent PCR-derived clones of B. subtilis yacB (remaned herein as coaX).

Figure 23A-D depicts a multiple sequence alignment (MSA) of the amino acid sequences encoded by fourteen known or predicted microbial coaX genes. SEQ ID NOs:9, 74, 7-8, 75, 11, 10 and 12-18 correspond to the amino acid sequences of Bacillus 25 subtilis (SwissProt™ Accession No. P37564), Clostridium acetobulyticum (WIT™ Accession No. RCA03301, Argonne National Laboratories), Streptomyces coelicolor (PIR™ Accession No. T36391), Mycobacterium tuberculosis (SwissProt™ Accession No. O06282), Rhodobacter capsulatus (WIT™ Accession No. RRC02473), Desulfovibrio vulgaris (DBJTM Accession No. BAA21476.1), Deinococcus radiodurans (SwissProt[™] Accession No. Q9RX54), Thermotoga maritima (GenBank[™] Accession No. AAD35964.1), Treponema pallidum (SwissProt™ Accession No. O83446), Borrelia burgdorferi (SwissProt™ Accession No.O51477), Aquifex aeolicus (SwissProt[™] Accession No. O67753), Synechocystis sp. (SwissProt[™] Accession No. P74045), Helicobacter pylori (SwissProt™ Accession No. O25533), and Bordetella pertussis (SwissProtTM Accession No. Q45338), respectively. The alignment was generated using Clustal W MSA software at the GenomeNet CLUSTAL W Server at the

Institute for Chemical Research, Kyoto University. The following parameters were used: Pairwise Alignment, K-tuple (word) size = 1, Window size = 5, Gap Penalty = 3, Number of Top Diagonals = 5, Scoring Method = Percent; Multiple Alignment, Gap Open Penalty = 10, Gap Extension Penalty = 0.0, Weight Transition = No, Hydrophilic residues = Gly, Pro, Ser, Asn, Asp, Gln, Glu, Arg and Lys, Hydrophobic Gaps = Yes; and Scoring Matrix = BLOSUM.

Figure 24 depicts a multiple sequence alignment of a portion of the protein sequences of the coaA gene products from the following microorganisms: Bacillus subtilis, Escherichia coli, Haemophilus influenzae, Mycobacterium leprae,

Mycobacterium tuberculosis, and Streptomyces coelicolor. The residues that are mutated in E. coli coaA15(Ts) and B. subtilis coaA282A are indicated below and above the alignment, respectively. The portions correspond to amino acid residues 168-187 of SEQ ID NO:3, 167-186 of SEQ ID NO:2, 165-184 of SEQ ID NO:1, 169-188 of SEQ ID NO:4, 169-188 of SEQ ID NO:5 and 179-198 of SEQ ID NO:6, respectively.

Figure 25 is a schematic representation of the structure of pAN294, a plasmid for integrating mutagenized B. subtilis coaA at its native locus.

Figure 26 is a schematic representation of the structure of pAN336, a plasmid designed to delete B. subtilis coaX from its chromosomal locus and replace it with a kanamycin resistence gene.

20

25

30

35

Detailed Description of the Invention

The present invention features new and improved methods of producing pantothenate and other key compounds of the pantothenate biosynthetic pathway (referred to herein as "panto-compounds", for example, pantothenate, ketopantoate, pantoate and β -alanine) using microorganisms in which the pantothenate biosynthetic pathway has been manipulated such that pantothenate or other desired panto-compounds are produced.

The new and improved methodologies of the present invention include methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one enzyme of the pantothenate biosynthetic pathway manipulated such that pantothenate or other desired panto-compounds are produced (e.g., produced at an increased level). For example, the invention features methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase or aspartate- α -decarboxylase manipulated such that pantothenate or other desired panto-compounds are produced. The methodologies of the present invention also include methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one

20

30

valine-isoleucine biosynthetic enzyme, described herein, manipulated such that pantothenate or other desired panto-compounds are produced. For example, the invention features methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one of acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase or dihydroxyacid dehydratase manipulated such that pantothenate or other desired panto-compounds are produced.

The invention also features methods of producing panto-compounds that involve culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism under conditions such that the panto-compound is produced. The invention also features methods of producing pantothenate in a manner independent of precursor feed that involve culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that pantothenate is produced. Also featured are β -alanine independent high yield pantothenate production methods as well as methods of producing β -alanine. The present invention also features methods for enhancing production of panto-compounds that involve culturing pantothenate kinase mutants. In particular, the present invention features new and improved methods of producing pantothenate and other key compounds of the pantothenate biosynthetic pathway (e.g., panto-compounds) utilizing microorganisms having modified pantothenate kinase activity, for example, microorganisms that include the coaX gene or that include a mutant coaX gene, having reduced pantothenate kinase activity.

In order that the present invention may be more readily understood, certain terms are first defined herein.

The term "pantothenate biosynthetic pathway" includes the biosynthetic pathway involving pantothenate biosynthetic enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of pantothenate. The term "pantothenate biosynthetic pathway" includes the biosynthetic pathway leading to the synthesis of pantothenate in a microorganisms (e.g., in vivo) as well as the biosynthetic pathway leading to the synthesis of pantothenate in vitro. Figure 1 includes a schematic representation of the pantothenate biosynthetic pathway. Pantothenate biosynthetic enzymes are depicted in bold and their corresponding genes indicated in italics.

The term "pantothenate biosynthetic enzyme" includes any enzyme utilized in the formation of a compound (e.g., intermediate or product) of the pantothenate biosynthetic pathway. According to Figure 1, synthesis of pantoate from α -ketoisovalerate (α -KIV) proceeds via the intermediate, ketopantoate. Formation of ketopantoate is catalyzed by the pantothenate biosynthetic enzyme ketopantoate

hydroxymethyltransferase (the panB gene product). Formation of pantoate is catalyzed by the pantothenate biosynthetic enzyme ketopantoate reductase (the panE gene product). Synthesis of β-alanine from aspartate is catalyzed by the pantothenate biosynthetic enzyme aspartate-α-decarboxylase (the panD gene product). Formation of pantothenate from pantoate and β -alanine (e.g., condensation) is catalyzed by the pantothenate biosynthetic enzyme pantothenate synthetase (the panC gene product).

The term "isoleucine-valine biosynthetic pathway" includes the biosynthetic pathway involving isoleucine-valine biosynthetic enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of conversion of pyruvate to valine or isoleucine. The term "isoleucine-valine biosynthetic pathway" includes the biosynthetic pathway leading to the synthesis of valine or isoleucine in a microorganisms (e.g., in vivo) as well as the biosynthetic pathway leading to the synthesis of valine or isoleucine in vitro. Figure 1 includes a 15 schematic representation of the isoleucine-valine biosynthetic pathway. Isoleucinevaline biosynthetic enzymes are depicted in bold italics and their corresponding genes indicated in italics

10

20

25

30

35

The term "isoleucine-valine biosynthetic enzyme" includes any enzyme utilized in the formation of a compound (e.g., intermediate or product) of the isoleucine-valine biosynthetic pathway. According to Figure 1, synthesis of valine from pyruvate proceeds via the intermediates, acetolactate, α,β -dihydroxyisovalerate (α,β -DHIV) and α -ketoisovalerate (α -KIV). Formation of acetolactate from pyruvate is catalyzed by the isoleucine-valine biosynthetic enzyme acetohydroxyacid synthetase (the ilvBN gene product, or alternatively, the alsS gene product). Formation of α,β -DHIV from acetolactate is catalyzed by the isoleucine-valine biosynthetic enzyme acetohydroxyacidisomero reductase (the *ilvC* gene product). Synthesis of α -KIV from α,β-DHIV is catalyzed by the isoleucine-valine biosynthetic enzyme dihydroxyacid dehydratase (the ilvD gene product). Moreover, valine and isoleucine can be interconverted by branched chain amino acid transaminases.

As used herein, each of ketopantoate, pantoate, \beta-alanine and pantothenate are "panto-compounds". The term "panto-compound" includes a compound (e.g., a substrate, intermediate or product) in the pantothenate biosynthetic pathway which is downstream from a particular pantothenate biosynthetic enzyme. In one example, a panto-compound is downstream of the pantothenate biosynthetic enzyme ketopantoate hydroxymethyltransferase (the panB gene product) and can include ketopantoate. pantoate and/or pantothenate. In another example, a panto-compound is downstream of the pantothenate biosynthetic enzyme ketopantoate reductase (the panE gene product)

20

25

30

012177282 L

DISCOUNT -WA

and can include pantoate and/or pantothenate. In yet another example, a panto-compound is downstream of the pantothenate biosynthetic enzyme pantothenate synthetase (the panC gene product) and can include pantothenate. In yet another example, a panto-compound is downstream of the pantothenate biosynthetic enzyme aspartate- α -decarboxylase (the panD gene product) and can include β -alanine and/or pantothenate.

Preferred panto-compounds include pantothenate and pantoate. The term "pantothenate" includes the free acid form of pantothenate, also referred to as "pantothenic acid" as well as any salt thereof (e.g., derived by replacing the acidic hydrogen of pantothenate or pantothenic acid with a cation, for example, calcium, sodium, potassium, ammonium), also referred to as a "pantothenate salt". The term "panto-compound" also includes alcohol derivatives of pantothenate. Preferred pantothenate salts are calcium pantothenate or sodium pantothenate. A preferred alcohol derivative is pantothenol. Pantothenate salts and/or alcohols of the present invention include salts and/or alcohols prepared via conventional methods from the free acids described herein. In another embodiment, calcium pantothenate is synthesized directly by a microorganism of the present invention. A pantothenate salt of the present invention can likewise be converted to a free acid form of pantothenate or pantothenic acid by conventional methodology.

The term "pantoate" includes the free acid form of pantoate, also referred to as "pantoic acid" as well as any salt thereof (e.g., derived by replacing the acidic hydrogen of pantoate or pantoic acid with a cation, for example, calcium, sodium, potassium, ammonium), also referred to as a "pantoate salt". Preferred pantoate salts are calcium pantoate or sodium pantoate. Pantoate salts of the present invention include salts prepared via conventional methods from the free acids described herein. A pantoate salt of the present invention can likewise be converted to a free acid form of pantoate or pantoic acid by conventional methodology. Moreover, a free acid form of pantoate or pantoic acid can be converted to pantolactone by conventional methodology.

The term "CoA biosynthetic pathway" includes the biosynthetic pathway involving CoA biosynthetic enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of CoA from pantothenate. A schematic representation of the CoA biosynthetic pathway in E. coli is set forth as Figure 16. (The pathway depicted is also presumed to be that utilized by other microorganisms.) The term "CoA biosynthetic pathway" includes the biosynthetic pathway leading to the synthesis of CoA in microorganisms (e.g., in vivo) as well as the biosynthetic pathway leading to the synthesis of CoA in vitro. The term "Coenzyme A

30

or CoA biosynthetic enzyme" includes any enzyme utilized in the formation of a compound (e.g., intermediate or product) of the CoA biosynthetic pathway, for example, the coaA, panK or coaX gene product which catalyzes the phosphorylation of pantothenate to form 4'-phosphopantothenate, or the coaD gene product which catalyzes the conversion of 4'-phosphopantetheine to dephosphocoenzyme A.

I. Recombinant Microorganisms and Methods for Culturing Microorganisms Such That Panto-Compounds are Produced

The methodologies of the present invention feature microorganisms, e.g., recombinant microorganisms, preferably including vectors or genes (e.g., wild-type 10 and/or mutated genes) as described herein and/or cultured in a manner which results in the production of a desired product (e.g. a panto-compound or panto-compounds). The term "recombinant" microorganism includes a microorganism (e.g., bacteria, yeast cell, fungal cell, etc.) which has been genetically altered, modified or engineered (e.g., genetically engineered) such that it exhibits an altered, modified or different genotype 15 and/or phenotype (e.g., when the genetic modification affects coding nucleic acid sequences of the microorganism) as compared to the naturally-occurring microorganism from which it was derived. Preferably, a "recombinant" microorganism of the present invention has been genetically engineered such that it overexpresses at least one bacterial gene or gene product (e.g., a pantothenate or isoleucine-valine biosynthetic 20 enzyme encoding-gene) as described herein, preferably a biosynthetic enzyme encodinggene included within a recombinant vector as described herein and/or a biosynthetic enzyme expressed from a recombinant vector. The ordinary skilled will appreciate that a microorganism expressing or overexpressing a gene product produces or overproduces the gene product as a result of expression or overexpression of nucleic acid sequences 25 and/or genes encoding the gene product.

The term "manipulated microorganism" includes a microorganism that has been engineered (e.g., genetically engineered) or modified such that the microorganism has at least one enzyme of the pantothenate biosynthetic pathway and/or at least one enzyme of the isoleucine-valine biosynthetic pathway modified such that pantothenate or other desired panto-compounds are produced. Modification or engineering of such microorganisms can be according to any methodology described herein including, but not limited to, deregulation of a biosynthetic pathway and/or overexpression of at least one biosynthetic enzyme. A "manipulated" enzyme (e.g., a "manipulated" biosynthetic enzyme) includes an enzyme, the expression or production of which has been altered or modified such that at least one upstream or downstream precursor, substrate or product

20

of the enzyme is altered or modified, for example, as compared to a corresponding wildtype or naturally occurring enzyme.

The term "overexpressed" or "overexpression" includes expression of a gene product (e.g., a pantothenate biosynthetic enzyme or isoleucine-valine biosynthetic 5 enzyme) at a level greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. In one embodiment, the microorganism can be genetically manipulated (e.g., genetically engineered) to overexpress a level of gene product greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. Genetic manipulation can include, but is not limited to, altering or modifying regulatory sequences or sites associated with expression of a particular gene (e.g., by adding strong promoters, inducible promoters or multiple promoters or by removing regulatory sequences such that expression is constitutive), modifying the chromosomal location of a particular gene, altering nucleic acid sequences adjacent to a particular gene such as a ribosome binding site or transcription terminator, increasing the copy number of a particular gene, modifying proteins (e.g., regulatory proteins, suppressors, enhancers, transcriptional activators and the like) involved in transcription of a particular gene and/or translation of a particular gene product, or any other conventional means of deregulating expression of a particular gene routine in the art (including but not limited to use of antisense nucleic acid molecules, for example, to block expression of repressor proteins).

In another embodiment, the microorganism can be physically or environmentally manipulated to overexpress a level of gene product greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. For example, a microorganism can be treated with or cultured in the presence of an agent known or suspected to increase transcription of a particular gene and/or translation of a particular gene product such that transcription and/or translation are enhanced or increased. Alternatively, a microorganism can be cultured at a temperature selected to increase transcription of a particular gene and/or translation of a particular gene product such that transcription and/or translation are enhanced or increased.

The term "deregulated" or "deregulation" includes the alteration or modification of at least one gene in a microorganism that encodes an enzyme in a biosynthetic pathway, such that the level or activity of the biosynthetic enzyme in the microorganism is altered or modified. Preferably, at least one gene that encodes an enzyme in a biosynthetic pathway is altered or modified such that the gene product is enhanced or increased. The phrase "deregulated pathway" can also include a biosynthetic pathway in which more than one gene that encodes an enzyme in a biosynthetic pathway is altered or modified such that the level or activity of more than one biosynthetic enzyme is altered or modified. The ability to "deregulate" a pathway (e.g., to simultaneously deregulate more than one gene in a given biosynthetic pathway) in a microorganism arises from the particular phenomenon of microorganisms in which more than one enzyme (e.g., two or three biosynthetic enzymes) are encoded by genes occurring adjacent to one another on a contiguous piece of genetic material termed an "operon".

The term "operon" includes a coordinated unit of gene expression that contains a promoter and possibly a regulatory element associated with one or more, preferably at least two, structural genes (e.g., genes encoding enzymes, for example, biosynthetic enzymes). Expression of the structural genes can be coordinately regulated, for example, by regulatory proteins binding to the regulatory element or by anti-termination of transcription. The structural genes can be transcribed to give a single mRNA that encodes all of the structural proteins. Due to the coordinated regulation of genes 15 included in an operon, alteration or modification of the single promoter and/or regulatory element can result in alteration or modification of each gene product encoded by the operon. Alteration or modification of the regulatory element can include, but is not limited to removing the endogenous promoter and/or regulatory element(s), adding strong promoters, inducible promoters or multiple promoters or removing regulatory sequences such that expression of the gene products is modified, modifying the 20 chromosomal location of the operon, altering nucleic acid sequences adjacent to the operon or within the operon such as a ribosome binding site, increasing the copy number of the operon, modifying proteins (e.g., regulatory proteins, suppressors, enhancers, transcriptional activators and the like) involved in transcription of the operon and/or translation of the gene products of the operon, or any other conventional means of deregulating expression of genes routine in the art (including but not limited to use of antisense nucleic acid molecules, for example, to block expression of repressor proteins). Deregulation can also involve altering the coding region of one or more genes to yield, for example, an enzyme that is feedback resistant or has a higher or lower 30 specific activity.

A particularly preferred "recombinant" microorganism of the present invention has been genetically engineered to overexpress a bacterially-derived gene or gene product. The term "bacterially-derived" or "derived-from", for example bacteria, includes a gene which is naturally found in bacteria or a gene product (e.g., ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase, aspartate- α -decarboxylate, acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase or

dihydroxyacid dehydratase) which is encoded by a bacterial gene (e.g., encoded by panB, panE, panC, panD, ilvB, ilvN, alsS, ilvC, or ilvD).

The methodologies of the present invention feature recombinant microorganisms which overexpress at least one of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase or aspartate-a-decarboxylase. A particularly preferred recombinant microorganism of the present invention has been genetically engineered to overexpress a Bacillus (e.g., Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus, etc.) biosynthetic enzyme (e.g., has been engineered to overexpress at least one of B. subtilis ketopantoate reductase (the panE gene product) (e.g., ketopantoate reductase having the amino acid sequence of SEQ ID NO:30 or encoded by the nucleic acid sequence of SEQ ID NO:29), B. subtilis ketopantoate hydroxymethyltransferase (the panB gene product) (e.g., ketopantoate hydroxymethyltransferase having the amino acid sequence of SEQ ID NO:24 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:23), B. subtilis pantothenate synthetase (the panC-gene product) (e.g., 15 pantothenate synthetase having the amino acid sequence of SEQ ID NO:26 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:25) and/or B. subtilis aspartate-α-decarboxylase (the panD gene product) (e.g., aspartate-αdecarboxylase having the amino acid sequence of SEQ ID NO:28 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:27). 20

In an exemplary embodiment, the invention features a microorganism (e.g., a KPAR-O microorganism) that has been transformed with a vector comprising a panE nucleic acid sequence (e.g., a panE nucleic acid sequence as set forth in SEQ ID NO:29). In another embodiment, the invention features a microorganism that has been transformed with a vector comprising a panB nucleic acid sequence (e.g., a panB nucleic acid sequence as set forth in SEQ ID NO:23), a vector comprising a panC nucleic acid sequence (e.g., a panC nucleic acid sequence as set forth in SEQ ID NO:25) or a vector comprising a panD nucleic acid sequence (e.g., a panD nucleic acid sequence as set forth in SEQ ID NO:27). In yet another embodiment, the invention features a microorganism having a deregulated panBCD operon (e.g., SEQ ID NO:59).

Other preferred "recombinant" microorganisms of the present invention have a deregulated isoleucine-valine (ilv) pathway. The phrase "microorganism having a deregulated isoleucine-valine (ilv) pathway" includes a microorganism having an alteration or modification in at least one gene encoding an enzyme of the isoleucine-valine (ilv) pathway or having an alteration or modification in an operon including more than one gene encoding an enzyme of the isoleucine-valine (ilv) pathway. A preferred "microorganism having a deregulated isoleucine-valine (ilv) pathway" has been

30

15

20

25

30

35

genetically engineered to overexpress a Bacillus (e.g., B. subtilis) ilv biosynthetic enzyme (e.g., has been engineered to overexpress at least one of acetohydroxyacid synthetase (the ilvBN gene products or the alsS gene product) (e.g., acetohydroxyacid synthetase having subunits having the amino acid sequences of SEQ ID NO:32 and SEQ ID NO:34 or encoded by nucleic acid molecules having the nucleotide sequence of SEO ID NO:31 and SEQ ID NO:33 or the nucleotide sequence of SEQ ID NO:58 from nucleotides 1-2246 or acetohydroxyacid synthetase encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:86), acetohydroxyacid isomeroreductase (the ilvC gene product) (e.g., acetohydroxyacid isomeroreductase having the amino acid sequence of SEQ ID NO:36 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:35), dihydroxyacid dehydratase (the ilvD gene product) (e.g., dihydroxyacid dehydratase having the amino acid sequence of SEQ ID NO:38 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:37), and/or has been transformed with a vector comprising an ilvBNC nucleic acid sequence (SEQ ID NO:58, coding regions from nucleotides 1-1725, 1722-2246 and 2263-3291) and/or an ilvD nucleic acid sequence (SEQ ID NO:37).

In another preferred embodiment, a recombinant microorganism is designed or engineered such that a mutant CoaA and/or CoaX biosynthetic enzyme is expressed and at least one pantothenate biosynthetic enzyme and/or at least one isoleucine-valine biosynthetic enzyme is overexpressed or deregulated.

In another preferred embodiment, a microorganism of the present invention overexpresses or is mutated for a gene or biosynthetic enzyme (e.g., a CoA biosynthetic enzyme, pantothenate biosynthetic enzyme or isoleucine-valine biosynthetic enzyme) which is bacterially-derived. The term "bacterially-derived" or "derived-from", for example bacteria, includes a gene product (e.g., ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase, aspartate-α-decarboxylate, acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase, dihydroxyacid dehydratase or pantothenate kinase) which is encoded by a bacterial gene (e.g., panB, panE, panC, panD, ilvBN (or alsS), ilvC, ilvD, or encoded by coaA or coaX).

Still other preferred recombinant microorganisms of the present invention are mutant microorganisms. As used herein, the term "mutant microorganism" includes a recombinant microorganism that has been genetically engineered to express a mutated gene or protein that is normally or naturally expressed by the microorganism.

Preferably, a mutant microorganism expresses a mutated gene or protein such that the microorganism exhibits an altered, modified or different phenotype (e.g., has been engineered to express a mutated CoaA biosynthetic enzyme, for example, pantothenate kinase). In one embodiment, a mutant microorganism is designed or engineered such

that it includes a mutant coaX gene, as defined herein. In another embodiment, a recombinant microorganism is designed or engineered such that it includes a mutant coaA gene, as defined herein. In another embodiment, a mutant microorganism is designed or engineered such that a coaX gene has been deleted (i.e., the protein encoded by the coaX gene is not produced). In another embodiment, a mutant microorganism is designed or engineered such that a coaA gene has been deleted (i.e., the protein encoded by the coaA gene is not produced). Preferably, a mutant microorganism has a mutant coaX gene or a mutant coaA gene, or has been engineered to have a coaX gene and/or coaA deleted, such that that the mutant microorganism encodes a "reduced pantothenate kinase activity". In the context of a whole microorganism, a "reduced pantothenate kinase activity" can be determined by measuring or assaying for a decrease in an intermediate or product of the CoA biosynthetic pathway, for example, measuring or assaying for 4'-phosphopantothenate, 4'-phosphopantothenylcysteine, 4'phosphopantetheine, dephosphocoenzyme A, Coenzyme A, apo-acyl carrier protein (apo-ACP) or holo-acyl carrier protein (ACP) in the microorganism (e.g., in a lysate isolated or derived from the microorganism) or in the medium in which the microorganism is cultured (see e.g., Figure 16). Alternatively, a "reduced pantothenate kinase activity" can be determined by measuring or assaying for decreased growth of the microorganism. Alternatively, a "reduced pantothenate kinase activity" can be determined by measuring or assaying for an increase in a panto-compound (e.g., pantothenate) in the microorganism or surrounding media, as panto-compounds lie upstream of the CoA biosynthetic pathway, the first step of which is catalyzed by pantothenate kinase. The invention also features recombinant microorganisms that, in addition to having reduced pantothenate kinase activity (e.g., expressing mutant-coaA and/or mutant coaX genes) have a deregulated pantothenate biosynthesis pathway and/or 25 a deregulated isoleucine-valine (ilv) biosynthetic pathway.

In one embodiment, a recombinant microorganism of the present invention is a Gram positive organism (e.g., a microorganism which retains basic dye, for example, crystal violet, due to the presence of a Gram-positive wall surrounding the microorganism). In a preferred embodiment, the recombinant microorganism is a microorganism belonging to a genus selected from the group consisting of Bacillus, Cornyebacterium, Lactobacillus, Lactococci and Streptomyces. In a more preferred embodiment, the recombinant microorganism is of the genus Bacillus. In another preferred embodiment, the recombinant microorganism is selected from the group consisting of Bacillus subtilis, Bacillus lentimorbus, Bacillus lentus, Bacillus firmus, Bacillus pantothenticus, Bacillus amyloliquefaciens, Bacillus cereus, Bacillus circulans, Bacillus coagulans, Bacillus licheniformis, Bacillus megaterium, Bacillus pumilus,

30

15

20

25

30

35

Bacillus thuringiensis, and other Group 1 Bacillus species, for example, as characterized by 16S rRNA type (Priest (1993) in Bacillus subtilis and Other Gram-Positive Bacteria eds. Sonenshein et al., ASM, Washington, D.C., p. 6). In another preferred embodiment, the recombinant microorganism is Bacillus brevis or Bacillus stearothermophilus. In another preferred embodiment, the recombinant microorganism is selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus.

In another embodiment, the recombinant microorganism is a Gram negative (excludes basic dye) organism. In a preferred embodiment, the recombinant microorganism is a microorganism belonging to a genus selected from the group consisting of Salmonella, Escherichia, Klebsiella, Serratia, and Proteus. In a more preferred embodiment, the recombinant microorganism is of the genus Escherichia. In an even more preferred embodiment, the recombinant microorganism is Escherichia coli. In another embodiment, the recombinant microorganism is Saccharomyces (e.g., S. cerevisiae).

An important aspect of the present invention involves culturing the recombinant microorganisms described herein, such that a desired compound (e.g., a desired pantocompound) is produced. The term "culturing" includes maintaining and/or growing a living microorganism of the present invention (e.g., maintaining and/or growing a culture or strain). In one embodiment, a microorganism of the invention is cultured in liquid media. In another embodiment, a microorganism of the invention is cultured in solid media or semi-solid media. In a preferred embodiment, a microorganism of the invention is cultured in media (e.g., a sterile, liquid media) comprising nutrients essential or beneficial to the maintenance and/or growth of the microorganism (e.g., carbon sources or carbon substrate, for example complex carbohydrates such as bean or grain meal, starches, sugars, sugar alcohols, hydrocarbons, oils, fats, fatty acids, organic acids and alcohols; nitrogen sources, for example, vegetable proteins, peptones, peptides and amino acids derived from grains, beans and tubers, proteins, peptides and amino acids derived form animal sources such as meat, milk and animal byproducts such as peptones, meat extracts and casein hydrolysates; inorganic nitrogen sources such as urea, ammonium sulfate, ammonium chloride, ammonium nitrate and ammonium phosphate; phosphorus sources, for example, phosphoric acid, sodium and potassium salts thereof; trace elements, for example, magnesium, iron, manganese, calcium, copper, zinc, boron, molybdenum, and/or cobalt salts; as well as growth factors such as amino acids, vitamins, growth promoters and the like).

25

Preferably, microorganisms of the present invention are cultured under controlled pH. The term "controlled pH" includes any pH which results in production of the desired product (e.g., a panto-compound). In one embodiment, microorganisms are cultured at a pH of about 7. In another embodiment, microorganisms are cultured at a pH of between 6.0 and 8.5. The desired pH may be maintained by any number of methods known to those skilled in the art.

Also preferably, microorganisms of the present invention are cultured under controlled aeration. The term "controlled aeration" includes sufficient aeration (e.g., oxygen) to result in production of the desired product (e.g., panto-compound). In one embodiment, aeration is controlled by regulating oxygen levels in the culture, for example, by regulating the amount of oxygen dissolved in culture media. Preferably, aeration of the culture is controlled by agitating the culture. Agitation may be provided by a propeller or similar mechanical agitation equipment, by revolving or shaking the growth vessel (e.g., fermentor) or by various pumping equipment. Aeration may be further controlled by the passage of sterile air or oxygen through the medium (e.g., through the fermentation mixture). Also preferably, microorganisms of the present invention are cultured without excess foaming (e.g., via addition of antifoaming agents).

Moreover, microorganisms of the present invention can be cultured under controlled temperatures. The term "controlled temperature" includes any temperature which results in production of the desired product (e.g., a panto-compound). In one embodiment, controlled temperatures include temperatures between 15°C and 95°C. In another embodiment, controlled temperatures include temperatures between 15°C and 70°C. Preferred temperatures are between 20°C and 55°C, more preferably between 30°C and 45°C or between 30°C and 50°C.

Microorganisms can be cultured (e.g., maintained and/or grown) in liquid media and preferably are cultured, either continuously or intermittently, by conventional culturing methods such as standing culture, test tube culture, shaking culture (e.g., rotary shaking culture, shake flask culture, etc.), aeration spinner culture, or fermentation. In a preferred embodiment, the microorganisms are cultured in shake flasks. In a more preferred embodiment, the microorganisms are cultured in a fermentor (e.g., a fermentation process). Fermentation processes of the present invention include, but are not limited to, batch, fed-batch and continuous methods of fermentation. The phrase "batch process" or "batch fermentation" refers to a closed system in which the composition of media, nutrients, supplemental additives and the like is set at the beginning of the fermentation and not subject to alteration during the fermentation, however, attempts may be made to control such factors as pH and oxygen concentration to prevent excess media acidification and/or microorganism death. The phrase "fed-

20

25

30

35

010177080 1 5

batch process" or "fed-batch" fermentation refers to a batch fermentation with the exception that one or more substrates or supplements are added (e.g., added in increments or continuously) as the fermentation progresses. The phrase "continuous process" or "continuous fermentation" refers to a system in which a defined fermentation media is added continuously to a fermentor and an equal amount of used or "conditioned" media is simultaneously removed, preferably for recovery of the desired product (e.g., panto-compound). A variety of such processes have been developed and are well-known in the art.

The phrase "culturing under conditions such that a desired compound (e.g., a panto-compound, for example, pantothenate) is produced" includes maintaining and/or growing microorganisms under conditions (e.g., temperature, pressure, pH, duration, etc.) appropriate or sufficient to obtain production of the desired compound or to obtain desired yields of the particular compound being produced. For example, culturing is continued for a time sufficient to produce the desired amount of a panto-compound (e.g., 15 pantothenate, pantoate or β-alanine). Preferably, culturing is continued for a time sufficient to substantially reach maximal production of the panto-compound. In one embodiment, culturing is continued for about 12 to 24 hours. In another embodiment, culturing is continued for about 24 to 36 hours, 36 to 48 hours, 48 to 72 hours, 72 to 96 hours, 96 to 120 hours, 120 to 144 hours, or greater than 144 hours. In another embodiment, culturing is continued for a time sufficient to reach production yields of panto-compound, for example, cells are cultured such that at least about 15 to 20 g/L of panto-compound are produced, at least about 20 to 25 g/L panto-compound are produced, at least about 25 to 30 g/L panto-compound are produced, at least about 30 to 35 g/L panto-compound are produced, at least about 35 to 40 g/L panto-compound are produced (e.g., at least about 37 g/L panto-compound) or at least about 40 to 50 g/L panto compound are produced. In yet another embodiment, microorganisms are cultured under conditions such that a preferred yield of panto-compound, for example, a yield within a range set forth above, is produced in about 24 hours, in about 36 hours, in about 48 hours, in about 72 hours, or in about 96 hours.

The methodology of the present invention can further include a step of recovering a desired compound (e.g., a panto-compound). The term "recovering" a desired compound (e.g., a panto-compound) includes extracting, harvesting, isolating or purifying the compound from culture media. Recovering the compound can be performed according to any conventional isolation or purification methodology known in the art including, but not limited to, treatment with a conventional resin (e.g., anion or cation exchange resin, non-ionic adsorption resin, etc.), treatment with a conventional adsorbent (e.g., activated charcoal, silicic acid, silica gel, cellulose, alumina, etc.),

alteration of pH, solvent extraction (e.g., with a conventional solvent such as an alcohol, ethyl acetate, hexane and the like), dialysis, filtration, concentration, crystallization, recrystallization, pH adjustment, lyophilization and the like. For example, a compound (e.g., a panto-compound) can be recovered from culture media by first removing the microorganisms from the culture. Media is then passed through or over a cation exchange resin to remove unwanted cations and then through or over an anion exchange resin to remove unwanted inorganic anions and organic acids having stronger acidities than the panto-compound of interest (e.g., pantothenate). The resulting panto-compound (e.g., pantothenate) can subsequently be converted to a pantothenate salt (e.g., calcium pantothenate) as described herein.

Preferably, a desired compound of the present invention is "extracted", "isolated" or "purified" such that the resulting preparation is substantially free of other components (e.g., free of media components and/or fermentation byproducts). The language "substantially free of other components" includes preparations of desired compound in which the compound is separated (e.g., purified or partially purified) from media components or fermentation byproducts of the culture from which it is produced. In one embodiment, the preparation has greater than about 80% (by dry weight) of the desired compound (e.g., less than about 20% of other media components or fermentation byproducts), more preferably greater than about 90% of the desired compound (e.g., less than about 10% of other media components or fermentation byproducts), still more preferably greater than about 95% of the desired compound (e.g., less than about 5% of other media components or fermentation byproducts), and most preferably greater than about 98-99% desired compound (e.g., less than about 1-2% other media components or fermentation byproducts). When the desired compound is a panto-compound that has been derivatized to a salt (e.g. a pantothenate salt or pantoate salt), the panto-compound is preferably further free (e.g., substantially free) of chemical contaminants associated with the formation of the salt. When the desired compound is a panto-compound that has been derivatized to an alcohol, the panto-compound is preferably further free (e.g., substantially free) of chemical contaminants associated with the formation of the alcohol.

In an alternative embodiment, the desired panto-compound is not purified from the microorganism, for example, when the microorganism is biologically non-hazardous (e.g., safe). For example, the entire culture (or culture supernatant) can be used as a source of product (e.g., crude product). In one embodiment, the culture (or culture supernatant) supernatant is used without modification. In another embodiment, the culture (or culture supernatant) is concentrated. In yet another embodiment, the culture (or culture supernatant) is dried or lyophilized.

30

10

20

Panto-Compound Production Methodologies Featuring Ketopantoate Reductase-Overexpressing Microorganisms

One aspect of the invention features methods of producing a panto-compounds that involve culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism under conditions such that the panto-compound is produced. The term "ketopantoate reductase-overexpressing (KPAR-O) microorganism" includes a microorganism which has been manipulated such that ketopantoate reductase is overexpressed (e.g., a B. subtilis ketopantoate reductase protein having the amino acid sequence of SEQ ID NO:30) and/or has been transformed with a vector comprising a panE1 nucleic acid sequence (e.g., a B. subtilis panE1 nucleic acid sequence as set forth in SEQ ID NO:29). In one embodiment, the panto-compound is pantothenate. In another embodiment, the panto-compound is pantoate. In another embodiment, the ketopantoate reductase is bacterial-derived. In another embodiemnt, the ketopantoate reductase is derived from Bacillus (e.g., is derived from Bacillus subtilis). In yet 15 another embodiment, the KPAR-O microorganism is Gram positive. In yet another embodiment, the KPAR-O microorganism is a microorganism belonging to a genus selected from the group consisting of Bacillus, Cornyebacterium, Lactobacillus, Lactococci and Streptomyces. In a preferred embodiemnt, the KPAR-O microorganism is of the genus Bacillus. In a more preferred embodiment, the KPAR-O microorganism is selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis and Bacillus pumilus. In a particularly preferred embodiemnt, the KPAR-O microorganism is Bacillus subtilis.

In still other embodiments, the KPAR-O microorganism further overexpresses at least one pantothenate biosynthetic enzyme in addition to ketopantoate reductase. In an exemplary embodiment, the KPAR-O microorganism further overexpresses at least one of ketopantoate hydroxymethyltransferase, pantothenate synthetase and aspartate-adecarboxylase. Also featured are methods of producing panto-compounds, for example, methods that involve culturing a KPAR-O microorganism, which further include the step of recovering the panto-compound.

30

35

Methods of Producing Panto-Compounds Independent of Precursor Feed *III*. Requirements

Depending on the biosynthetic enzyme or combination of biosynthetic enzymes manipulated, it may be desirable or necessary to provide (e.g., feed) microorganisms of the present invention at least one pantothenate biosynthetic precursor such that pantothenate or other desired panto-compounds are produced. The term "pantothenate biosynthetic precursor" or "precursor" includes an agent or compound which, when

20

25

provided to, brought into contact with, or included in the culture medium of a microorganism, serves to enhance or increase pantothenate biosynthesis. In one embodiment, the pantothenate biosynthetic precursor or precursor is aspartate. In another embodiment, the pantothenate biosynthetic precursor or precursor is \(\beta-alanine.

The amount of aspartate or β-alanine added is preferably an amount that results in a concentration in the culture medium sufficient to enhance productivity of the microorganism (e.g., a concentration sufficient to enhance production of a pantocompound, for example, β-alanine, ketopantoate, pantoate or pantothenate).

Pantothenate biosynthetic precursors of the present invention can be added in the form of a concentrated solution or suspension (e.g., in a suitable solvent such as water or buffer) or in the form of a solid (e.g., in the form of a powder). Moreover, pantothenate biosynthetic precursors of the present invention can be added as a single aliquot, continuously or intermittently over a given period of time.

In yet another embodiment, the pantothenate biosynthetic precursor is valine, see e.g., Example III. In yet another embodiment, the pantothenate biosynthetic precursor is α -ketoisovalerate. Preferably, valine or α -ketoisovalerate is added in an amount that results in a concentration in the medium sufficient for production of the desired product (e.g., panto-compound) to occur. Pantothenate biosynthetic precursors are also referred to herein as "supplemental pantothenate biosynthetic substrates".

Providing pantothenate biosynthetic precursors in the pantothenate biosynthetic methodologies of the present invention, can be associated with high costs, for example, when the methodologies are used to produce high yields of panto-compounds. Accordingly, preferred methodologies of the present invention feature microorganisms having at least one biosynthetic enzyme or combination of biosynthetic enzymes (e.g., at least one pantothenate biosynthetic enzyme and/or valine-isoleucine biosynthetic enzyme) manipulated such that pantothenate or other desired panto-compounds are produced in a manner independent of precursor feed. The phrase "a manner independent of precursor feed", for example, when referring to a method for producing a desired compound (e.g., a panto-compound), includes an approach to or a mode of producing the desired compound that does not depend or rely on precursors being provided (e.g., fed) to the microorganism being utilized to produce the desired compound. For example, microorganisms featured in the methodologies of the present invention can be used to produce panto-compounds in a manner requiring no feeding of the precursors aspartate, β-alanine, valine and/or α-KIV.

Alternative preferred methodologies of the present invention feature microorganisms having at least one biosynthetic enzyme or combination of biosynthetic enzymes manipulated such that pantothenate or other desired panto-compounds are

20

25

30

35

produced in a manner substantially independent of precursor feed. The phrase "a manner substantially independent of precursor feed" includes an approach to or a method of producing the desired compound that depends or relies to a lesser extent on precursors being provided (e.g., fed) to the microorganism being utilized. For example, microorganisms featured in the methodologies of the present invention can be used to produce panto-compounds in a manner requiring feeding of substantially reduced amounts of the precursors aspartate, β -alanine, valine and/or α -KIV. In one embodiment, the invention features methods of producing panto-compounds (e.g., pantothenate) in a manner that requires feeding of less than 5%-10% of the amount of precursor required by a control microorganism (e.g., a microorganism that is dependent, for example is wholly dependent, on precursor feed to efficiently produce the desired compound). In another embodiment, the invention features methods of producing panto-compounds in a manner that requires feeding of less than 15-20% of the amount of precursor required by a control microorganism. In another embodiment, the 15 invention features methods of producing panto-compounds in a manner that requires feeding of less than 25-30%, 35-40%, 45-50% or 55-60% of the amount of precursor required by a control microorganism. As described in Examples I-III herein, particular microorganisms featured in the methodologies of the present invention require, for example, 5 g/L of aspartate, β-alanine, valine or α-KIV (e.g., in test tube or in shake flask cultures). Accordingly, in a preferred embodiment, the present invention features methods of producing panto-compounds (e.g., pantothenate) in a manner requiring feeding of less than 0.25 g/L, 0.5 g/L, 0.75 g/L, 1 g/L, 1.25 g/L, 1.5 g/L, 1.75 g/L, 2 g/L, 2.25 g/L, 2.5 g/L, 2.75 g/L or 3 g/L.

Preferred methods of producing desired compounds (e.g., panto-compounds) in a manner independent of precursor feed or alternatively, in a manner substantially independent of precursor feed, involve culturing microorganisms which have been manipulated (e.g., designed or engineered, for example, genetically engineered) such that expression of at least one pantothenate biosynthetic enzyme, and/or at least one isoleucine-valine biosynthetic enzyme, is modified. For example, in one embodiment, a microorganism is manipulated (e.g., designed or engineered) such that the production of at least one pantothenate biosynthetic enzyme, and/or at least one isoleucine/valine biosynthetic enzyme is deregulated. In a preferred embodiment, a microorganism is manipulated (e.g., designed or engineered) such that it has a deregulated biosynthetic pathway, for example, a deregulated pantothenate biosynthesis pathway and/or a deregulated isoleucine-valine biosynthetic pathway, as defined herein. In another preferred embodiment, a microorganism is manipulated (e.g., designed or engineered)

such that at least one pantothenate biosynthetic enzyme, and/or at least one isoleucinevaline biosynthetic enzyme is overexpressed.

Preferred methods of producing desired compounds (e.g., panto-compounds) in a manner independent of precursor feed or alternatively, in a manner substantially independent of precursor feed, are as follows. In one embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism having a deregulated isoleucine-valine (ilv) pathway under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism having a deregulated pantothenate (pan) pathway'and a deregulated isoleucine-valine (ilv) pathway, under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of aspartate or β -alanine feed comprising culturing an aspartate- α decarboxylase-overexpressing (AaD-O) microorganism under conditions such that pantothenate is produced. In yet another embodiment, the invention features a method of producing pantothenate in a manner independent of valine or α-ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced.

The term "aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism" includes a microorganism which has been manipulated such that aspartate- α -decarboxylase is overexpressed. A preferred "aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism" has been transformed with a vector comprising a *B. subtilis* panD nucleic acid sequence (e.g., a panD nucleic acid sequence that encodes an aspartate- α -decarboxylase protein having the amino acid sequence of SEQ ID NO:28, for example, a panD nucleic acid sequence as set forth in SEQ ID NO:27).

The phrase "microorganism having a deregulated isoleucine-valine (ilv) pathway" includes a microorganism having an alteration or modification in at least one gene encoding an enzyme of the isoleucine-valine (ilv) pathway or having an alteration or modification in an operon including more than one gene encoding an enzyme of the isoleucine-valine (ilv) pathway. A preferred "microorganism having a deregulated isoleucine-valine (ilv) pathway" overexpresses acetohydroxyacid synthetase (e.g., acetohydroxyacid synthetase having subunits having the amino acid sequences of SEQ ID NO:34 or acetohydroxyacid synthetase having the amino acid sequence of SEQ ID NO:87), acetohydroxyacid isomeroreductase (having the amino acid sequence of SEQ ID NO:36), or dihydroxyacid dehydratase (having the amino acid

012177242 | -

sequence of SEQ ID NO:38) and/or has been transformed with a vector comprising *ilvB*, *ilvN*, *ilvC*, *ilvBN*, *ilvBNC* or *alsS* nucleic acid sequences (SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, nucleotides 1-2246 of SEQ ID NO:58, SEQ ID NO:58 having coding regions from nucleotides 1-1725, 1722-2246 and 2263-3291, or SEQ ID NO:86, respectively) and/or an *ilvD* nucleic acid sequence (SEQ ID NO:37).

IV. High Yield Production Methodologies

A particularly preferred embodiment of the present invention is a high yield production method for producing pantothenate comprising culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly 10 high yield. The phrase "high yield production method", for example, a high yield production method for producing a desired compound (e.g., for producing a pantocompound) includes a method that results in production of the desired compound at a level which is elevated or above what is usual for comparable production methods. 15 Preferably, a high yield production method results in production of the desired compound at a significantly high yield. The phrase "significantly high yield" includes a level of production or yield which is sufficiently elevated or above what is usual for comparable production methods, for example, which is elevated to a level sufficient for commercial production of the desired product (e.g., production of the product at a commercially feasible cost). In one embodiment, the invention features a high yield 20 production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 2 g/L. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 10 g/L. In 25 another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 20 g/L. In yet another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under 30 conditions such that pantothenate is produced at a level greater than 30 g/L. In yet another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 40 g/L.

The invention further features a high yield production method for producing a desired compound (e.g., for producing a panto-compound) that involves culturing a manipulated microorganism under conditions such that a sufficiently elevated level of

compound is produced within a commercially desireable period of time. In an exemplary embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 15-20 g/L in 36 hours. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 25-30 g/L in 48 hours. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 35-40 g/L in 72 hours, for example, greater that 37 g/L in 72 hours. In another-embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 30-40 g/L in 60 hours, for example, greater that 30, 35 or 40 g/L in 60 hours. Values and ranges included and/or intermediate within the ranges 15 set forth herein are also intended to be within the scope of the present invention. For example, pantothenate production at levels of at least 31, 32, 33, 34, 35, 36, 37, 38 and 39 g/L in 60 hours are intended to be included within the range of 30-40 g/L in 60 hours. In another example, ranges of 30-35 g/L or 35-40 g/L are intended to be included within the range of 30-40 g/L in 60 hours. Moreover, the skilled artisan will appreciate that culturing a manipulated microorganism to achieve a production level of, for example, "30-40 g/L in 60 hours" includes culturing the microorganism for additional time periods (e.g., time periods longer than 60 hours), optionally resulting in even higher yields of pantothenate being produced.

25

30

35

V. Panto-Compound Production Methodologies Featuring Pantothenate Kinase Mutant Microorganisms

The present invention relates to methods of producing pantothenate using microorganisms engineered to produce high yields of pantothenate as well as other panto-compounds. Cells overproducing pantothenate result in high intracellular pantothenate levels that could overcome the feedback inhibition of pantothenate kinase by CoA, leading to overproduction of CoA. Besides consuming pantothenate, increased synthesis of CoA may cause increased feedback inhibition of the PanB, PanD, PanE or PanC reaction, thereby limiting pantothenate production. Accordingly, a reduction in pantothenate kinase activity may lead to a decrease in CoA levels with resulting increases in PanB, PanD, PanE or PanC activity and pantothenate production.

15

20

25

30

35

Thus, certain methodologies of the present invention are based, at least in part, on the identification and characterization of the *B. subtilis coaA* gene and the demonstration that the gene is neither essential for *B. subtilis* growth (*i.e.*, deletion of the coaA gene from the chromosome of *B. subtilis* is not a lethal event) nor for pantothenate kinase activity in *B. subtilis*. A second pantothenate kinase-encoding gene has been identified and characterized in *B. subtilis*, and is termed "coaX". This gene complements an *E. coli* mutant that contains a temperature sensitive pantothenate kinase and is not related by homology to any previously known pantothenate kinase gene.

In one aspect, the methodologies of the invention feature recombinant microorganisms that include the coaX gene or that include a mutant coaX gene, having reduced pantothenate kinase activity. In one embodiment, the methodologies feature such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway. In another embodiment, the methodologies feature such recombinant microorganisms further having a deregulated isoleucine-valine (ilv) pathway. In a preferred embodiment, the microorganisms belong to the genus Bacillus (e.g., B. subtilis).

The methodologies of the invention also feature recombinant microorganisms (e.g., microorganisms belong to the genus Bacillus, for example, B. subtilis) that include the coaA gene or that include a mutant coaA gene, optionally including a coaX gene or mutant thereof, having reduced pantothenate kinase activity. In one embodiment, the methodologies feature such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway or having a deregulated isoleucine-valine (ilv) pathway. Also featured are vectors that include isolated coaX or coaA genes as well as mutant coaX and/or coaA genes. Isolated nucleic acid molecules that include isolated coaX genes or mutant coaX genes are features in addition to isolated CoaX proteins and mutant CoaX proteins.

The above-described nucleic acid molecules (e.g., genes), proteins, vectors, and recombinant microorganisms (e.g., mutant microorganisms), are particularly suited for use in methods of producing panto-compounds and/or methods of enhancing panto-compound production. In one embodiment, the invention features a method for producing a panto-compound (e.g., pantothenate) that includes culturing a pantothenate kinase mutant (e.g., a recombinant microorganism that misexpresses, e.g., is mutated for, pantothenate kinase, as defined herein) under conditions such that panto-compound is produced. In another embodiment, the invention features a method for enhancing production of a panto-compound (e.g., pantothenate) that includes culturing a pantothenate kinase mutant (e.g., a recombinant microorganism that misexpresses, e.g., is mutated for, pantothenate kinase, as defined herein) under conditions such that

production of the panto-compound is produced. As used herein, the term "enhancing" (for example, in the context of the phrase "enhancing production") includes increasing the level or rate of production of panto-compound (e.g., pantothenate) as compared to the level or rate of production in a non-mutant microorganism (e.g., a microorganism having a normal pantothenate kinase gene(s) and/or having normal pantothenate production rates and/or levels.

Preferably, the level of panto-compound produced in methodologies featuring the pantothenate kinase mutants of the present invention is increased by at least 5% as compared to the level produced by a non-mutant (e.g., a recombinant microorganism expressing non-mutated pantothenate kinase). Even more preferably, the level of panto-compound is increased 10% as compared to methodologies featuring non-mutants. Even more preferably, panto-compound levels (e.g., pantothenate levels) are increased 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, are increased 2-fold, 5-fold, 10-fold, 50-fold, 100-fold or more as compared to methodologies featuring non-mutants.

15

20

VI. Additional Mutations Resulting in Enhanced Panto-Compound Production

The methodologies of the present invention further can include, for example in addition to overexpressing or deregulating a pantothenate biosynthetic enzyme and/or an isoleucine-valine biosynthetic enzyme, or in addition to mutating a pantothenate-kinase encoding gene, deleting or mutating an enzyme that catalyzes the conversion of key pantothenate biosynthesis substrates or precursors to unwanted or undesirable products. For example, mutating the ilvE gene (Kuramitsu et al. (1985) J. Biochem. 97:993-999) or a homologue thereof (SEQ ID NO:62 or SEQ ID NO:64), thereby limiting the conversion of a-ketoisovalerate to valine, in addition to mutating a pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or increased production of panto-compound. Alternatively, mutating the ansB gene (Sun and Seflow (1991) J. Bacteriol. 173:3831-3845) or a homologue thereof (SEQ ID NO:66), thereby limiting the degradation of aspartate, in addition to mutating a pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or increased production of pantocompound. Alternatively, mutating the alsD gene (Renna et al. (1993) J. Bacteriol. 175:3863-3875) or a homologue thereof (SEQ ID NO:68), thereby limiting the conversion of acetolactate to acetoin, in addition to mutating a pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or increased production of panto-compound. Alternatively, mutating the avtA gene encoding alanine-valine transaminase or a homologue thereof, thereby limiting the conversion of aketoisovalerate to valine, in addition to mutating a pantothenate kinase encoding

20

35

0121772A2 I >

enzyme, is predicted to result in even further enhanced or increased production of panto-compound. Mutating the *avtA* gene can include mutating, for example, an *avtA* gene having the nucleotide sequence of SEQ ID NO:70 (*e.g.*, the *E. coli avtA* gene), or a structural homolog thereof (*e.g.*, a homologue encoding a protein having 30-40%, 40-50%, 50-60%, 60-70%, 70-80%, 80-90%, 90-95% or more identity with the amino acid sequence of SEQ ID NO:71) or a functional homologue (*e.g.*, a gene encoding a structurally unrelated protein having alanine-valine transmainase activity. Such mutations can be accomplished using the methodologies as exemplified in the Examples (*e.g.*, Examples XIII, XV, XVI and XVII).

Accordingly, in one embodiment, the invention features a method of producing a panto-compound which includes culturing a microorganism having a mutant pantothenate kinase-encoding gene and which further has a deletion or mutation in an avtA, ilvE, ansB, and/or alsD gene, or homologue thereof. In another embodiment, the invention features a method of producing a panto-compound which includes culturing a microorganism having a mutant pantothenate-kinase encoding gene and a deregulated pantothenate biosynthetic pathway enzyme and which further has a deletion or mutation in an avtA, ilvE, ansB, and/or alsD gene, or homologue thereof. In another embodiment, the invention features a method of producing a panto-compound which includes culturing a microorganism having a mutant pantothenate-kinase encoding gene and a deregulated isoleucine-valine biosynthetic pathway enzyme and which further has a deletion or mutation in an avtA, ilvE, ansB, and/or alsD gene, or homologue thereof.

Mutating the *alsD* gene can be particularly useful when accomplished in conjunction with overexpression or deregulation of the *alsS* gene, for example, to prevent carbon (*e.g.*, acetolactate) from being drawn away from the precursor pool utilized for α-KIV production. Accordingly, to maximize the contribution of the *als* locus to panto-compound production, it is desirable to disrupt the *alsD* gene in addition to overexpressing the *alsS* gene. To disrupt the *alsD* gene, appropriate fragments of the *als* operon, flanking the *alsD* gene, are amplified by PCR and cloned to provide homology for creating the disruptions. A drug resistance gene, such as the *cat* gene, is cloned between the flanking DNA fragments in place of the *alsD* gene, and the linearized DNA is transformed into a pantothenate production strain such as PA824, selecting for drug-resistance. To overexpress *alsS*, the *alsS* coding sequence (*e.g.*, an *alsS* coding sequence that has been engineered by PCR for expression) is cloned into an expression vector. Vectors which express *alsS* (or alternatively, vectors which express *alsS* plus *ilvC*) are the introduced into panto-compound production strains (*e.g.*, the pantothenate producing strain PA824).

10

15

20

25

30

35

The methodologies of the present invention further can include, for example in addition to overexpressing or deregulating a pantothenate biosynthetic enzyme and/or an isoleucine-valine biosynthetic enzyme, or in addition to mutating a pantothenate-kinase encoding gene, deleting or mutating an enzyme that catalyzes the conversion of desired panto-compounds to unwanted or undesireable downstream products.

VII. Isolated Nucleic Acid Molecules and Genes

Another aspect of the present invention features isolated nucleic acid molecules that encode *Bacillus* proteins (e.g., B. subtilis proteins), for example, *Bacillus* pantothenate biosynthetic enzymes (e.g., B. subtilis pantothenate biosynthetic enzymes) or *Bacillus* valine-isoleucine biosynthetic enzymes (e.g., B. subtilis valine-isoleucine biosynthetic enzymes). Also featured are isolated coaX and/or coaA nucleic acid molecules (e.g., isolated coaX and/or coaA genes) as well as isolated nucleic acid molecules that include such coaX and/or coaA nucleic acid molecules or genes.

The term "nucleic acid molecule" includes DNA molecules (e.g., linear, circular, cDNA or chromosomal DNA) and RNA molecules (e.g., tRNA, rRNA, mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA. The term "isolated" nucleic acid molecule includes a nucleic acid molecule which is free of sequences which naturally flank the nucleic acid molecule (i.e., sequences located at the 5' and 3' ends of the nucleic acid molecule) in the chromosomal DNA of the organism from which the nucleic acid is derived. In various embodiments, an isolated nucleic acid molecule can contain less than about 10 kb, 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb, 0.1 kb, 50 bp, 25 bp or 10 bp of nucleotide sequences which naturally flank the nucleic acid molecule in chromosomal DNA of the microorganism from which the nucleic acid molecule is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular materials when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

The term "gene", as used herein, includes a nucleic acid molecule (e.g., a DNA molecule or segment thereof), for example, a protein or RNA-encoding nucleic acid molecule, that in an organism, is separated from another gene or other genes, by intergenic DNA (i.e., intervening or spacer DNA which naturally flanks the gene and/or separates genes in the chromosomal DNA of the organism). A gene may direct synthesis of an enzyme or other protein molecule (e.g., may comprise coding sequences, for example, a contiguous open reading frame (ORF) which encodes a protein) or may itself be functional in the organism. A gene in an organism, may be clustered in an

15

20

25

30

35

operon, as defined herein, said operon being separated from other genes and/or operons by the intergenic DNA. Individual genes contained within an operon may overlap without intergenic DNA between said individual genes. An "isolated gene", as used herein, includes a gene which is essentially free of sequences which naturally flank the gene in the chromosomal DNA of the organism from which the gene is derived (i.e., is free of adjacent coding sequences which encode a second or distinct protein or RNA molecule, adjacent structural sequences or the like) and optionally includes 5' and 3' regulatory sequences, for example promoter sequences and/or terminator sequences. In one embodiment, an isolated gene includes predominantly coding sequences for a protein (e.g., sequences which encode Bacillus proteins). In another embodiment, an isolated gene includes coding sequences for a protein (e.g., for a Bacillus protein) and adjacent 5' and/or 3' regulatory sequences from the chromosomal DNA of the organism from which the gene is derived (e.g., adjacent 5' and/or 3' Bacillus regulatory sequences). Preferably, an isolated gene contains less than about 10 kb, 5 kb, 2 kb, 1 kb, 0.5 kb, 0.2 kb, 0.1 kb, 50 bp, 25 bp or 10 bp of nucleotide sequences which naturally flank the gene in the chromosomal DNA of the organism from which the gene is derived.

In one aspect, the present invention features isolated *panB* nucleic acid sequences or genes, isolated *panC* nucleic acid sequences or genes, isolated *panD* nucleic acid sequences or genes, isolated *panE* nucleic acid sequences or genes, isolated *ilvB*, *ilvN*, *ilvBN* nucleic acid sequences or genes, isolated *alsS* nucleic acid sequences or genes, isolated *ilvD* nucleic acid sequences or genes and/or isolated *ilvD* nucleic acid sequences or genes.

In a preferred embodiment, the nucleic acid or gene is derived from Bacillus (e.g., is Bacillus-derived). The term "derived from Bacillus" or "Bacillus-derived" includes a nucleic acid or gene which is naturally found in microorganisms of the genus Bacillus. Preferably, the nucleic acid or gene is derived from a microorganism selected from the group consisting of Bacillus subtilis, Bacillus lentimorbus, Bacillus lentus, Bacillus firmus, Bacillus pantothenticus, Bacillus amyloliquefaciens, Bacillus cereus, Bacillus circulans, Bacillus coagulans, Bacillus licheniformis, Bacillus megaterium, Bacillus pumilus, Bacillus thuringiensis, and other Group 1 Bacillus species, for example, as characterized by 16S rRNA type (Priest, supra). In another preferred embodiment, the nucleic acid or gene is derived from Bacillus brevis or Bacillus stearothermophilus. In another preferred embodiment, the nucleic acid molecules and/or genes of the present invention are derived from a microorganism selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus. In a particularly preferred

embodiment, the nucleic acid or gene is derived from Bacillus subtilis (e.g., is Bacillus subtilis-derived). The term "derived from Bacillus subtilis" or "Bacillus subtilis-derived" includes a nucleic acid or gene which is naturally found in Bacillus subtilis. In yet another preferred embodiment, the nucleic acid or gene is a Bacillus gene homologue (e.g., is derived from a species distinct from Bacillus but having significant homology to a Bacillus gene of the present invention, for example, a Bacillus pan gene or Bacillus ilv gene).

Included within the scope of the present invention are bacterial-derived nucleic acid molecules or genes and/or Bacillus-derived nucleic acid molecules or genes (e.g., B. subtilis-derived nucleic acid molecules or genes), for example, the genes identified by 10 the present inventors, for example, Bacillus or B. subtilis coaX genes, coaA genes, pan genes and/or ilv genes. Further included within the scope of the present invention are bacterial-derived nucleic acid molecules or genes and/or Bacillus-derived nucleic acid molecules or genes (e.g., B. subtilis-derived nucleic acid molecules or genes) (e.g., B. subtilis nucleic acid molecules or genes) which differ from naturally-occurring bacterial and/or Bacillus nucleic acid molecules or genes (e.g., B. subtilis nucleic acid molecules or genes), for example, nucleic acid molecules or genes which have nucleic acids that are substituted, inserted or deleted, but which encode proteins substantially similar to the naturally-occurring gene products of the present invention. In one embodiment, an isolated nucleic acid molecule comprises at least one of the nucleotide sequences set 20 forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:86, SEQ ID NO:35 or SEQ ID NO:37. In another preferred embodiment, an isolated nucleic acid molecule comprises at least two, three or four of the nucleotide sequences set forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37. For example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequences of SEQ ID NO:23, SEQ ID NO:25 and SEQ ID NO:27, preferably linked such that the proteins encoded by the nucleotide sequences of SEQ ID NO:23, SEQ ID NO:25 and SEQ ID NO:27 are each produced when the isolated nucleic acid molecule is expressed in a microorganism (e.g., 30 SEQ ID NO:59). In another example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequences of SEQ ID NO:31 and SEQ ID NO:33, preferably linked such that the proteins encoded by the nucleotide sequences of SEQ ID NO:31 and SEQ ID NO:33 are each produced when the isolated nucleic acid molecule is expressed in a microorganism (e.g., nucleotides 1-2246 of SEQ ID NO:58). 35 In another example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequence of SEQ ID NO:86. In another example, a preferred

isolated nucleic acid molecule of the present invention can include the nucleotide sequences of SEQ ID NO:31, SEQ ID NO:33 and SEQ ID NO:35, preferably linked such that the proteins encoded by the nucleotide sequences of SEQ ID NO:31, SEQ ID NO:33 and SEQ ID NO:35 are each produced when the isolated nucleic acid molecule is expressed in a microorganism (e.g., SEQ ID NO:58).

In another embodiment, an isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which is at least about 60-65%, preferably at least about 70-75%, more preferable at least about 80-85%, and even more preferably at least about 90-95% or more identical to a nucleotide sequence set forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37. In another embodiment, an isolated nucleic acid molecule hybridizes under stringent conditions to a nucleic acid molecule having a nucleotide sequence set forth as SEQ ID NO:23, SEQ ID NO:25, SEO ID NO:27, SEO ID NO:29, SEO ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEO ID NO:37. Such stringent conditions are known to those skilled in the art and can be found in Current Protocols in Molecular Biology, John Wiley & Sons. N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent (e.g. high stringency) hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that 20 hybridizes under stringent conditions to the sequence of SEQ ID NO:23, SEQ ID NO:25, SEO ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature. 25

A nucleic acid molecule of the present invention (e.g., a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37 can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, nucleic acid molecules can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) or can be isolated by the polymerase chain reaction using synthetic oligonucleotide primers designed based upon the sequence of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEO ID NO:35 or SEQ ID NO:37. A nucleic acid of the invention can be

30

amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:33, SEQ ID NO:33, SEQ ID NO:35.

Additional panC nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:25, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:26 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:26 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:25 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:26, or are complementary to a panC nucleotide sequence as set forth herein.

Aditional panD nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:27, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:28 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:28 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:27 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:28, or are complementary to a panD nucleotide sequence as set forth herein.

Additional panE nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:29, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:30 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:30 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:29 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:30, or are complementary to a panE nucleotide sequence as set forth herein.

35

20

25

30

35

Additional ilvB nucleic acid sequences are those that comprise the nucleotide sequence of SEQ ID NO:31, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:32 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:32 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:31 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:32, or are complementary to an ilvB nucleotide sequence as set forth herein.

Additional ilvN nucleic acid sequences are those that comprise the nucleotide sequence of SEQ ID NO:33, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:34 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having 15 the amino acid sequence as set forth in SEQ ID NO:34 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:33 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:34, or are complementary to an ilvN nucleotide sequence as set forth herein.

Additional ilvC nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:35, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:36 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:36 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:35 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:36, or are complementary to an ilvC nucleotide sequence as set forth herein.

Additional ilvD nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:37, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:38 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:38 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:37 or

to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:38, or are complementary to an *ilvD* nucleotide sequence as set forth herein.

Additional alsS nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:86, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:87 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:87 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:86 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:87, or are complementary to an alsS nucleotide sequence as set forth herein.

In another embodiment, an isolated nucleic acid molecule is or includes a coaX gene, or portion or fragment thereof. In one embodiment, an isolated coaX nucleic acid molecule or gene comprises the nucleotide sequence as set forth in SEQ ID NO:19 (e.g., comprises the B. subtilis coaX nucleotide sequence). In another embodiment, an isolated coaX nucleic acid molecule or gene comprises a nucleotide sequence that encodes the amino acid sequence as set forth in SEQ ID NO:9 (e.g., encodes the B. subtilis CoaX amino acid sequence). In yet another-embodiment, an isolated coaX 20 nucleic acid molecule or gene encodes a homologue of the CoaX protein having the amino acid sequence of SEQ ID NO:9. As used herein, the term "homologue" includes a protein or polypeptide sharing at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identity with the amino acid sequence of a wild-type protein or 25 polypeptide described herein and having a substantially equivalent functional or biological activity as said wild-type protein or polypeptide. For example, a CoaX homologue shares at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identity with the protein having the amino acid sequence set forth as SEQ ID NO:9 and has a substantially equivalent functional or biological activity (i.e., is a functional equivalent) of the protein having the amino acid sequence set forth as SEQ ID NO:9 (e.g., has a substantially equivalent pantothenate kinase activity). In a preferred embodiment, an isolated coaX nucleic acid molecule or gene comprises a nucleotide sequence that encodes a polypeptide as set forth in any one of SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID

NO:74 or SEQ ID NO:75. In another embodiment, an isolated coaX nucleic acid molecule hybridizes to all or a portion of a nucleic acid molecule having the nucleotide sequence set forth in SEQ ID NO:19 or hybridizes to all or a portion of a nucleic acid molecule having a nucleotide sequence that encodes a polypeptide having the amino acid sequence of any of SEQ ID NOs:7-18, 74 or 75. Such hybridization conditions are known to those skilled in the art and can be found in Current Protocols in Molecular Biology, Ausubel et al., eds., John Wiley & Sons, Inc. (1995), sections 2, 4 and 6. Additional stringent conditions can be found in Molecular Cloning: A Laboratory Manual, Sambrook et al., Cold Spring Harbor Press, Cold Spring Harbor, NY (1989), chapters 7, 9 and 11. A preferred, non-limiting example of stringent hybridization 10 conditions includes hybridization in 4X sodium chloride/sodium citrate (SSC), at about 65-70°C (or hybridization in 4X SSC plus 50% formamide at about 42-50°C) followed by one or more washes in 1X SSC, at about 65-70°C. A preferred, non-limiting example of highly stringent hybridization conditions includes hybridization in 1X SSC, 15 at about 65-70°C (or hybridization in 1X SSC plus 50% formamide at about 42-50°C) followed by one or more washes in 0.3X SSC, at about 65-70°C. A preferred, nonlimiting example of reduced stringency hybridization conditions includes hybridization in 4X SSC, at about 50-60°C (or alternatively hybridization in 6X SSC plus 50% formamide at about 40-45°C) followed by one or more washes in 2X SSC, at about 50-60°C. Ranges intermediate to the above-recited values, e.g., at 65-70°C or at 42-50°C 20 are also intended to be encompassed by the present invention. SSPE (1X SSPE is 0.15 M NaCl, 10mM NaH₂PO₄, and 1.25 mM EDTA, pH 7.4) can be substituted for SSC (1X SSC is 0.15 M NaCl and 15 mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes each after hybridization is complete. The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length 25 should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, $T_m(^{\circ}C) = 2(\# \text{ of } A + T \text{ bases}) + 4(\# \text{ of } G + C \text{ bases})$. For hybrids between 18 and 49 base pairs in length, $T_m(^{\circ}C) = 81.5 + 16.6(\log_{10}[Na^+]) + 0.41(\%G+C) - (600/N)$, where N is the number of bases in the hybrid, and [Na⁺] is the concentration of sodium 30 ions in the hybridization buffer ($[Na^+]$ for 1X SSC = 0.165 M). It will also be recognized by the skilled practitioner that additional reagents may be added to hybridization and/or wash buffers to decrease non-specific hybridization of nucleic acid molecules to membranes, for example, nitrocellulose or nylon membranes, including but not limited to blocking agents (e.g., BSA or salmon or herring sperm carrier DNA), 35 detergents (e.g., SDS), chelating agents (e.g., EDTA), Ficoll, PVP and the like. When using nylon membranes, in particular, an additional preferred, non-limiting example of

stringent hybridization conditions is hybridization in 0.25-0.5M NaH₂PO₄, 7% SDS at about 65°C, followed by one or more washes at 0.02M NaH₂PO₄, 1% SDS at 65°C, see e.g., Church and Gilbert (1984) Proc. Natl. Acad. Sci. USA 81:1991-1995, (or, alternatively, 0.2X SSC, 1% SDS). In another preferred embodiment, an isolated nucleic acid molecule comprises a nucleotide sequence that is complementary to a coaX nucleotide sequence as set forth herein (e.g., is the full complement of the nucleotide sequence set forth as SEQ ID NO:19).

In another preferred embodiment, an isolated nucleic acid molecule is or includes a coaA gene, for example, a Bacillus (e.g., B. subtilis) coaA gene, or portion or fragment thereof. Exemplary isolated coal nucleic acid molecules and/or genes include (1) an isolated coaA nucleic acid molecule or gene comprising the nucleotide sequence as set forth in any one of SEQ ID NOs:20-22; (2) an isolated coal nucleic acid molecule or gene comprising a nucleotide sequence that encodes the amino acid sequence as set forth in SEQ ID NO:3; (3) an isolated coal nucleic acid molecule or gene comprising a nucleotide sequence which encodes a CoaA homologue (e.g., a polypeptide having an amino acid sequence at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identical to the amino acid sequence set forth as SEQ ID NO:3 and having a substantially equivalent enzymatic activity; (4) an isolated coaA nucleic acid molecule or gene comprising a nucleotide sequence that encodes a polypeptide as set forth in any one of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6; (5) an isolated nucleic acid molecule that hybridizes under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence set forth in SEQ ID NO:20, SEQ ID NO:21 or SEQ ID NO:22 or hybridizes to all or a portion of a nucleic acid molecule having a nucleotide sequence that encodes a polypeptide having the amino acid sequence of SEQ ID NO:3; and (6) an isolated nucleic acid molecule comprising a nucleotide sequence that is complementary to a coal nucleotide sequence as set forth herein (e.g., is the full complement of the nucleotide sequence set forth in SEQ ID NO:20, SEQ ID NO:21 or SEQ ID NO:22).

A nucleic acid molecule of the present invention (e.g., a coaX nucleic acid molecule or gene or a coaA nucleic acid molecule or gene), can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, nucleic acid molecules can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) or can be isolated by the polymerase chain reaction using synthetic oligonucleotide primers designed

20

based upon the coaX or coaA nucleotide sequences set forth herein, or flanking sequences thereof. A nucleic acid of the invention (e.g., a coaX nucleic acid molecule or gene or a coaA nucleic acid molecule or gene), can be amplified using cDNA, mRNA or alternatively, chromosomal DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques.

Yet another embodiment of the present invention features mutant coaX and coaA nucleic acid molecules or genes. The phrase "mutant nucleic acid molecule" or "mutant gene" as used herein, includes a nucleic acid molecule or gene having a nucleotide sequence which includes at least one alteration (e.g., substitution, insertion, deletion) such that the polypeptide or protein that may be encoded by said mutant exhibits an activity that differs from the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene. Preferably, a mutant nucleic acid molecule or mutant gene (e.g., a mutant coaA or coaX gene) encodes a polypeptide or protein having a reduced activity (e.g., having a reduced pantothenate kinase activity) as compared to the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene, for example, when assayed under similar conditions (e.g., assayed in microorganisms cultured at the same temperature). A mutant gene also can encode no polypeptide or have a reduced level of production of the wild-type polypeptide.

As used herein, a "reduced activity" or "reduced enzymatic activity" is one that is at least 5% less than that of the polypeptide or protein encoded by the wild-type 20 nucleic acid molecule or gene, preferably at least 5-10% less, more preferably at least 10-25% less and even more preferably at least 25-50%, 50-75% or 75-100% less than that of the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene. Ranges intermediate to the above-recited values, e.g., 75-85%, 85-90%, 90-95%, are also intended to be encompassed by the present invention. As used herein, a "reduced activity" or "reduced enzymatic activity" also includes an activity that has been deleted or "knocked out" (e.g., approximately 100% less activity than that of the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene). Activity can be determined according to any well accepted assay for measuring activity of a particular protein of interest. Activity can be measured or assayed directly, for example, measuring an activity of a protein isolated or purified from a cell. Alternatively, an activity can be measured or assayed within a cell or in an extracellular medium. For example, assaying for a mutant coaA gene or a mutant coaX gene (i.e., said mutant encoding a reduced pantothenate kinase activity) can be accomplished by expressing the mutated gene in a microorganism, for example, a mutant microorganism which expresses pantothenate kinase in a temperature-sensitive manner, assaying the mutant gene for the ability to complement a temperature sensitive (Ts) mutant for

15

20

30

35

pantothenate kinase activity. A coaX mutant gene or coaA mutant gene that encodes a "reduced pantothenate kinase activity" is one that complements the Ts mutant less effectively than, for example, a corresponding wild-type coaX gene or coaA gene.

It will be appreciated by the skilled artisan that even a single substitution in a nucleic acid or gene sequence (e.g., a base substitution that encodes an amino acid change in the corresponding amino acid sequence) can dramatically affect the activity of an encoded polypeptide or protein as compared to the corresponding wild-type polypeptide or protein. A mutant nucleic acid or mutant gene (e.g., encoding a mutant polypeptide or protein), as defined herein, is readily distinguishable from a nucleic acid or gene encoding a protein homologue, as described above, in that a mutant nucleic acid or mutant gene encodes a protein or polypeptide having an altered activity, optionally observable as a different or distinct phenotype in a microorganism expressing said mutant gene or nucleic acid or producing said mutant protein or polypeptide (i.e., a mutant microorganism) as compared to a corresponding microorganism expressing the wild-type gene or nucleic acid or producing said mutant protein or polypeptide. By contrast, a protein homologue has an identical or substantially similar activity, optionally phenotypically indiscernable when produced in a microorganism, as compared to a corresponding microorganism expressing the wild-type gene or nucleic acid. Accordingly it is not, for example, the degree of sequence identity between nucleic acid molecules, genes, protein or polypeptides that serves to distinguish between homologues and mutants, rather it is the activity of the encoded protein or polypeptide that distinguishes between homologues and mutants: homologues having, for example, low (e.g., 30-50% sequence identity) sequence identity yet having substantially equivalent functional activities, and mutants, for example sharing 99% sequence identity yet having dramatically different or altered functional activities. Exemplary 25 homologues are set forth in Figure 20 (i.e., CoaA homologues) and in Figure 23 (i.e., CoaX homologues). Exemplary mutants are described in Examples XV and XVIII herein.

Recombinant Nucleic Acid Molecules and Vectors

The present invention further features recombinant nucleic acid molecules (e.g., recombinant DNA molecules) that include nucleic acid molecules and/or genes described herein (e.g., isolated nucleic acid molecules and/or genes), preferably Bacillus genes, more preferably Bacillus subtilis genes, even more preferably Bacillus subtilis pantothenate kinase genes (e.g., coaX genes or coaA genes), pantothenate biosynthetic genes (e.g., genes encoding pantothenate biosynthetic enzymes, for example, panB genes encoding ketopantoate hydroxymethyltransferase, panE genes encoding

10

20

25

30

35

0101770A0 L -

ketopantoate reductase, panC genes encoding pantothenate synthetase, and/or panD genes encoding aspartate-α-decarboxylase) and/or isoleucine-valine (ilv) biosynthetic genes (e.g., ilvBN or alsS genes encoding acetohydroxyacid synthetase, ilvC genes encoding acetohydroxyacid isomeroreductase and/or ilvD genes encoding dihydroxyacid dehydratase).

The present invention further features vectors (e.g., recombinant vectors) that include nucleic acid molecules (e.g., isolated or recombinant nucleic acid molecules and/or genes) described herein. In particular, recombinant vectors are featured that include nucleic acid sequences that encode bacterial gene products as described herein, preferably Bacillus gene products, more preferably Bacillus subtilis gene products, even more preferably Bacillus subtilis pantothenate biosynthetic gene products (e.g., pantothenate biosynthetic enzymes, for example, ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase, and/or aspartate-α-decarboxylase) and/or isoleucine-valine biosynthetic gene products (e.g., 15 acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase and/or dihydroxyacid dehydratase).

The term "recombinant nucleic acid molecule" includes a nucleic acid molecule (e.g., a DNA molecule) that has been altered, modified or engineered such that it differs in nucleotide sequence from the native or natural nucleic acid molecule from which the recombinant nucleic acid molecule was derived (e.g., by addition, deletion or substitution of one or more nucleotides). Preferably, a recombinant nucleic acid molecule (e.g., a recombinant DNA molecule) includes an isolated nucleic acid molecule or gene of the present invention (e.g., an isolated coaX, coaA, pan or ilv gene) operably linked to regulatory sequences.

The term "recombinant vector" includes a vector (e.g., plasmid, phage, phasmid, virus, cosmid or other purified nucleic acid vector) that has been altered, modified or engineered such that it contains greater, fewer or different nucleic acid sequences than those included in the native or natural nucleic acid molecule from which the recombinant vector was derived. Preferably, the recombinant vector includes a coaX, coaA, pan or ilv gene or recombinant nucleic acid molecule including such coaX, coaA, pan or ilv gene, operably linked to regulatory sequences, for example, promoter sequences, terminator sequences and/or artificial ribosome binding sites (RBSs), as defined herein.

The phrase "operably linked to regulatory sequence(s)" means that the nucleotide sequence of the nucleic acid molecule or gene of interest is linked to the regulatory sequence(s) in a manner which allows for expression (e.g., enhanced, increased, constitutive, basal, attenuated, decreased or repressed expression) of the

nucleotide sequence, preferably expression of a gene product encoded by the nucleotide sequence (e.g., when the recombinant nucleic acid molecule is included in a recombinant vector, as defined herein, and is introduced into a microorganism).

The term "regulatory sequence" includes nucleic acid sequences which affect (e.g., modulate or regulate) expression of other nucleic acid sequences. In one embodiment, a regulatory sequence is included in a recombinant nucleic acid molecule or recombinant vector in a similar or identical position and/or orientation relative to a particular gene of interest as is observed for the regulatory sequence and gene of interest as it appears in nature, e.g., in a native position and/or orientation. For example, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector 10 operably linked to a regulatory sequence which accompanies or is adjacent to the gene of interest in the natural organism (e.g., operably linked to "native" regulatory sequences, for example, to the "native" promoter). Alternatively, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence which accompanies or is adjacent to another (e.g., a 15 different) gene in the natural organism. Alternatively, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence from another organism. For example, regulatory sequences from other microbes (e.g., other bacterial regulatory sequences, bacteriophage regulatory sequences and the like) can be operably linked to a particular gene of interest. 20

In one embodiment, a regulatory sequence is a non-native or non-naturallyoccurring sequence (e.g., a sequence which has been modified, mutated, substituted, derivatized, deleted including sequences which are chemically synthesized). Preferred regulatory sequences include promoters, enhancers, termination signals, anti-termination signals and other expression control elements (e.g., sequences to which repressors or inducers bind and/or binding sites for transcriptional and/or translational regulatory proteins, for example, in the transcribed mRNA). Such regulatory sequences are described, for example, in Sambrook, J., Fritsh, E. F., and Maniatis, T. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989. Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in a microorganism (e.g., constitutive promoters and strong constitutive promoters), those which direct inducible expression of a nucleotide sequence in a microorganism (e.g., inducible promoters, for example, xylose inducible promoters) and those which attenuate or repress expression of a nucleotide sequence in a microorganism (e.g., attenuation signals or repressor sequences). It is also within the scope of the present invention to regulate expression of a gene of interest by removing or deleting regulatory sequences. For

25

example, sequences involved in the negative regulation of transcription can be removed such that expression of a gene of interest is enhanced.

In one embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes a nucleic acid sequence or gene that encodes at least one bacterial gene product (e.g., a pantothenate biosynthetic enzyme, an isoleucinevaline biosynthetic enzyme, or a CoaA biosynthetic enzyme, for example CoaA or CoaX) operably linked to a promoter or promoter sequence. Preferred promoters of the present invention include Bacillus promoters and/or bacteriophage promoters (e.g., bacteriophage which infect Bacillus). In one embodiment, a promoter is a Bacillus promoter, preferably a strong *Bacillus* promoter (e.g., a promoter associated with a biochemical housekeeping gene in Bacillus or a promoter associated with a glycolytic pathway gene in Bacillus). In another embodiment, a promoter is a bacteriophage promoter. In a preferred embodiment, the promoter is from the bacteriophage SP01. In a particularly preferred embodiment, a promoter is selected from the group consisting of 15 P_{15} , P_{26} or P_{veg} , for example, the promoters set forth in SEQ ID NO:39, SEQ ID NO:40 or SEQ ID NO:41. Additional preferred promoters include tef (the translational elongation factor (TEF) promoter) and pyc (the pyruvate carboxylase (PYC) promoter), which promote high level expression in Bacillus (e.g., Bacillus subtilis). Additional preferred promoters, for example, for use in Gram positive microorganisms include, but are not limited to, the amy E promoter or phage SP02 promoters. Additional preferred promoters, for example, for use in Gram negative microorganisms include, but are not limited to tac, trp, tet, trp-tet, lpp, lac, lpp-lac, lacIq, T7, T5, T3, gal, trc, ara, SP6, λ-P_R or $\lambda - P_L$.

In another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes a terminator sequence or terminator sequences (e.g., transcription terminator sequences). The term "terminator sequences" includes regulatory sequences which serve to terminate transcription of a gene. Terminator sequences (or tandem transcription terminators) can further serve to stabilize mRNA (e.g., by adding structure to mRNA), for example, against nucleases.

20

25

30

35

In yet another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes sequences which allow for detection of the vector containing said sequences (i.e., detectable and/or selectable markers), for example, sequences that overcome auxotrophic mutations, for example, ura3 or ilvE, fluorescent markers, and/or colorimetric markers (e.g., lacZ/β-galactosidase), and/or antibiotic resistance genes (e.g., amp or tet).

In yet another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes an artificial ribosome binding site (RBS). The term "artificial ribosome binding site (RBS)" includes a site within an mRNA molecule (e.g., coded within DNA) to which a ribosome binds (e.g., to initiate translation) which differs from a native RBS (e.g., a RBS found in a naturally-occurring gene) by at least one nucleotide. Preferred artificial RBSs include about 5-6, 7-8, 9-10, 11-12, 13-14, 15-16, 17-18, 19-20, 21-22, 23-24, 25-26, 27-28, 29-30 or more nucleotides of which about 1-2, 3-4, 5-6, 7-8, 9-10, 11-12, 13-15 or more differ from the native RBS (e.g., the native RBS of a gene of interest). Preferably, nucleotides which differ are substituted. such that they are identical to one or more nucleotides of an ideal RBS (e.g., SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47 or SEQ ID NO:48), when optimally aligned for comparisons. Artificial RBSs can be used to replace the naturallyoccurring or native RBS associated with a particular gene. Artificial RBSs preferably increase translation of a particular gene. Preferred artificial RBSs (e.g., RBSs for increasing the translation of panB, for example, of B. subtilis panB) are depicted in Table IA (e.g., SEQ ID NO:49 and SEQ ID NO:50).

Table 1A: Preferred panB Ribosome Binding Sites

```
20 10 20

-----AGAAAGGAGGTGA ideal RBS (SEQ ID NO:44)

CCCTCT-AG-AAGGAGGAGAAAACATG RBS1 (SEQ ID NO:49)

CCCTCT-AG--AGGAGGAGAAAACATG RBS2 (SEQ ID NO:50)

25 TAAACAT-G--AGGAGGAGAAAACATG panB native RBS (SEQ ID NO:42)
```

Additional preferred artificial RBSs (e.g., RBSs for increasing the translation of panD, for example, of B. subtilis panD) are depicted in Table 1B (e.g., SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53 and SEQ ID NO:54).

Table 1B: Preferred panD Ribosome Binding Sites

	10 20	
	. 1 1	
35	CTAGAAAAGGAGGAATTTAAATG	pAN423 RBS (SEQ ID NO:88)
,دد	TTAAGAAAGGAGGTGANNNATG	ideal RBS (SEQ ID NO:45)
40	TTAGAAAGGAGGATTTAAATATG TTAGAAAGGAGGTTTAATTAA	new design A (SEQ ID NO:51) new design B (SEQ ID NO:52) new design C1 (SEQ ID NO:53) new design C2 (SEQ ID NO:54) ideal RBS (SEQ ID NO:46)

Additional preferred artificial RBSs (e.g., RBSs for increasing the translation of panD, for example, of B. subtilis panD) are depicted in Table 1C (e.g., SEQ ID NO:55, SEQ ID NO:56 and SEQ ID NO:57). The predicted amino acid sequence at the C-terminus of the PanC protein is shown. The start codon for PanD translation is underlined.

Table 1C: Additional Preferred panD Ribosome Binding Sites

			10		20	כ					
10			L			<u> </u>					
		A	GAA	AGG	AGG	TGA	NNN	NNN	N	ATG	ideal RBS (SEQ ID NO:47)
	ATT	CGA	GAA	ATG	GAG	AGA	ATA	TAA	т	ATG	native panD RBS (SEQ ID NO:43)
15	Ile	Arg	Glu	Met	Glu	Arg	Ile	*		Met	SEQ ID NO:89
		A	GAA	AGG	AGG	TGA	NNN	NNN	N	ATG	ideal RBS (SEQ ID NO:47)
							ATA	TAA	T	ATG	NDI (SEQ ID NO:55)
20	Ile	Arg	Glu	Arg	Arg	*				Met	SEQ ID NO:90
	ATT	CGA	GAA	AGG	AGG	TGA	ATA	ATA	_	ATG	NDII (SEQ ID NO:56)
	Ile	Arg	Glu	Arg	Arg	*	1 .			Met	SEQ ID NO:90
	ATT	CGT	AGA	AAG	GAG	GTG	AAT	TAA	т	ATG	NDIII (SEQ ID NO:57)
25	Ile	Arg	Arg	Lys	Glu	Val	Asn	*		Met	SEQ ID NO:91
			AGA	AAG	GAG	GTG	ANN	NNN	N	ATG	ideal RBS (SEQ ID NO:48)

Accordingly, in one embodiment, a vector of the present invention includes an artificial RBS as set forth in SEQ ID NO:49 or SEQ ID NO:50. In another embodiment, a vector of the present invention includes an artificial RBS as set forth in SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53 or SEQ ID NO:54. In yet another embodiment, a vector of the present invention includes an artificial RBS as set forth in SEQ ID NO:55, SEQ ID NO:56 or SEQ ID NO:57.

In another embodiment, a recombinant vector of the present invention includes sequences that enhance replication in bacteria (e.g., replication-enhancing sequences). In one embodiment, replication-enhancing sequences are derived from E. coli. In another embodiment, replication-enhancing sequences are derived from pBR322 (e.g., sequences included within the pBR322 derived portion of any of the pAN vectors as set forth in the Figures, i.e., the Not I-Not I sequences from about 5.0 kB to 9.0 kB of the vector depicted in Figure 3A).

15

20

In yet another embodiment, a recombinant vector of the present invention includes antibiotic resistance genes. The term "antibiotic resistance genes" includes sequences which promote or confer resistance to antibiotics on the host organism (e.g., Bacillus). In one embodiment, the antibiotic resistance genes are selected from the group consisting of cat (chloramphenicol resistance) genes, tet (tetracycline resistance) genes, erm (erythromycin resistance) genes, neo (neomycin resistance) genes and spec (spectinomycin resistance) genes. Recombinant vectors of the present invention can further include homologous recombination sequences (e.g., sequences designed to allow recombination of the gene of interest into the chromosome of the host organism). For example, amyE sequences can be used as homology targets for recombination into the host chromosome.

Preferred vectors of the present invention include, but are not limited to, vectors set forth in Figures 2-15, 17, 19, 22, 25 and 26. It will further be appreciated by one of skill in the art that the design of a vector can be tailored depending on such factors as the choice of microorganism to be genetically engineered, the level of expression of gene product desired and the like.

IX. Isolated Proteins

Another aspect of the present invention features isolated proteins (e.g., isolated pantothenate biosynthetic enzymes and/or valine-isoleucine biosynthetic enzymes and/or isolated CoA biosynthetic enzymes, for example isolated CoaA or CoaX). In one embodiment, proteins (e.g., isolated pantothenate biosynthetic enzymes and/or valineisoleucine biosynthetic enzymes and/or isolated CoaA biosynthetic enzymes, for example isolated CoaA or CoaX) are produced by recombinant DNA techniques and can be isolated from microorganisms of the present invention by an appropriate purification 25 scheme using standard protein purification techniques. In another embodiment, proteins are synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein (e.g., an isolated or purified biosynthetic enzyme) is substantially free of cellular material or other contaminating proteins from the microorganism from which the protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. In one embodiment, an isolated or purified protein has less than about 30% (by dry weight) of contaminating protein or chemicals, more preferably less than about 20% of contaminating protein or chemicals, still more preferably less than about 10% of contaminating protein or chemicals, and most preferably less than about 5% contaminating protein or chemicals.

30

35

In a preferred embodiment, the protein or gene product is derived from Bacillus (e.g., is Bacillus-derived). The term "derived from Bacillus" or "Bacillus-derived" includes a protein or gene product which is encoded by a Bacillus gene. Preferably, the gene product is derived from a microorganism selected from the group consisting of Bacillus subtilis, Bacillus lentimorbus, Bacillus lentus, Bacillus firmus, Bacillus pantothenticus, Bacillus amyloliquefaciens, Bacillus cereus, Bacillus circulans, Bacillus coagulans, Bacillus licheniformis, Bacillus megaterium, Bacillus pumilus, Bacillus thuringiensis, and other Group 1 Bacillus species, for example, as characterized by 16S rRNA type (Priest, supra). In another preferred embodiment, the protein or gene product is derived from Bacillus brevis or Bacillus stearothermophilus. In another 10 preferred embodiment, the protein or gene product is derived from a microorganism selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus. In a particularly preferred embodiment, the protein or gene product is derived from Bacillus subtilis (e.g., is Bacillus subtilis-derived). The term "derived from Bacillus subtilis" or "Bacillus 15 subtilis-derived" includes a protein or gene product which is encoded by a Bacillus subtilis gene. In yet another preferred embodiment, the protein or gene product is encoded by a Bacillus gene homologue (e.g., a gene derived from a species distinct from Bacillus but having significant homology to a Bacillus gene of the present invention, for 20 example, a Bacillus pan gene or Bacillus ilv gene).

Included within the scope of the present invention are bacterial-derived proteins or gene products and/or Bacillus-derived proteins or gene products (e.g., B. subtilisderived gene products) that are encoded by naturally-occurring bacterial and/or Bacillus genes (e.g., B. subtilis genes), for example, the genes identified by the present inventors, for example, Bacillus or B. subtilis coaX genes, coaA genes, pan genes and/or ilv genes. Further included within the scope of the present invention are bacterial-derived proteins or gene products and/or Bacillus-derived proteins or gene products (e.g., B. subtilisderived gene products) that are encoded bacterial and/or Bacillus genes (e.g., B. subtilis genes) which differ from naturally-occurring bacterial and/or Bacillus genes (e.g., B. subtilis genes), for example, genes which have nucleic acids that are mutated, inserted or deleted, but which encode proteins substantially similar to the naturally-occurring gene products of the present invention. For example, it is well understood that one of skill in the art can mutate (e.g., substitute) nucleic acids which, due to the degeneracy of the genetic code, encode for an identical amino acid as that encoded by the naturallyoccurring gene. Moreover, it is well understood that one of skill in the art can mutate (e.g., substitute) nucleic acids which encode for conservative amino acid substitutions. It is further well understood that one of skill in the art can substitute, add or delete

amino acids to a certain degree without substantially affecting the function of a gene product as compared with a naturally-occurring gene product, each instance of which is intended to be included within the scope of the present invention.

In a preferred embodiment, an isolated protein of the present invention (e.g., an isolated pantothenate biosynthetic enzyme and/or an isolated isoleucine-valine biosynthetic enzyme and/or an isolated CoaA biosynthetic enzymes, for example isolated CoaA or CoaX) has an amino acid sequence shown in SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87. In other embodiments, an isolated protein of the present invention is a homologue of the at least one of the proteins set forth as SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87 (e.g., comprises an amino acid sequence at least about 30-40% identical, preferably about 40-50% identical, more preferably about 50-60% identical, and even more preferably about 60-70%, 70-80%, 80-90%, 90-95% or 15 more identical to the amino acid sequence of SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87, and has an activity that is substantially similar to that of the protein encoded by the amino acid sequence of SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87, respectively.

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = # of identical positions/total # of positions x 100), preferably taking into account the number of gaps and size of said gaps necessary to produce an optimal alignment.

The comparison of sequences and determination of percent homology between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264-68, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-77. Such

10

15

20

an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul et al. (1990) J. Mol. Biol. 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997) Nucleic Acids Research 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See http://www.ncbi.nlm.nih.gov. Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller (1988) Comput Appl Biosci. 4:11-17. Such an algorithm is incorporated into the ALIGN program available. for example, at the GENESTREAM network server, IGH Montpellier, FRANCE (http://vega.igh.cnrs.fr) or at the ISREC server (http://www.ch.embnet.org). When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

In another preferred embodiment, the percent homology between two amino acid sequences can be determined using the GAP program in the GCG software package (available at http://www.gcg.com), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 12, 10, 8, 6, or 4 and a length weight of 2, 3, or 4. In yet another preferred embodiment, the percent homology between two nucleic acid sequences can be accomplished using the GAP program in the GCG software package (available at http://www.gcg.com), using a gap weight of 50 and a length weight of 3.

25

30

35

X. Biotransformations and Bioconversions

Another aspect of the present invention includes biotransformation processes which feature recombinant microorganisms (e.g., mutant microorganisms) and/or isolated CoA, pantothenate or isoleucine-valine biosynthetic enzymes described herein. The term "biotransformation process", also referred to herein as "bioconversion processes", includes biological processes which result in the production (e.g., transformation or conversion) of any compound (e.g., intermediate or product) which is upstream of a CoA, pantothenate or isoleucine-valine biosynthetic enzyme to a compound (e.g., substrate, intermediate or product) which is downstream of said CoA, pantothenate or isoleucine-valine biosynthetic enzyme.

In one embodiment, the invention features a biotransformation process for the production of a panto-compound comprising contacting a microorganism which overexpresses at least one pantothenate biosynthetic enzyme with at least one appropriate substrate or precursor under conditions such that said panto-compound is produced and recovering said panto-compound. In a preferred embodiment, the invention features a biotransformation process for the production of pantoate comprising contacting a microorganism which overexpresses ketopantoate reductase (the panE gene product) with an appropriate substrate (e.g., ketopantoate) under conditions such that pantoate is produced and recovering said pantoate. In another preferred embodiment, the invention features a biotransformation process for the production of pantothenate comprising contacting a microorganism which overexpresses ketopantoate reductase and pantothenate synthetase with appropriate substrates (e.g., ketopantoate and β-alanine) under conditions such that pantothenate is produced and recovering said pantothenate. In yet another preferred embodiment, the invention features a biotransformation process for the production of pantothenate comprising contacting a microorganism which overexpresses ketopantoate hydroxymethyltransferase, ketopantoate reductase and pantothenate synthetase with appropriate substrates (e.g., α -ketoisovalerate and β alanine) under conditions such that pantothenate is produced and recovering said pantothenate. Preferred recombinant microorganisms for carrying out the abovedescribed biotransformations include pantothenate kinase mutants. Conditions under which pantoate or pantothenate are produced can include any conditions which result in the desired production of pantoate or pantothenate, respectively.

In yet another embodiment, the present invention includes a method of producing β -alanine that includes culturing a microorganism which overexpresses aspartate- α -decarboxylase under conditions such that β -alanine is produced. Preferably, the aspartate- α -decarboxylase-overexpressing microorganism has a mutation in a nucleic acid sequence encoding a pantothenate biosynthetic enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase and pantothenate synthetase.

The invention further features a method of producing β -alanine that includes contacting a composition comprising aspartate with an isolated *Bacillus* aspartate- α -decarboxylase enzyme under conditions such that β -alanine is produced (e.g., an in vitro synthesis method).

The microorganism(s) and/or enzymes used in the biotransformation reactions are in a form allowing them to perform their intended function (e.g., producing a desired compound). The microorganisms can be whole cells, or can be only those portions of the cells necessary to obtain the desired end result. The microorganisms can be

25

30

suspended (e.g., in an appropriate solution such as buffered solutions or media), rinsed (e.g., rinsed free of media from culturing the microorganism), acetone-dried, immobilized (e.g., with polyacrylamide gel or k-carrageenan or on synthetic supports, for example, beads, matrices and the like), fixed, cross-linked or permeablized (e.g., have permeablized membranes and/or walls such that compounds, for example, substrates, intermediates or products can more easily pass through said membrane or wall).

Purified or unpurified CoA biosynthetic enzyme(s) (e.g., CoaA and/or CoaX), pantothenate biosynthetic enzyme(s) and/or valine-isoleucine biosynthetic enzyme(s) can also be used in biotransformation reactions. The enzyme can be in a form that allows it to perform its intended function (e.g., obtaining the desired compound). For example, the enzyme can be in free form or immobilized. Purified or unpurified CoA biosynthetic enzyme(s), pantothenate biosynthetic enzyme(s) and/or valine-isoleucine biosynthetic enzyme(s) can be contacted in one or more in vitro reactions with appropriate substrate(s) such that the desired product is produced.

With respect to at least the above-described methodologies (e.g., the production methodologies of the present invention), at least one aspect of the invention features the folowing: embodiments is which the methods do not use microorganisms of the genus Corynebacterium and/or microorganisms of the genus Escherichia; embodiments in which the methods do not use microorganisms selected from the group consiting of Escherichia coli and Corynebacterium glutamicum; embodiments in which the methods do not use gram negative microorganisms; embodiments in which the microorganisms utilized do not include, express or produce nucleic acid molecules, genes or proteins (e.g., biosynthetic emzymes) derived from microorganisms of the genus

25 Corynebacterium and/or microorganisms of the genus Escherichia; embodiments in which the microorganisms to not include, express or produce nucleic acid molecules, genes or proteins (e.g., biosynthetic emzymes) derived from microorganisms selected from the group consisting of Escherichia coli and Corynebacterium glutamicum.

30 XI. Screening Assays

Because CoA is an essential factor in bacteria, proteins (e.g., enzymes) involved in the biosynthesis of CoA provide valuable tools in the search for novel anti-biotics. In particular, the CoaX protein is a valuable target for identifying bacteriocidal compounds because it bears no resemblance in primary sequence to mammalian pantothenate kinase enzymes. Accordingly, the present invention also provides a method (also referred to herein as a "screening assay") for identifying modulators, i.e., candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other drugs)

10

15

20

30

35

which bind to CoaX, or have a stimulatory or inhibitory effect on, for example, coaX expression or CoaX activity.

In one embodiment, the invention provides assays for screening candidate or test compounds which are capable of binding to CoaX proteins or a biologically active 5 portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which modulate the activity of CoaX proteins or biologically active portions thereof. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'onebead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) Anticancer Drug Des. 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt et al. (1993) Proc. Natl. Acad. Sci. U.S.A. 90:6909; Erb et al. (1994) Proc. Natl. Acad. Sci. USA 91:11422; Zuckermann et al. (1994). J. Med. Chem. 37:2678; Cho et al. (1993) Science 261:1303; Carrell et al. (1994) Angew. Chem. Int. Ed. Engl. 33:2059; Carell et al. (1994) Angew. Chem. Int. Ed. Engl. 33:2061; and in Gallop et al. (1994) J. Med. Chem. 37:1233. Libraries of compounds may be presented in solution (e.g., Houghten (1992) Biotechniques 13:412-421), or on beads (Lam (1991) Nature 354:82-84), chips (Fodor (1993) Nature 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull et al. (1992) Proc Natl Acad Sci USA 89:1865-1869) or on phage (Scott and Smith (1990) Science 249:386-390); (Devlin (1990) Science 249:404-406); (Cwirla et al. (1990) Proc. Natl. Acad. Sci. 87:6378-6382); (Felici (1991) J. Mol. Biol. 222:301-310); (Ladner supra.).

In one embodiment, an assay is a microorganism-based assay in which a recombinant microorganism which expresses a CoaX protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate CoaX activity is determined. Determining the ability of the test compound to modulate CoaX activity can be accomplished by monitoring, for example, intracellular phosphopanthoate or CoA concentrations or secreted pantothenate concentrations (as compounds that inhibit CoaX will result in a buildup of pantothenate in the test microorganism). CoaX substrate can be labeled with a radioisotope or enzymatic label such that modulation of CoaX activity can be determined by detecting a conversion of labeled substrate to intermediate or product. For example, CoaX substrates can be

labeled with ³²P, ¹⁴C, or ³H, either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Determining the ability of a compound to modulate CoaX activity can alternatively be determined by detecting the induction of a reporter gene (comprising a CoA-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a CoA-regulated cellular response.

In yet another embodiment, a screening assay of the present invention is a cell-free assay in which a CoaX protein or biologically active portion thereof is contacted with a test compound *in vitro* and the ability of the test compound to bind to or modulate the activity of the CoaX protein or biologically active portion thereof is determined. In a preferred embodiment, the assay includes contacting the CoaX protein or biologically active portion thereof with known substrates to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to modulate enzymatic activity of the CoaX on its substrates.

10

15 .

20

25

30

35

Screening assays can be accomplished in any vessel suitable for containing the microorganisms, proteins, and/or reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either CoaX protein or a recombinant microorganism expressing CoaX protein to facilitate separation of products and/or substrates, as well as to accommodate automation of the assay. For example, glutathione-S-transferase/CoaX fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates. Other techniques for immobilizing proteins on matrices (e.g., biotin-conjugation and streptavidin immobilization or antibody conjugation) can also be used in the screening assays of the invention.

In another embodiment, modulators of CoaX expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of coaX mRNA or CoaX polypeptide in the cell is determined. The level of expression in the presence of the candidate compound is compared to the level of expression in the absence of the candidate compound (or to a suitable control, for example, an appropriate buffer control or standard). The candidate compound can then be identified as a modulator of coaX mRNA or CoaX polypeptide expression based on this comparison.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an CoaX modulating agent identified as described herein (e.g., an anti-bactericidal

compound) can be used in an infectious animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents, patent applications (including U.S. Patent Application Serial No. 09/400,494, filed September 21, 1999 (pending), provisional U.S. Patent Application Serial No. 60/210,072, filed June 7, 2000, provisional U.S. Patent Application Serial No. 60/221,938, filed July 28, 2000 and provisional U.S. Patent Application Serial No. 60/227,860, filed August 24, 2000, to which this application relates) and published patent applications cited throughout this application are incorporated herein by reference.

EXAMPLES

General Methodology:

Strains. Bacillus subtilis strains of the present invention are generally derived from either of two strains. The first is variously named "168", "1A1", or "RL-1". The genotype is trpC2. This strain was derived from the wild type "Marburg" strain by mutagenesis and has been the basis of much of the molecular biology work done on B. subtilis. The second strain is PY79, a prototrophic derivative of 168 that was made Trp⁺ by transduction from the wild type strain W23.

Media. Standard minimal medium for B. subtilis is comprised of 1 x Spizizen salts and 0.5% glucose. Standard solid "rich medium" is Tryptone Blood Agar Broth (Difco), and standard liquid "rich medium" is VY, a mixture of veal infusion broth and yeast extract. For testing production of pantothenate in liquid test tube cultures, an enriched form of VY, called "Special VY" or "SVY" is used. For batch fermentations, SVYG and PFMG are used. The compositions of these media are given below.

15 VY, a rich liquid medium: 25 g Difco Veal Infusion Broth, 5 g Difco Yeast Extract, 1L water (autoclave).

TBAB, a rich solid medium: 33 g Difco Tryptone Blood Agar Broth, 1L water (autoclave).

MIN, a minimal medium: 100 ml 10 x Spizizen salts; 10 ml 50% glucose; 2 ml 10% arginine HCl*; 10 ml 0.8% tryptophan**; water to 1 liter. (*In some cases, arginine is omitted or replace by sodium glutamate at 0.04% final concentration. In general, B. subtilis grows faster in minimal medium when certain amino acids, such as arginine, glutamine, glutamate, or proline, are added as an auxiliary nitrogen source.) (**For strains that are tryptophan auxotrophs, tryptophan is routinely added to most minimal media.)

10 x Spizizen Salts: 174 g K₂HPO₄·3H₂O; 20 g (NH₄)₂SO₄; 60 g KH₂PO₄; 10 g Na₃Citrate·2H₂O; 2 g MgSO₄·7H₂O; water to 993 mls; then add 3.5 ml FeCl₃ solution and 3.5 ml Trace Elements solution.

FeCl₃Solution: 4 g FeCl₃·6H₂O; 197 g Na₃Citrate·2H₂O; water to 1 liter (filter 30 sterilize)

Trace Elements Solution: 0.15 g Na₂MoO₄·2H₂O; 2.5 g H₃BO₃; 0.7 g CoCl₂·6H₂O; 0.25 g CuSO₄·5H₂O; 1.6 g MnCl₂·4H₂O; 0.3 g ZnSO₄·7H₂O; water to 1 liter (filter sterilize).

SVY, Special VY, a supplemented* rich medium for testing pantothenate production in test tube cultures: 25 g Difco Veal Infusion Broth; 5 g Difco yeast extract; 5 g sodium glutamate; 2.7 g ammonium sulfate; 740 ml water (autoclave); add 200 ml 1 M potassium phosphate, pH 7.0; 60 ml 50% glucose. (*For testing pantothenate production in liquid SVY test tube cultures, Na α-ketoisovalerate and/or β-alanine can be added to a concentration of 5 g/L from filter-sterilized stocks.)

PFMG, a yeast extract based medium used in fermentors: 20 g Amberex
1003™ yeast extract; 5 g sodium glutamate, 2 g ammonium sulfate; 5 g tryptophan; 10 g
KH₂PO4; 20 g K₂HPO₄·3H₂O; 1 g MgCl₂·6H₂O; 0.1 g CaCl₂·2H₂O; 1 g sodium citrate;
10 0.01 g FeSO₄·7H₂O; 1 ml trace elements solution; 20 g glucose; add water to 1 L.
Glucose or other sugars are fed as needed. Feed solutions can contain minerals, defined or food grade nutrients.

PF, a chemically defined pantothenate free medium for testing pantothenate auxotrophy: 100 ml 10 x Spizizen Salts; 100 ml 1 x Difco Pantothenate Assay Medium; 10 ml 50% glucose; water to 1 liter.

For pantothenate auxotrophs, 1 mM Na pantothenate is added to both minimal and rich media, since there is generally not enough pantothenate in rich media to support B. subtilis pan mutants. Amino acids are at 100 mg per liter, when used.

Selection for antibiotic resistance is done with 5 mg/L chloramphenicol, 100 mg/L spectinomycin HCl, 15 mg/L tetracycline HCl, or 1 mg/L erythromycin plus 25 mg/L lincomycin.

Pantothenate Assays: Biological assay. The indicator organism, Lactobacillus plantarum, requires pantothenate for growth, and responds to low concentrations (μg/L).
 Thus, using serial dilutions, a wide range of concentrations can be assayed.
 Commercially available medium (e.g., Pantothenate Assay Medium (PAM), Difco), can be used. However, Difco PAM supplemented with pantothenate does not support growth to the same level as obtainable using a fresh-mixed version of Pantothenate Assay Medium (FM-PAM), made up of the individual components as specified by
 Difco, which is accordingly, routinely used instead of the commercial product.

Before assaying *B. subtilis* culture supernatants, the *B. subtilis* cells must be either removed or killed. *B. subtilis* culture supernatants give approximately the same pantothenate titer when the supernatants are autoclaved as when they are sterile filtered. Accordingly, routine procedures involve autoclaving samples for 5 minutes prior to the biological assay.

WO 01/21772 PCT/US00/25993

- 61 -

Pantothenate Assays: HPLC assay. Pantothenic acid production is measured by HPLC with a detector wavelength of 197 nm and a reference at 450 nm. The procedure is a modification of one recommended by Hewlett-Packard for water soluble vitamins. Samples of culture broth are diluted into an equal volume of 60% acetronitrile (ACN), centrifuged and filtered. Typically a further 10-fold dilution before analysis brings the final dilution to 20-fold. Higher concentrations of product are diluted further. Compounds are separated on a C18 Phenomenex 5µ Aqua 250 x 4.6 mm column with 5% acetronitrile (ACN) in 50 mM Na phosphate buffer at pH 2.5. An ACN gradient from 5% to 95% washes the column between every sample. The area of the pantothenate peak is proportional to the concentration between 5 to 1000 mg/L. Other panto-compounds are also separated and quantitated by this method.

Amino Acid Analysis: HPLC assay. Amino acids present in the fermentation medium and throughout the fermentation are measured by HPLC with a detector wavelength of 338 nm and a reference at 390 nm. The procedure is a modification of 15 one recommended by Hewlett-Packard for amino acid analysis. Samples of culture broth are prepared identically as for the panto-compound analysis. Compounds are separated on a C18 Hypersil 5µ ODS 200 x 2.1 mm column. Solvent A is 20 mM Na acetate buffer at pH 7.2. Solvent B contains 40% ACN and 40% methanol. A gradient from 100% Solvent A to 100% Solvent B separates amino acids and washes the column between every sample.

Batch Fermentations. Pantothenate producing strains are grown in stirred tank fermentors, for example, in CF3000 Chemap 14 liter vessels with 10 liter working volumes. Computer control and data collection is by commercial software, for example, B. Braun Biotech MFCS software. Fermentations can be batch processes but are preferably sugar-limited, fed batch processes. Some media components (e.g. of SVYG and PMFG) are added to the fermentor and sterilized in place. Portions of the media are sterilized separately and added to the fermentors aseptically. This procedure is well known to those familiar with the art. Additional nitrogen sources in feeds are sterilized separately and added to the carbon source after cooling.

The initial sugar in the medium is consumed in approximately 6 hours. Afterwards, glucose or other sugars are fed with the possible addition of minerals, and defined or food grade nutrients. Alternatively, feeds are scheduled based on a consensus profile of nutritional requirements from samples taken from earlier fermentations.

20

25

After inoculation, agitation is set at a relatively low speed, e.g. 200 rpm. When the dissolved oxygen (pO2) falls to 30%, computer control automatically adjusts the agitation to maintain a dissolved oxygen concentration between 25 and 30% pO2.

EXAMPLE I: Enhanced Production of a Panto-Compound Using Bacteria Overexpressing panBCD Gene Products.

This Example describes the cloning of the B. subtilis panBCD operon and the generation of microorganisms overexpressing the panBCD gene products.

To clone the B. subtilis panBCD operon, a plasmid library of B. subtilis GP275 (a derivative of 168) genomic DNA was transformed in E. coli BM4062 (birAis), and temperature resistant clones were selected at 42°C. By comparison of restriction maps to the genome sequence, one particular clone was deduced to contain the B. subtilis birA gene and the adjacent panBCD genes. This plasmid was named pAN201.

To overexpress the panBCD operon and produce pantothenate, the native promoter of the panBCD operon was replaced by either of two strong, constitutive promoters derived from the B. subtilis bacteriophage SP01. These two promoters are named P_{26} and P_{15} . In addition, either of two artificial ribosome binding sites (RBSs) were used to replace the native panB RBS. These two artificial RBSs (set forth as SEQ ID NO:49 and SEQ ID NO:50) were predicted to increase translation of panBCD; their sequences are shown in Table 1A. Three such engineered panBCD expression cassettes were built into circular plasmids capable of replicating in E. coli. Other features of the plasmids include a strong rho-independent transcription terminator from the E. coli ribosomal RNA transcription unit, called T1T2, a Gram-positive chloramphenicol resistance gene (cat), derived from pC194, and a pair of NotI restriction sites at the junctions between the E. coli replicon and the segment intended for integration into B. subtilis. Three plasmids of this series, pAN004, pAN005, and pAN006 were constructed. pAN004 contains the P26 promoter, RBS1, and a low copy £. coli replicon. pAN005 contains the P_{15} promoter, which in our experience is not as strong as P_{26} , RBS1, and the low copy replicon. pAN006 contains the P26 promoter, RBS2, and a medium copy replicon. 30

The three panBCD expression cassettes contained in the above-mentioned three plasmids were all ligated to a DNA fragment consisting of sequences that naturally occur immediately upstream from the native panB gene and integrated in single copy by homologous recombination into the panBCD locus of B. subtilis strains RL-1 and PY79, replacing the wild-type operon. This was accomplished in two steps. First a deletionsubstitution that replaced about two thirds of the panB coding region with a Gram-

35

positive spectinomycin resistance gene (*spec*) was integrated at *panB* to yield Spec, pantothenate auxotrophs. These intermediate strains were than transformed with the *panBCD* expression cassettes of pAN004, pAN005, and pAN006 after ligating them to a DNA fragment containing chromosomal sequences just upstream of *panB*. Selection of the incoming cassette was for pantothenate prototrophy. The resulting strains were named PA221, PA222 and PA223 (from RL-1), and PA235, PA232 and PA233 (from PY79), respectively. An example of a plasmid that contains the joined upstream sequence that is in the integrated strain in PA221 is pAN240 (see Figure 2). The nucleotide sequence of pAN240 is set forth as SEQ ID NO:76.

Polymerase chain reaction using appropriate primers was used to verify the correct chromosomal structures of these engineered strains. When extracts of strain PA221 were examined by SDS-PAGE, two proteins were found to be overexpressed. One protein had an apparent molecular weight of 29,000 and the other protein appeared to be 39,000 daltons. The 29,000 dalton bands is presumably PanB (predicted molecular weight of 29,761). The larger protein band presumably represents PanC (predicted size 31,960 daltons).

The ability of these strains to produce pantothenate in test tube cultures was assessed as follows. Each strain was grown in SVY medium supplemented with 5 g/L α-ketoisovalerate (α-KIV) and 5 g/L β-alanine, to ensure that these precursors were not limiting. Culture supernatants were autoclaved and assayed using the bioassay. Relative to the parent strains, RL-1 and PY79, the engineered strains produced about 8-to 30-fold more pantothenate, attaining 1 g/L pantothenate in some cases.

PHICOCOLD AND 012177242 L

Table 2. Production of pantothenate by engineered B. subtilis strains in liquid test tube cultures grown in SVY medium with $5 \, \text{g/L} \, \alpha$ -KIV and $5 \, \text{g/L} \, \beta$ -alanine.

		المحاصر بالمحاصر والمحاصر والم			
Strain	Promoter	RBS at panB	[pantothenate] mg/L		
RL-1	Native	Native	30		
PA221	P ₂₆	RBS1	990 790		
PA222	P,,	RBS1	250 250		
PA223	P ₂₆	RBS2	790 790		
PY79	Native	Native	40		
PA235	P ₂₆	RBS1	930 ¹ 860		
PA221	P ₂₆	RBS1	1100 1030		
	RL-1 PA221 PA222 PA223 PY79 PA235	RL-1 Native PA221 P ₂₆ PA222 P ₁₅ PA223 P ₂₆ PY79 Native PA235 P ₂₆	Strain Promoter panB RL-1 Native Native PA221 P ₁₆ RBS1 PA222 P ₁₆ RBS1 PA223 P ₂₆ RBS2 PY79 Native Native PA235 P ₂₆ RBS1		

The P_{26} promoter was about 3- to 4-fold more effective than the P_{15} promoter, 5 while RBS1 and RBS2 were roughly equivalent. Plasmids such as pAN004, pAN005, pAN006 can also be recombined as circles into the B. subtilis wild type panBCD locus by Campbell-type (single crossover) integration, selecting for chloramphenicol resistance at 5 mg/L. Strains obtained in this fashion produce about the same amount of 10 pantothenate as strains PA221, PA222, and PA223, respectively. pAN004 containing the P_{26} promoter, RBS1 and a low copy E. coli replicon, is depicted schematically in Figure 3A. The nucleotide sequence of plasmid pAN004 is set forth as SEQ ID NO:93. pAN006 containing the P_{26} promoter, RBS2 and a medium copy E. coli replicon, is depicted schematically in Figure 3B. The nucleotide sequence of plasmid pAN006 is set forth as SEQ ID NO:94. The nucleotide sequence of panBCD is set forth as SEQ ID 15 NO:59 and the predicted amino acid sequences of PanB, PanC and PanD are set forth as SEQ ID NO:24, SEQ ID NO:26 and SEQ ID NO:28, respectively. Methods for manipulating Bacilli are described, for example, in Harwood, C.R. and Cutting, S.M. (editors), Molecular Biological Methods for Bacillus (1990) John Wiley & Sons, Ltd., Chichester, England, the content of which is incorporated herein by reference. 20

EXAMPLE II: Enhanced Production of a Panto-Compound Using Bacteria Overexpressing the *panE1* Gene Product – Ketopantoate Reductase.

This Example describes the cloning of the *B. subtilis panE1* gene and the generation of microorganisms overexpressing the *panE1* gene product.

Pan B. subtilis strains (e.g., B. subtilis mutants blocked in the synthesis of pantothenic acid) had previously been isolated, one of which was reported to be affected in ketopantoate reductase activity (Baigori et al. (1991) J. Bacteriol. 173:4240-4242). However, the mutations in these strains were incorrectly mapped to the purE-tre interval of the B. subtilis genetic map which does not contain the panE or panBCD genes.

Furthermore as shown below, a panE mutant does not have a Pan phenotype as the ilvC gene product can substitute for the panE gene product in B. subtilis as in other bacterial strains such as E. coli. More recently, the S. typhimuruim panE gene has been located and determined to be allelic to apbA, a gene required for anaerobic purine biosynthesis (Frodyma et al. (1998) J. Biol. Chem. 273:5572-5576). E. coli carries a highly

homologous gene at the same map location. Identification of the panE genes in E. coli and S. typhimurium was complicated by the fact that the ilvC gene product, acetohydroxy acid isomeroreductase, is also capable of carrying out the ketopantoate reductase reaction. As a result, pantothenate auxotrophy is not obtained unless both panE and ilvC are mutated.

To identify the *B. subtilis panE1* gene, the *B. subtilis* genome was searched using the protein sequence of *E. coli* or *S. typhimurium* ApbA (PanE), and two open reading frames were identified having homology to ApbA, named ylbQ and ykpB. These genes were renamed panE1 and panE2, due to their proposed function in pantothenate biosynthesis. Both panE1 and panE2 were cloned as PCR products generated from

RL-1 genomic DNA as a template. Both genes were disrupted by either a spectinomycin resistance gene (spec) or a chloramphenicol resistance gene (cat). The interrupted genes were each integrated by double crossover into PY79 to give PA240 (ΔpanEl::spec) and PA241 (ΔpanE2::cat). Neither of these strains were pantothenate auxotrophs when tested on pantothenate-free (PF) plates, although PA240 containing
ΔpanEl::spec grew slightly more slowly on TBAB without added pantothenate than with a 1 mM pantothenate supplement. By comparison, a ΔpanB::spec strain does not produce single colonies on TBAB, presumably because B. subtilis has no active uptake

system for pantothenate.

It was hypothesized that the B. subtilis gene, ilvC, could function for panE as had been shown for E. coli. Accordingly, the panE1 and panE2 disruptions were introduced

into a strain, CU550, which is reported to be trpC2 ilvC4 leuC124. Both the single

35

panE1 and the double panE1, panE2 disruptants were pantothenate auxotrophs on PF medium.

Table 3. Phenotypes of various panE1 and panE2 mutants on rich and defined media.

Strain	Medium	Growth*:	+ pan
PY79	TBAB	+++	+++
	PF	++	++
PA240	TBAB spec	+	+++
	PF	++	++
PA241	TBAB cam	+++	+++
	PF	++	` ++
CU550	TBAB	+++	+++
•	PF	++	++
PA256	TBAB spec	-	+++
	PF	-	++ ,
PA258	TBAB spec, cam	-	+++
	PF	· · · · · · · · · · · · · · · · · · ·	++

^{*}Each "+" represents about 1 mm of colony diameter after overnight at 37°C.

Thus, mutating both panE1 and ilvC results in pantothenate auxotrophy, while mutating only panE1 does not, similar to what has been reported for E.coli and S. typhimurium.

Next, the quantitative effect of panE1 and panE2 knockouts in a pantothenate overproducing strain (PA235 described herein) was examined. The panE1 and panE2 disruptions were introduced into PA235, either singly or together to produce PA245 (ΔpanE1::spec), PA248 (ΔpanE2::cat) and PA244 (ΔpanE1::cat, ΔpanE2::spec). The effect of each mutation on pantothenate production was then tested in liquid test tube cultures.

10

15

20

25

Table 4. Pantothenate production by PA235 derivatives containing panE1 and panE2 disruptions.

Strain	[pan] mg/L	% of PA235
PA235	990	(100)
PA235	940	95
PA245	59	6
PA245	82	8
PA248	1060	106
PA248	1030	104
PA244	25	3
PA244	50	5

Thus, deletion analysis indicated that the panE1 gene contributes to over 90% of the pantothenate production, while deletion of panE2 did not have a significant effect on pantothenate production. It is therefore concluded that panE1 accounts for most, but not necessarily all, of the ketopantoate reductase activity in B. subtilis. The rest of the ketopantoate reductase activity is predicted to be supplied by ilvC.

Having identified panE1 as an important gene for pantothenate production, increased panE1 expression was tested to determine whether it could enhance pantothenate production in strains such as PA221 or PA235. The panE1 coding sequence was installed downstream of the P26 promoter and RBS2 in a vector, pOTP61, designed to integrate and amplify at either the bpr locus (a non-essential protease gene) or at the locus of the cloned insert. The resulting plasmid, pAN236 (Figure 4) was transformed into PA221, selecting for resistance to tetracycline at 15 mg/L. The nucleotide sequence of pAN236 is set forth as SEQ ID NO:77. One transformant, named PA236 was chosen for further study.

PA236 was shown to overexpress a protein of about 31,000 daltons, which is close to the expected molecular weight of 33,290 daltons for panE1 protein. Briefly, whole cell extracts were prepared from PY79, RL-1, PA221, PA221/pOTP61and PA236 (2 samples). Cell extracts were separated by gel electrophoresis and the gels were coomassie stained to visualize proteins. In cells engineered to overexpress panE (PA236-1 and PA236-2), a band was visible having an approximate molecular weight of ~31,000 daltons (as compared to molecular weight markers). Moreover, PA221 and PA236 expressed increased levels of a ~29,000 dalton band, corresponding to the panB

gene product, and a ~39,000 dalton band, presumably corresponding the panC gene product. Furthermore, E. coli transformed with pAN006 (Figure 3B) expressed bands correlating to the panB and panC gene products and E. coli transfected with PAN236 expressed a ~31,000 dalton band corresponding to the panE gene product.

Next, PA236 was compared to PA221 carrying the empty vector pOTP61 for pantothenate production in liquid test tube cultures supplemented with 5 g/L β -alanine and 5 g/L α -KIV.

Table 5. Effect of overexpression of panE1 and panE2 on pantothenate production by engineered strains in liquid test tube cultures.

Strain	Additional Plasmid	Gene Overexpressed	[Pantothe mg/L	nate]
PA221	pOTP61	none		1,000
				940
PA236	pAN236	panE1		2,030
	-	•		2,050
PA238	pAN238	panE2		530
,	•	-		680

Overexpression of panE1 caused a two-fold increase in pantothenate production when compared to the parent strain (e.g., to slightly over 2 g/L) whereas overexpression of panE2 resulted in a strain that produced about 35% less pantothenate than the parent strain. The panE1 nucleotide sequence and predicted amino acid sequence are set forth as SEQ ID NO:29 and SEQ ID NO:30.

EXAMPLE III: Enhanced Production of a Panto-Compound by Culturing Bacteria Overexpressing panE1 or panBCD in the Presence of Valine.

The ability of valine to function as a media supplement (e.g., as a substitute for α-KIV) in strains engineered to overexpress the panBCD operon and panE1 was evaluated. Valine is closely related to α-KIV by transamination, is less expensive than α-KIV, and is commercially available in kilogram quantities. Valine was substituted for α-KIV in the standard liquid test tube cultures in SVY medium. The concentration of valine was varied from 5 to 50 g/L. Although valine at 5 g/L was slightly less effective

than α -KIV in promoting pantothenate production, valine at 10 or 20 g/L equaled or surpassed 5 g/L α -KIV in promoting pantothenate production.

EXAMPLES IV-X Generation of Microorganisms Capable of Producing Pantothenate in a Precursor-Independent Manner

B. subtilis strains such as PA221 and PA235 (engineered to overexpress panBCD) and PA236 (engineered to overexpress panBCD and panE1) need to be fed α-ketoisovalerate (α-KIV) (or valine) and aspartate (or β-alanine) to achieve maximal pantothenate production, as both these precursors are limiting for pantothenate
synthesis. Accordingly, manipulated microorganisms were designed to eliminate the need to feed limiting precursors of pantothenate biosynthesis in the production of pantothenate. These strains are also useful in the production of various pantothenate biosynthetic pathway intermediates.

15 EXAMPLE IV: Generation of Microorganisms Capable of Producing Pantothenate in an Aspartate- (or β-Alanine) Independent Manner

The panD gene was cloned into B. subtilis expression vector pOTP61 to construct pAN423 (Figure 5). The nucleotide sequence of pAN423 is set forth as SEQ ID NO:78. The NotI restriction fragment containing panD was isolated from pAN423, self ligated and used to transform PA221. Transformants resistant to Tet¹⁵, Tet³⁰, and Tet⁶⁰ were isolated and saved for further analysis.

Six of the pAN423 transformants plus two control transformants were grown in SVY containing 5 g/l α -KIV with and without 10 g/l aspartate and then assayed for pantothenate production (Table 6).

Table 6. Effect of overproducing PanD on pantothenate production with and without added aspartate.

Culture* (PA221 transformants)	Asp (10 g/L)	TetR** (μg/ml)	OD550	[pan] (mg/L)
pOTP61-1	-	60	8.0	76
pOTP61-2	-	60	7.7	91
423#1-1	-	15	8.5	180
423#1-2	-	15	8.0	150
423#1-3	-	30	8.3	220
423#1-4	-	30	8.5	280
423#1-5	-	60	8.9	580
423#1-6	-	60	8.8	280

pOTP61-1	+	60	7.5	380
pOTP61-2	+ 1	60	6.9	560
423#1-1	+	15	8.5	1200
423#1-2	+ 1	15]	8.6	1000
423#1-3	+	30	8.8	1200
423#1-4	+	30	9.0	, 1200
423#1-5	+	60]	9.0	1200
423#1-6	+	60	9.0	1200

^{*}Test tubes cultures were grown in SVY + α -KIV (5 g/L) with Asp (10 g/L) where indicated.

The pAN423 transformants produced at least twice the amount of pantothenate as the controls (i.e., to a level at or near that which was obtained in earlier experiments by the addition of β -alanine to the culture medium). The data also show that in the absence of added aspartate, transformants containing additional copies of the panD gene expression cassette produce more pantothenate than the control transformants. One of the transformants, 423#1-5, produced about five times as much pantothenate as the controls. These results indicated that increased levels of PanD protein "pull" the conversion of available aspartate towards β -alanine, and that increasing panD gene expression can result in enhancement of pantothenate production both in the presence and absence of added aspartate.

Transformant 423#1-5 was re-named strain PA401 and studied further in shake flask fermentations. The shake flask medium was SVY with maltose instead of SVY with glucose. Results of shake flask experiments agreed well with test tube experiments during the first 24 hours. In shake flask experiments without the addition of β-alanine, PA401 produced approximately 1.5 g/l of pantothenate in 24 hours. Addition of β-alanine to the culture medium did not further improve pantothenate titers (Table 7), indicating that with this strain and these fermentation conditions, β-alanine is not limiting pantothenate production. In fact, when no β-alanine is fed, one can observe that PA401 is secreting β-alanine in significant amounts into the medium.

^{**}TetR = Approximate Tet-resistance of transformant

Table 7. Shake flask cultures with strain PA401 (panD) with and without \(\beta\)-alanine.

	Amino ao	cids (g/l)		24 hours			
Initial β-ala Added	β-ala	Val	рН	OD ₆₀₀	Pantothenate (g/l)		
0	0.7	1.5	7.5	13.7	1.5		
5 g/l	7.1	1.4	7.6	12.4	1.5		

Each value represents the average of duplicate 250 ml baffled flasks containing 50 ml of medium, incubated at 37°C with shaking (200 rpm).

Base Medium: SVY with 10 g/l α -KIV, 30 g/l maltose

2% Inoculum: SVY with Tet 15 grown 24 hours.

EXAMPLE V: Engineering the *panD* gene for Further Increased Synthesis of Aspartate Decarboxylase and Enhanced Production of Pantothenate

This Example describes the generation of improved ribosome binding sites (RBSs) in the *panD* gene to increase the translation of *panD* mRNA.

Increasing the translation of the panD gene mRNA by generation of synthetic panD RBSs

- The RBS (SEQ ID NO:88) used to express panD in pAN423 is a synthetic RBS and has been used to successfully produce other proteins in B. subtilis at a high level. However, it contains six mismatches when aligned to the "ideal" B. subtilis RBS (SEQ ID NO:45) (e.g., an RBS having a sequence which is complementary to the 16S RNA sequence within the B.subtilis ribosome). (See e.g., Table 1B, mismatches in bold).
- Two new RBSs were designed to more closely mimic the ideal RBS. These synthetic RBSs, named new design A (NDA) and new design B (NDB) (also referred to herein as RBS3 and RBS4), are set forth as SEQ ID NO:51 and SEQ ID NO:52 and are aligned with the ideal RBS in Table 1B.

Oligonucleotides corresponding to the top and bottom strands of each new RBS were synthesized, annealed, then used to replace the RBS in pAN420, generating plasmids pAN426 and pAN427. These constructions are illustrated in Figure 6. The presence of the NDA and NDB RBS in pAN426 and pAN427 was confirmed by DNA sequence analysis. Next, the *panD* genes from pAN426 and pAN427 were transferred to *B. subtilis* expression vector pOTP61 as shown in Figure 7, creating pAN428 and pAN429. The nucleotide sequence of pAN429 is set forth as SEQ ID NO:79.

Not restriction fragments lacking the E. coli vector sequences were isolated from pAN428 and pAN429, self-ligated, and used to transform strain PA221 to resistance to Tet¹⁵. Four isolates resistant to Tet⁶⁰ were picked from each transformation and assayed for pantothenate and β-alanine production along with PA221 transformed with the empty vector (pOTP61) and PA221 transformed with pAN423 (strain PA401) (see Table 8).

Table 8. Panthothenate production by test tube cultures of PA221 transformed with pAN428 and pAN429

•	^
1	()

Plasmid		Medium Supplements	OD550	Pan g/l	β-Ala g/l
рОТР61		α-KIV ⁵	10	UND	0.04
pAN423		α-KIV ⁵	10	0.4	0.04
pAN428-1	*	α-KIV ⁵	12	0.6	0.04
pAN428-2		α-KIV ^S	11]	0.5	0.03
pAN428-3		α -KIV ⁵	- 11	0.3	0:03
pAN428-4		α-KIV ⁵	10	0.1	UND
pAN429-1		α-KIV ^S	12	0.6	0.04
pAN429-2		α-KIV ⁵	11	0.5	0.04
pAN429-3		α-KIV ⁵	11	0.6	0.05
pAN429-4	#	α-KIV ⁵	12	0.8	0.10
pOTP61		α -KIV ₅ + Asp ₁₀	11	0.5	0.08
pAN423		α -KIV + Asp α -KIV ⁵ + Asp α -KIV ⁵ + Asp	12	0.9	1.32
•	*	\	12	0.8	1.97
pAN428-1	•	α -KIV ⁵ + Asp ¹⁰	12	0.8	1.51
pAN428-2		α -KIV + Asp 10 α -KIV + Asp 10	12	0.8	1.02
pAN428-3			11	0.8	0.30
pAN428-4		α -KIV ⁵ + Asp ¹⁰		. 0.8	0.50
pAN429-1		α -KIV ⁵ + Asp ¹⁰	· 12	0.8	1.78
pAN429-2		α-KIV + ASP	12	0.8	1.66
pAN429-3		α -KIV ³ + Asp ¹⁰	12	0.8	1.78
pAN429-4	#	α -KIV ⁵ + Asp ¹⁰	13	0.8	2.28

UND: Below the limits of detection. * Renamed PA402 # Renamed PA403

When grown in medium supplemented with α-KIV at 5 g/l (α-KIV⁵), the pAN428-1 transformant and all four of the pAN429 transformants produced more

pantothenate than did PA401, suggesting that these transformants contain higher levels of aspartate decarboxylase activity. When grown in medium supplemented with α-KIV⁵ and Asp¹⁰ none of the pAN428 or pAN429 transformants produced more pantothenate than PA401. However, the pAN428-1 transformant and all four of the pAN429 transformants produced significantly more β-alanine than did PA401. It is possible that the excess β-alanine produced from added aspartate causes inhibition of pantothenate production. Alternatively, β-alanine may accumulate because pantoate is limiting in these strains.

The strains that produced the highest level of β-alanine, the pAN428-1 and pAN429-4 transformants, were renamed PA402 and PA403, respectively. These two strains were grown in SVY medium supplemented with various intermediates and reassayed for pantothenate and β-alanine production. PA221 and PA401 were included as controls. The results of the assays are presented in Table 9.

15 Table 9. Pantothenate production of PA402 and PA403 in test tube cultures.

Strain	Medium Supplements	OD ₅₅₀	Pan g/l	β-Ala g/l	Val g/l
PA221	α-ΚΙV ⁵	7.9	UND	UND	0.9
PA401	α-ΚΙV ⁵	8.7	0.3	0.04	0.9
PA402	α-ΚΙV ⁵	8.5	0.5	0.04	0.9
PA403	α-ΚΙV ⁵	9.4	0.7	0.07	0.9
PA221 PA401 PA402 PA403	$\alpha\text{-KIV}^{5} + \text{Asp}^{10}$ $\alpha\text{-KIV}^{5} + \text{Asp}^{10}$ $\alpha\text{-KIV}^{5} + \text{Asp}^{10}$ $\alpha\text{-KIV}^{5} + \text{Asp}^{10}$	9.8 9.1 9.4 9.7	0.4 0.8 0.8 0.7	0.11 1.15 2.02 2.40	0.8 0.8 0.8
PA221	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	8.9	UND	UND	0.2
PA401		8.7	0.3	0.02	0.2
PA402		10.6	0.5	0.02	0.2
PA403		10.5	0.7	0.02	0.2
PA221	Pantoate ⁵ + Asp ¹⁰	9.5	0.4	0.06	0.2
PA401		9.2	2.2	0.62	0.2
PA402		9.1	2.8	1.17	0.2
PA403		10.2	2.9	1.58	0.2

UND: Below the limits of detection.

When grown in medium supplemented with either α-KIV⁵ or Pantoate⁵, PA402 and PA403 produced significantly more pantothenate than did PA401. As before, even though PA402 and PA403 produced significantly more β-alanine than PA401 when grown in medium supplemented with α-KIV⁵ and Asp¹⁰, they did not produce a proportional increase in pantothenate. However, when grown in medium supplemented with Pantoate⁵ plus Asp¹⁰, both PA402 and PA403 produced significantly more pantothenate than PA401, about a 30% increase.

It can be concluded from these experiments that the improved NDA and NDB panD ribosome binding sites, engineered into pAN428 and pAN429, respectively, lead to increased levels of aspartate decarboxylase activity.

Increasing the translation of the panD gene mRNA by generation of synthetic panD RBSs within the panBCD operon

15

20

25

The native B. subtilis panD gene ribosome binding site (RBS) (SEQ ID NO:43), which is found in the $P_{26}panBCD$ operon cassette present in PA221 (and in other engineered pantothenate production strains described herein), is shown in Table 1C aligned with the ideal ribosome binding site (SEQ ID NO:47). The alignment shows mismatches between the native B. subtilis panD gene RBS, which is located within the coding sequence for PanC, and the the ideal RBS. Three new RBSs (within the P26 panBCD operon cassette) were generated to increase translation of the panD gene mRNA and to yield increased synthesis of aspartate decarboxylase. These synthetic RBSs (termed NDI, NDII, and NDIII, also referred to herein as RBS5, RBS6 and RBS7, respectively) are set forth as SEQ ID NO:55, SEQ ID NO:56 and SEQ ID NO:57, respectively) and are included in Table 1C. It should be noted that although changes in the panD RBS within the panBCD operon also changes the C-terminal amino acid sequence of the PanC protein encoded by that operon, an alignment of known and suspected PanC protein amino acid sequences showed that the sequence of the last nine amino acids of the B. subtilis PanC protein could be altered without affecting any conserved amino acid residues indicating that such changes should not reduce pantothenate synthetase activity or expression. The new RBSs were synthesized and incorporated into the P_{26} panBCD operon expression cassette as follows.

First, PCR primers were designed to contain the following elements: (1) a nucleic acid sequence encoding the first five amino acids of PanD up to and including a unique *BsiWI* restriction site that had been previously introduced into *panD* by PCR; (2)

a stop codon for panC, (3) at least one synthetic RBS; and (4) 30-39 bp of nucleic acid sequence having 100% identity with panC upstream of the panD RBS. The primers were named TP102, TP103, and TP104 and contain the NDI, NDII, and NDIII ribosome binding sites, respectively. These three primers were used in conjunction with the 5' primer TP101, which hybridizes near the start codon of panC, in three independent PCR reactions to generate the NDI, NDII, and NDIII PCR products. The PCR products were purified, digested with XbaI, then cloned into plasmid vector pASK-1BA3 which had been digested with XbaI and SmaI. The resulting plasmids were named pAN431, pAN432, and pAN433. The construction of pAN431 is illustrated in Figure 8 and is representative of all three plasmid constructions. The presence of the desired synthetic panD gene RBS in each new plasmid was confirmed by DNA sequencing.

Next, the modified panC genes containing the new panD RBSs were joined with the panD gene utilizing the unique BsiWI restriction site. This was accomplished by isolating the appropriate NsiI-BsiWI restriction fragments from pAN431, pAN432, and pAN433 and ligating them with a 2395 bp NsiI-BsiWI restriction fragment from pAN420, which supplied the BsiWI-modified panD gene. These constructions resulted in plasmids pAN441, pAN442, and pAN443, respectively. A representative construction (pAN441) is illustrated in Figure 9. The nucleotide sequence of pAN443 is set forth as SEQ ID NO:80.

The new panD gene RBSs were then substituted into the P₂₆panBCD operon expression cassette as follows. First, a deletion-insertion mutation which removes the region of panC containing the panD RBS was created. This was constructed by digesting pAN430 with a mixture of BspE1 and BgIII and recovering the 4235 bp fragment which is now missing the 3' end of panC and the 5' end of panD. This fragment was ligated with an AvaI-BamHI restriction fragment from plasmid pECC4, which contains the chloramphenicol acetyl transferase (cat) gene. The 5' extension produced by AvaI digestion is compatible with that produced by BspEI while the BgIII and BamHI extensions are also compatible. The resulting plasmid was named pAN440, and its construction is illustrated in Figure 10.

The resulting deletion-insertion mutation was crossed into the P_{26} panBCD operon via homologous recombination by transforming PA221 with linearized pAN440 and selecting for resistance to chloramphenicol on Cam⁵ plates containing 1 mM pantothenate. Several transformants were tested, and were all found to require 1 mM pantothenate for growth, as expected. Two of these transformants were remaned PA408A and PA408B and were assayed for pantothenate production. Neither strain synthesized measurable quantities of pantothenate, even when grown in medium

20

30

containing pantoate and β -alanine at 5 g/l, indicating that the strains are deficient in pantothenate synthetase activity. Next, the new panD RBSs were crossed into the P_{26} panBCD operon by transforming PA408 with linearized pAN441, pAN442, and pAN443 plasmid DNA and selecting for growth on TBAB plates without pantothenate supplementation. A transformation with linearized pAN430 (including the native panD RBS) was included as a control and was expected to give rise to transformants identical to PA221 described herein. Four isolates from each transformation were assayed for pantothenate and β -alanine production in SVY medium supplemented with various intermediates (Tables 10 and 11).

10

Table 10. Pantothenate production of PA410 - PA413 in test tube cultures.

Strain	RBS	Medium Supplements	OD ₅₅₀	Pan g/l	β-Ala g/l
PA221	native	Pantoate ⁵	11	UND	UND
PA410-1 PA410-2 PA410-3 PA410-4	native	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 12 12 12	UND UND UND UND	UND UND UND UND
PA411-1 PA411-2 PA411-3 PA411-4	NDI	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 12 12 12	0.23 0.20 0.19 UND	UND UND UND UND
PA412-1 PA412-2 PA412-3 PA412-4	NDII	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 11 13 12	UND UND 0.18 0.18	UND UND UND UND
PA413-1 PA413-2 PA413-3 PA413-4	NDIII	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 12 12 12	0.18 0.17 0.16 0.17	UND UND UND UND

UND: Below the limits of detection.

Table 11. Pantothenate production of PA410 - PA413 in test tube cultures.

Strain	RBS	Medium Supplements	OD ₅₅₀	Pan g/i	β-Ala g/l
PA221	native	Pantoate ⁵ + Asp ¹⁰	11	0.3	UND
PA410-1 PA410-2 PA410-3 PA410-4	native	Pantoate 5 + Asp 10	12 12 12 12	0.4 0.4 0.4 0.4	UND UND UND UND
PA411-1 PA411-2 PA411-3 PA411-4	NDI	Pantoate 5 + Asp 10	13 13 13 13	1.7 1.7 1.8 0.4	0.4 0.4 0.3 UND
PA412-1 PA412-2 PA412-3 PA412-4	NDII	Pantoate 5 + Asp 10	13 12 12 12	0.4 0.4 1.6 1.5	UND UND 0.3 0.2
PA413-1 PA413-2 PA413-3 PA413-4	NDIII	Pantoate ⁵ + Asp ¹⁰	13 13 13 13	1.6 1.6 1.7 1.7	0.3 0.4 0.4 0.4

UND: Below the limits of detection.

As expected from previous experiments using PA221, none of the transformants that contained the native panD RBS produced measurable quantities of pantothenate when grown in medium supplemented with pantoate. However, nine of the twelve transformants expected to contain modified panD RBSs produced significant quantities of pantothenate (160-230 mg/l) under these conditions, indicating that they possess elevated levels of aspartate decarboxylase activity. When grown in medium supplemented with both pantoate and aspartate, these same nine transformants produced approximately four times more pantothenate than those with the native panD RBS. In addition, these nine transformants accumulated measurable quantities of β-alanine (230-410 mg/l). All transformants produced roughly equivalent quantities of pantothenate when grown in medium containing pantoate and \beta-alanine, demonstrating that each 15 contains a functional pantothenate synthetase.

5

These data demonstrate that the synthetic panD RBSs are about four times more effective than the native panD RBS in directing translation of the panD gene mRNA and evidence the utility of such synthetic RBSs in enhancing pantothenate production.

Additional approaches to increasing pantothenate production can include, for example, increasing the half-life of the panD gene mRNA, increasing the strength of the promoter for panD transcription and/or increasing the stability of the PanD protein.

EXAMPLE VI: Construction of Strains Containing an Integrated P_{26} panE1 Cassette without an Antibiotic Resistance Gene.

10

15

20

Example II describes the identification of the B. subtilis panE1 gene that encodes the enzyme responsible for the majority of the ketopantoate reductase activity in B. subtilis. PA236 (containing the pAN236 plasmid) produced about twice as much pantothenate (2 g/l) as its parent strain, PA221 (1 g/l) in 24 hour SVY test tube cultures. PA236 was presumed to contain an amplified (~3 copies) integrated pAN236 plasmid based on selection for tetracycline resistance (the tetR gene product being encoded on the pAN236 plasmid in addition to the P_{26} panE1 cassette). Also useful in the methodologies of the present invention are strains that contain a single integrated unamplifiable copy of P_{26} panE1 at the panE1 locus, for example, without an antibiotic resistance gene in the strain. Such a strain was generated as follows.

A plasmid named pAN251 was derived from pAN236 by inserting additional chromosomal sequences just upstream and just downstream from the P_{26} panE1 cassette. These additional sequences, which provide homology to allow integration of the P_{26} panE1 cassette at panE1 by double crossover, were obtained by PCR from chromosomal DNA as a template. pAN251 is shown in Figure 11. The nucleotide sequence of pAN251 is set forth as SEQ ID NO:81.

Next, a strain was constructed which allowed selection for the incoming P_{26} panE1 cassette. The strain included the following three components: (1) P_{26} panBCD; (2) $\Delta panE1$; and (3) ilvC, since both panE1 and ilvC must be mutated to have a Pan phenotype. The starting strain was CU550 (trpC2, ilvC4, leuC124). The P_{26} panBCD cassette from PA221 chromosomal DNA was introduced in two steps to create strain PA290. Next, $\Delta panE1::spec$ was transformed into PA290, using chromosomal DNA from strain PA240, to give strain PA294 (trpC2, ilvC4, leuC124, P_{26} panBCD, $\Delta panE1::spec$), which is a strict pantothenate auxotroph. Finally, PA294 was transformed with plasmid pAN251, selecting for pantothenate prototrophy, to give strain PA303. This strain was expected to have the genotype trpC2, ilvC4, leuC124, P_{26}

20

panBCD, P_{26} panE1. PA303 was checked for the correct chromosomal structure at the panE1 locus by PCR using primers that flank the P_{26} insertion just upstream of panE1. The PCR product from PA303 was of the expected size, with a concomitant loss of the PCR product from the wild type panE1 gene, consistent with having obtained the desired double crossover event. Furthermore, PA303 was tetracycline sensitive, which is also consistent with the desired double crossover event, as opposed to a Campbell-type single crossover of the plasmids into the chromosome. The trp, ilv, and leu auxotrophies from the parent strain were all maintained in PA303.

In 24 hour liquid SVY test tube cultures, PA303 produced almost the same level of pantothenate as positive control PA236, and about twice as much as PA221, which does not contain engineered *panE1* as indicated in Table 12.

Table 12. Pantothenate production by 24 hr. test tube cultures of PA303 and controls grown in SVY plus 5 g/l α -KIV and 5 g/l β -alanine.

Strain	OD600	[pan] g/l
D		
PA221-1	10.9	0.85
PA221-2	10.5	0.85
PA236-1	9.5	1.74
PA236-2	9.3	1.70
PA303-1	10.8	1.66
PA303-2	10.7	1.61

EXAMPLE VII: Generation of Microorganisms Capable of Producing Pantothenate in an α -KIV (or Valine) Independent Manner

 α -ketoisovalerate (α -KIV) is a rate limiting intermediate for pantothenate production in certain strains deregulated for pantothenate synthesis. Addition of either α -KIV or valine at 5 g/l increases pantothenate production about 5-fold in test tube cultures with strains such as PA221. In order to alleviate the need to feed either α -KIV or valine, strains were engineered that have an increased capacity to synthesize α -KIV.

25 α-KIV is produced in *B. subtilis* from pyruvate by the sequential action of three enzymes encoded by four genes, *ilvB* and *ilvN*, *ilvC*, and *ilvD*. In a wild type *B. subtilis*, three of the genes (*ilvB*, *ilvN*, and *ilvC*) are the first three genes of the large *ilv-leu* operon. The fourth gene necessary for α-KIV synthesis, *ilvD*, is located by itself elsewhere on the chromosome. The *B. subtilis ilv-leu* operon is thought to be regulated

15

25

30

35

only by leucine levels. Feeding of exogenous leucine reduces transcription of the ilv-leu operon by about 13-fold, probably by an attenuation mechanism (Grandoni et al. (1992) J. Bacteriol. 174: 3212-3219). The only known feedback regulation in the ilv-leu pathway is the inhibition of the leuA gene product by leucine.

As a first step to deregulate the synthesis of α-KIV, a copy of the ilvBNC region from the wild type B. subtilis ilv-leu operon was isolated by PCR, and installed adjacent to the P_{26} promoter and RBS2 on a vector, pOLL8, that was designed to integrate a single P_{26} expression cassette by double recombination at the amy E locus. The amy E gene encodes a nonessential a-amylase, and is a useful locus for installing expression 10 cassettes. The resulting plasmid, pAN267, is illustrated in Figure 12. The nucleotide sequence of pAN267 is set forth as SEQ ID NO:82. pAN267 readily gave stable transformants by double crossover at the amyE locus of B. subtilis strains, as described in detail below.

Construction of pantothenate overproducing strains that are leucine prototrophs

Initially, a B. subtilis strain containing ilvC4 and ApanE1 was used to introduce a single copy of P_{26} panE1 into the chromosome without using an antibiotic resistance gene. The double mutant was required to select for the incoming P_{26} panE1 cassette because a ApanE1 mutation alone does not result in pantothenate auxotrophy. A strain named CU550 was obtained containing ilvC4 to be used as a basis for this type of strain construction. However, CU550 also contains a closely linked leuC124 mutation, so all strains derived from CU550 required leucine. Having shown that the combination of P_{26} panBCD and P_{26} panE1 was favorable for pantothenate production, the next step was to reassemble this combination of two cassettes in a leucine prototroph.

Accordingly, the two cassettes were combined in two different strain backgrounds, RL-1 and PY79. To introduce chromosomal P26 panE1 into the PY79 and RL-1 strain backgrounds without using an antibiotic resistance gene, a strategy was used that did not rely on ilvC4. (The strategy took advantage of the observation that the ApanEl mutation causes a pantothenate bradytrophy, manifested by relatively small colonies on TBAB (rich) plates). First, ΔpanB::cat and ΔpanE::spec were introduced into both strain backgrounds. Next, the resulting strains were transformed simultaneously with DNA from two strains, PA221 (P26 panBCD) and PA303 (P26 panEl), selecting for Pan+ on TBAB plates. Colonies of two distinct sizes grew on the selective plates, with the larger size comprising about 2% of the colonies. The larger colonies were presumed to represented co-transformants that received both P_{26} panBCD and P_{26} panE1, and that the smaller colonies had received only P_{26} panBCD. Consistent with this prediction, the larger colonies had lost both Cam^r and Spec^r, while the smaller colonies had lost only the *cat* gene, and retained the *spec* gene. Furthermore, a representative derivative of PY79 named PA327, and a representative derivative of RL-1, named PA328, both produced the elevated levels of pantothenate in test tube cultures which was about 1.6 to 1.7 g/l (Table 13).

Table 13. Pantothenate production of PA327, PA328, and controls from 24 hr test tube cultures grown in SVY plus 5 g/l α-KIV and β-alanine.

Strain	Background	P26 panE1 copy number	[pan] g/l	
DA221 1	DI 1	•		_
PA221-1	RL-1	0	0.92	
PA221-2	RL-1	0	0.95	
PA236-1	RL-1	amplified (~3)	1.60	
PA236-2	RL-1	amplified (~3)	1.73	
PA327-1	PY79	1	1.66	
PA327-2	PY79	1	1.65	
PA328-1	RL-1	1	1.61	
PA328-2	RL-I	1	1.91	

10

Thus, PA327 and PA328 were concluded to contain both P_{26} panBCD and P_{26} panE1, and were used for further constructions as described below. PCR analysis confirmed the presence of the two cassettes.

15 Installation of a stable P26 ilvBNC cassette into two lineages of pantothenate overproducing strains

Having constructed PA327 and PA328, derivatives of PY79 and RL-1 that contain P26 panBCD and P26 panE1, and that are Leu⁺, the next step was to introduce stable copies of P26 ilvBNC. This was accomplished by transforming PA327 and PA328 with plasmid pAN267, selecting for Spec^r. Screening by PCR showed that about 85% of the obtained transformants contain P26 ilvBNC integrated at amyE by double crossover. One transformant of PA327, named PA340, and one transformant of PA328, named PA342, were chosen for further study.

In test tube cultures grown in SVY medium plus 5 g/l β-alanine but without added α-KIV, both PA340 and PA342 gave the expected increase in pantothenate production over that of PA327 and PA328, to about 1.3 to 2 g/l (Table 14).

15

Table 14. Pantothenate and valine production by PA340 and PA342, both containing P₂₆ ilvBNC in 24 hr test tube cultures-grown in SVY with 5 g/l β-alanine and with or without 5 g/l α-KIV

Strain	Back- ground	OD ₆₀₀ - α-KIV	+α-KIV	[pan] g/l - α-KIV	+α-KIV	{val] g/l - α-KIV	+ α-KIV
PA340-1 PA340-2	PY79 PY79	11.8	7.1 7.5	2.02	2.10 2.03	0.38 0.40	0.90 0.91
PA342-1 PA342-2	RL-I RL-I	10.2 9.6	8.0 9.2	1.29	1.89 2.04	0.27 0.21	0.78 0.79

The two new strains also gave a slight increase in value secretion, indicating that the ilvBNC genes had been deregulated. However, when the same strains were grown with 5 g/l α -KIV added, a further increase in pantothenate production occurred from PA342, suggesting that α -KIV was still rate limiting in this strain background. Similar results, only with more growth and hence higher pantothenate levels, were seen in shake flask cultures (Table 15).

Table 15. Pantothenate and valine production by PA340 and PA342, both containing P_{26} ilvBNC in 24 hour shake flask cultures grown in SVY with 5 g/l β -alanine and with or without 5 g/l α -KIV.

Strain	Back- ground	OD600 - α-KIV	+ α-KIV	[pan] g/l - α-KIV	+ α-KIV	[val] g/l - a-KIV	+ α-KIV
PA327	PY79	21	22	0.6	3.0	0.5	1.3
PA340-1	PY79	20	20	3.5	4.1	1.0	1.9
PA340-2	PY79	22	19	3.0	2.1	0.8	1.4
PA328	RL-1	20	16	1.4	2.7	0.6	1.3
PA342-1	RL-1	17	16	3.3	3.6	0.9	1.6
PA342-2	RL-1	18	18	3.1	4.2	0.8	1.4

EXAMPLE VIII: Increasing panD Copy Number in Strains Engineered to Overproduce panE1 and the ilvBNC Gene Products Enhances Pantothenate Production

Experiments where β-alanine was fed to cultures of engineered B. subtilis strains consistently showed that β-alanine was a rate limiting intermediate in pantothenate synthesis. The effect of adding additional copies of panD on pantothenate production in PA340 and PA342 was examined. Strains PA340 and PA342 were transformed with chromosomal DNA isolated from PA401 with selection on plates containing 15 µg/ml of tetracycline (Tet¹⁵ plates). Transformants derived from each parent were patched onto Tet⁶⁰ plates to identify those which were likely to contain multiple copies of the 10 expression cassette. Twelve transformants from each transformation which grew on Tet⁶⁰ were streaked for single colonies on this medium and then assayed in SVY medium test tube cultures for pantothenate production. One transformant from each group was found to produce greater than 300 mg/l pantothenate in 24 hours. These two transformants were saved and named PA404 (PA340 strain background) and PA405 (PA342 strain background). Both strains were resistant to spectinomycin, indicating that the P_{26} ilvBNC expression cassette was still present at amy E. PCR analysis of chromosomal DNA isolated from each strain confirmed that the deregulated panE1 gene had also been retained.

Next, PA404 and PA405 were evaluated in shake flask cultures which were grown in SVY medium containing maltose as the carbon source and supplemented with various intermediates. The cultures were grown for 24 and 48 hours and then assayed for pantothenate, β-alanine, and valine production. The results of this experiment are presented in Table 16. Analogous shake flask culture data for the parent strains (PA340 and PA342) are included in the tables for comparison.

Table 16. Pantothenate production by PA404 and PA405 in shake flask cultures after 24 hours

Strain	Medium Supplements	OD600	Pan g/l	β-Ala g/l	Val g/l
PA340	none	20	0.4	<0.1	1.0
PA404	none	22	1.8	<0.1	0.7
PA342	none	19	0.3	0.2	0.7
PA405	none	19	1.4	0.4	0.5

20

1 P	PA340	β-alanine ⁵	18	3.6	3.2	0.6
	PA404	β-alanine ⁵	18	2.8	5.1	0.7
	PA342*	β-alanine ⁵	17	3.3	3.3	0.5
	PA405*	β-alanine ⁵	19	1.3	6.5	0.6

Values are the average of duplicate flasks except where indicated by *.

In the absence of any medium supplementation, PA404 and PA405 made four to five times more pantothenate in 24 hours compared to their isogenic parent strains (Table 16). The supply of β-alanine was clearly limiting in the parent strains PA340 and PA342. Addition of amplified P26 panD greatly increased the supply of β-alanine.

EXAMPLE IX: Deregulation of the *B. subtitis ilvD* Gene Enhances Pantothenate Production

To deregulate expression of the *ilvD gene*, standard procedures (described above) were used to integrate the constitutive P_{26} promoter and an artificial ribosome binding site, RBS2, just upstream of the *ilvD* coding region. The *ilvD* gene maps by itself, unlinked to the *ilvBNC* operon. First, a 2.4 kb region of the RL-1 chromosome that contains the *ilvD* coding region and 730 bp of upstream sequence was cloned by PCR into a low copy (about 15 per *E. coli* cell) vector called pOK12, to give plasmid pAN257, shown in Figure 13.

Taking advantage of a natural EcoRI site just upstream of the native ilvD gene promoter, and a natural NcoI site at the ilvD start codon, an artificial sequence containing P_{26} and RBS2 was inserted into pAN257 to give pAN263 (Figure 14). The nucleotide sequence of pAN263 is set forth as SEQ ID NO:83. In parallel with this construction, the cat gene was also inserted into pAN257, between the same upstream EcoRI site and a BgIII site in the middle of the ilvD coding region, to give pAN261, which is deleted for a large portion of the ilvD gene (Figure 15). Using pAN261 and pAN263, the P_{26} ilvD cassette could then be installed in the B subtilis chromosome in two steps. In the first step, pAN261 is introduced by transformation, selecting for chloramphenicol resistance, and then confirming an IIv phenotype. In the second step, pAN263 is introduced, selecting for IIv $^+$, checking for chloramphenicol sensitivity, and confirming correct local structure by PCR.

pAN261 was first transformed into strain RL-1 (highly competent) to give strain PA343 (*AilvD::cat*), and then chromosomal DNA from PA343 was used to transform PA340 and PA342 to IIv auxotrophy, yielding strains named PA348 and PA349, respectively. Chromosomal DNA is inherently more efficient than monomeric plasmid

in transforming *B. subtilis*. Similarly, pAN263 DNA was transformed into PA343 (moderately competent) to give strain PA345 (P_{26} ilvD), and then PA345 chromosomal DNA was used to transform PA348 and PA349 to Ilv^+ prototrophy, yielding strains PA374 and PA354, respectively.

As predicted, PA374 and PA354 gave further increases in pantothenate production, to about 2.5 to 2.9 g/l, in test tube cultures grown in SVY plus 5 g/l β-alanine (Table 17).

Table 17. Pantothenate and valine production by PA374 and PA354, containing P_{26} ilvD, and controls, in 24 hr test tube cultures grown in SVY with 5 g/l β -alanine and with or without 5 g/l α -KIV.

		,	OD ₆₀₀		[pan] g/l	•	[val] g/l	
Strain	Back- ground	ilvD status	α-KIV -	+	α-KIV -	+	α-KIV -	+
PA340	PY79	w.t.	9.2	9.0	2.14	2.23	0.38	0.90
PA348	PY79	ilvD::cat	11.7	10.0	0.19	2.23	0.19	0.91
PA374-1 PA374-2	PY79 PY79	P26 ilvD P26 ilvD	9.1 8.2	7.3 7.7	2.93 2.99	2.40 2.36	0.58 0.60	0.87 0.95
PA342	RL-1	w.t.	10.2	8.0	1.29	1.89	0.27	0.78
PA349	RL-1	ilvD::cat	8.1	7.7	0.17	1.87	0.22	0.88
PA354-1 PA354-2	RL-1 RL-1	P ₂₆ ilvD P ₂₆ ilvD	9.6 7.5	9.6 8.2	2.57 2.48	2.03 2.24	0.65 0.64	1.23 0.97

In the absence of added β -alanine, strains PA374 and PA354 produced only about 0.2 g/l pantothenate in test tube cultures, indicating that PanD activity is significantly rate limiting.

To alleviate this limitation, the amplifiable P_{26} panD cassette from strain PA401 was installed. PA401 chromosomal DNA was transformed into PA374 and PA354, selecting for Tet^r at 15 mg/l, to yield strains PA377 and PA365, respectively. After transformants were obtained, the strains were streaked on plates containing 30 and 60 mg/l tetracycline to reamplify the copy number of the P_{26} panD cassette integrated at the bpr locus. In test tube cultures grown in SVY without α -KIV or β -alanine, a substantial improvement in pantothenate titers over those of PA374 and PA354 was obtained (Tables 18 and 19).

WO 01/21772

5

10

Table 18. Pantothenate production by PA365, containing amplified P_{26} panD, and controls, in 24 and 36 hr test tube cultures grown in SVY-glucose without β -alanine or α -KIV.

	OD ₆₀₀		[pan] g/l	[pan] g/l	
Relevant genotype	24 hrs.	36 hrs	24 hrs.	36 hrs.	
w.t. <i>ilvD</i>	11.7	8.8	b.d.	0.27	
w.t. ilvD	12.8	8.8	b.d.	0.26	
P ₂₆ ilvD	n.d.	11.0	n.d.	0.19	
P ₂₆ ilvD	n.d.	8.4	n.d.	0.20	
P26 ilvD, P26 panD	9.8	10.0	1.01	2.07	
P ₂₆ ilvD, P ₂₆ panD	9.9	10.4	0.96	2.09	
	w.t. ilvD w.t. ilvD P26 ilvD P26 ilvD P26 ilvD, P26 panD	W.t. ilvD 11.7 w.t. ilvD 12.8 P26 ilvD n.d. P26 ilvD, P26 panD 9.8	Relevant genotype 24 hrs. 36 hrs w.t. ilvD 11.7 8.8 w.t. ilvD 12.8 8.8 P26 ilvD n.d. 11.0 P26 ilvD n.d. 8.4 P26 ilvD, P26 panD 9.8 10.0	Relevant genotype 24 hrs. 36 hrs 24 hrs. w.t. ilvD 11.7 8.8 b.d. w.t. ilvD 12.8 8.8 b.d. P26 ilvD n.d. 11.0 n.d. P26 ilvD n.d. 8.4 n.d. P26 ilvD, P26 panD 9.8 10.0 1.01	

'n.d. = not determined; b.d. = below detection

Table 19. Pantothenate production by PA377, containing amplified P_{26} panD, and controls, in 27 hr test tube cultures grown in SVY-glucose or SVY-maltose, without α -KIV, and with or without β -alanine.

		OD ₆₀₀			
•	Relevant genotype	- β-ala	+ β-ala	- β-ala	+β-ala
Strain		Glucose	Glucose	Maltose	Maltose
PA374-1	P ₂₆ ilvD	9.4	9.8	7.0	6.4
PA374-2	P ₂₆ ilvD	9.2	9.6	6.6	6.3
PA377-1	P26 ilvD, P26 panD	10.0	7.6	7.2	6.1
PA377-2	P ₂₆ ilvD, P ₂₆ panD	10.5	7.8	9.4	5.4
PA3/1-2	F 26 HVD, F 26 panD	10.5	7.0	9.4	3.4
		1.5 2.0			
		[pan].g/l			
	Relevant genotype	j - β-ala	+ β-ala	- β-ala	+ β-ala
Strain		Glucose	Glucose	Maltose	Maltose
		1		<u>-</u>	
PA374-1	P ₂₆ ilvD	0.04	2.76	0.14	1.31
PA374-2	P ₂₆ ilvD	0.10	2.65	0.15	1.33
PA377-1	P ₂₆ ilvD, P ₂₆ panD	1.25	2.76	1.26	1.10
PA377-2	P26 ilvD, P26 panD	1.25	2.35	1.31	1.26

In SVY with glucose, an increase in pantothenate production can still be
achieved by feeding 5 g/l β-alanine suggesting that increasing panD expression further
might increase pantothenate production. In SVY with maltose, no further increase in
pantothenate was obtained by feeding β-alanine suggesting that β-alanine and/or

aspartate synthesis is suppressed by glucose. Strains PA377 and PA365 have been evaluated in 10 liter fermentors, where they typically produce above 20 g/l pantothenate in 48 hours without supplemental β -alanine and α -KIV or valine, described in detail below.

5

EXAMPLE X: 10 liter Fermentations of Pantothenate-Producing Microbes

Engineering of the P_{26} ilvBNC and P_{26} ilvD cassettes to give strains PA342 and PA354 allowed the production of 22 and 26 g/l of pantothenate, respectively, without the addition of valine or α -KIV to the fermentation medium (Table 20). At 48 hours, both strains had secreted about 0.5 g/l of valine into the medium.

Table 20. 10-liter fermentations of five pantothenate overproducing strains.

Strain	Medium	Feed 40% Glucose plus	OD 600 48 hr	Valine 48 hours g/l	β-ala 48 hr g/l	Pa 36 hr	ntothena g/L 48 hr	te 72 hr
PA 236	SVYG	50 g/l β-ala 25 g/l α-KIV	108	added	. added	16	19	21
PA 342	SVYG	50 g/l β-ala	92	0.5	added	17	22	
PA 354	SVYG	50 g/l β-ala	90	0.5	added	19	26	
PA 365	svyg	25g/l YE	77	0.85	0.4	18	21	27
PA 377	svyg	25g/l YE	85	1.5	0.5	18	22	31
PA 377	PFMG	25g/l YE	96	0.8	0.4	19	25	29
PA377	PFMG	-	71	0.7	0.1	16	21	-

15 Pantothenate synthesis in fermentors

With the addition of the P_{26} panD cassette to strains PA354 and PA374 to create strains PA365 and PA377, neither β -alanine nor α -KIV needed to be added to the fermentors. Strain PA365 produced 21 g/l pantothenate in 48 hours and 27 g/l in 72 hours with no precursors added to the medium (Table 20). PA377 was somewhat better, producing 18 g/l of pantothenate in 36 hours, 22 g/l in 48 hours, and 31 g/l in 72 hours). Valine was measured at 0.85 and 1.5 g/l for strains PA365 and PA377, respectively, at

48 hours in SVYG medium. Strain PA377 maintained valine between 1-1.5 g/l throughout most of the fermentation and β-alanine between 0.2 and 0.5 g/l.

Strain PA377 was further evaluated in 10-liter fermentors in yeast extract based PFMG medium. Pantothenate yields in PFMG and SVYG medium were similar. In PFMG, PA377 produced 19 g/l of pantothenate in 36 hours, 25 g/l in 48 hours, and 29 g/l in 72 hours. In SVYG, PA377 produced 18 g/L pantothenate in 36 hours, 22 g/L in 48 hours and 31 g/L in 72 hours (Table 20).

EXAMPLE XI: Converting Strain PA377 to a Tryptophan Prototroph

10

PA377 (Trp) was transformed to Trp⁺ using chromosomal DNA from PY79 to give strain PA824. After re-amplification of the $P_{26}panD$ casette, PA824 was compared to PA377 for pantothenate production in test tube cultures grown in SVY glucose with or without 5 g/L β -alanine (Table 21).

15

Table 21: Trp⁺ derivatives of PA377: Pantothenate production in 48 hour test tube cultures grown in SVY glucose, ±β-alanine

•		OD ₆₀₀		[pan] g/L		
Strain	trpC donor	- β-alanine	+ β-alanine	- β-alanine	+ β-alanine	
PA377-1	RL-1	8	8	1.5	3.4	
PA377-2	RL-1	8	9	1.6	3.6	
PA824-1	PY79	12	10	0.7	3:7	
PA824-2	PY79	11	11	1.9	4.9	
		ļ				

The Trp+ strains grew to slightly higher densities than PA377. In the absence of exogenous β -alanine, all of the strains produced similar levels of pantothenate, while with the addition of β -alanine, the Trp+ derivatives produced somewhat more pantothenate.

25 Fermentor studies with PA824

PA824 was evaluated in CF3000 Chemap 14 liter vessels with 10 liter working volumes. Formulations for two of the media used in the fermentors are given in Tables 22 and 23.

Table 22: Formulation for PFMG-5 medium

BATCH

	BAICH	/ / / · · · · · · · · · · · · · · · · ·
	MATERIAL	g/L (final [])
1	Amberex 1003	10
2	Na Glutamate	5
3	(NH ₄) ₂ SO ₄	8
4	MAZU DF 37C	2.5
	Added After Sterilization and C	ool Down
1	KH ₂ PO ₄	10
2	K ₂ HPO ₄ ·3H ₂ O	20
1	Glucose	20
2	MgCl ₂ ·6H ₂ O	1
3	CaCl ₂ ·2H ₂ O	0.1
1	Sodium Citrate	1
2	FeSO ₄ ·7H ₂ O	0.01
3	SM-1000X	1.0 ml
	H ₂ O	qs to 6000 m

FEED

	MATERIAL	g/L
1	Glucose	600
2	CaCl ₂ ·2H ₂ O	0.6
	H ₂ O	qs to 3000 ml

Table 23: Formulation for SVY-4 medium

BATCH

	BAICH	
j	MATERIAL	g/L (final [])
1	Veal Infusion	25
2	Yeast Extract	5.
3	Na Glutamate	5
4	(NH ₄) ₂ SO ₄	4
5	MAZU DF 37C	2.5
	Added After Sterilization and Co	pol Down
1	KH₂PO₄	10
2	K₂HPO₄·3H₂O	20
1	Glucose	20
2	MgCl ₂ ·6H ₂ O	1
3	CaCl₂·2H₂O	0.1
1	Sodium Citrate	1
2	FeSO ₄ ·7H ₂ O	0.01
3	SM-1000X	1.0 ml
	Н,О	qs to 6000 m

FEED -

	MATERIAL	g/L
1	Glucose	. 600
2	CaCl ₂ ·2H ₂ O	0.6
		2000 1
	H ₂ O	qs to 3000 ml

All fermentations were glucose limited fed batch processes. Immediately after inoculation, agitation was set at 200 rpm. The initial batched 2% glucose was consumed during exponential growth. Afterwards, glucose concentrations were maintained

15

20

25

30

between 0.2 and 1.0 g/L by continuous feeding of a 60% glucose solution. The variable rate feed pump was computer controlled and linked to the dissolved oxygen concentration [pO₂] in the tank by an algorithm. When the [pO₂]fell to 30%, computer control began to automatically adjust the agitation rate to maintain a dissolved oxygen concentration between 25 and 30% [pO₂]. Computer control and data recording were by Braun MFCS software.

In one study, PA284 was grown in fermentors at two temperatures (40°C and 43°C) in the medium described in Table 22. Results of two experiments demonstrated that the highest pantothenate titers at early time points were produced at 43°C. The cell mass approached 150 optical density units at OD₆₀₀ and 56 hours at 43°C, and the pantothenate titers were 21 g/L, 28 g/L and 36 g/L at 36, 48 and 72 hours respectively. In the parallel fermentation at 40°C, the cell mass approached 120 optical density units at OD₆₀₀ and 56 hours, and the pantothenate titers were 18 g/L, 26 g/L and 37 g/L at 36, 48 and 72 hours, respectively.

In another study, PA824 was grown in a fermentor at 43°C in the medium described in Table 23. The cell mass exceeded 160 optical density units at OD₆₀₀ and 36 hours, and the pantothenate titers were 23 g/L, 34 g/L, 37 g/L and 40 g/L at 24, 36, 48 and 60 hours, respectively. In other fermentations, increasing the amount of trace elements in the glucose feed (e.g., increasing the concentration of SM from 1X to 2X) resulted in even higher titers of pantothenate.

EXAMPLE XII: Identification and characterization of the B. subtilis coaA gene product

The annotated version of the *B. subtilis* genome sequence available on the "Subtilist" web site contains no gene labeled as coaA. However a homology search using the protein sequence of *E. coli* pantothenate kinase as a query sequence gave a good match with *B. subtilis* gene yqjS, which is annotated as "unknown; similar to pantothenate kinase." This gene appears to be the penultimate gene in an operon containing five open reading frames (Figure 18). Two of the open reading frames encode proteins which are similar to D-serine dehydratase and to "ketoacyl reductase"; the other two have no known homologies. For the open reading frame corresponding to coaA, there are three possible start codons; each having a possible ribosome-binding site (RBS) associated with it. The three potential coaA ORFs were named coaA1, coaA2, and coaA3, from longest to shortest.

All three potential coal open reading frames were cloned along with their respective RBSs by PCR followed by ligation into expression plasmid pAN229. pAN229 is a low copy vector in E. coli that provides expression from the SP01 phage P_{15} promoter and can integrate by single crossover at bpr with tetracycline selection. A representative resulting plasmid, pAN281, is shown in Figure 19.

To determine if the cloned putative coaA ORFs actually encode a pantothenate kinase activity, several isolates of all three plasmids were transformed into the E. coli strain YH1, that contains the coaA15(Ts) allele. Transformants were streaked to plates incubated at 30° and 43°C to test for complementation of the temperature sensitive allele. All isolates of all three coaA variants, except for one isolate of pAN282, complemented well at 43°C, indicating that all three plasmid constructs encode an active pantothenate kinase. Accordingly, it can be concluded that the B. subtilis yqjS open reading frame codes for an active pantothenate kinase.

EXAMPLE XIII: Deletion of the coaA gene from the B. subtilis genome 15

The coaA gene of B. subtilis (yqjS) was deleted from the chromosome of a B. subtilis strain by conventional means. The majority of the coaA coding sequence was deleted from a plasmid clone and replaced by a chloramphenicol resistance gene (cat), while leaving approximately 1 kb of upstream and downstream sequence to allow homologous recombination within the chromosome, to give plasmid pAN296 (see Figure 17). pAN296 was then used to transform a B. subtilis strain (PY79), selecting for chloramphenicol resistance. The majority of transformants result from a double crossover event that effectively substitutes the cat gene for the coaA gene. The transformed strain containing the coaA deletion -cat insertion grew normally due the 25 presence of a second B. subtilis pantothenate kinase encoding gene described herein.

EXAMPLE XIV: Identification and characterization of a second B. subtilis gene encoding pantothenate kinase activity

As described in detail in the instant specification, in order to maximize pantothenate production, it is necessary to restrict the flow of pantothenate toward 30 Coenzyme A (CoA), for example, by reducing the activity of pantothenate kinase, the first enzyme in the pathway from pantothenate to CoA. After finding that deletion of the coaA gene from the chromosome of B. subtilis is not a lethal event (see Example XIII), it was concluded that B. subtilis must contain a second gene that encodes an active pantothenate kinase, since pantothenate kinase is an essential enzyme activity. 35

A second pantothenate kinase-encoding gene was identified by complementing the *E. coli* strain YH1 (coaA15(Ts)) with a *B. subtilis* gene bank and selecting for transformants that were able to grow at 43°C. Found among the transformants were two families of plasmids that had overlapping restriction maps within each family, but not between the families. As expected, the restriction map of one family was identical to that predicted from the *B. subtilis* genome sequence for the homologue of the *E. coli coaA* gene (which we named coaA also, see above) and surrounding sequences. The other family had a restriction map that was completely non-overlapping with the first.

DNA sequencing of the ends of the cloned inserts from the second family showed that the clones came from a region of the *B. subtilis* chromosome that includes the 3' end of the *fisH* gene, the 5' end of the *sul* gene, and all of the *yacB*, *yacC*, *yacD*, *cysK*, *pabB*, *pabA* and *pabC* genes. None of the open reading frames of these cloned inserts showed homology to any known pantothenate kinase sequences, either prokaryotic or eukaryotic.

15 Several deletions were created through the B. subtilis genomic sequences in the cloned inserts. Each deletion was tested for complementation of the E. coli temperature sensitive pantothenate kinase. In particular, a deletion that removed all DNA between a Stu I site in the cloning vector and a Swa I site in the yacC gene, leaves yacB as the only intact open reading frame in the cloned insert (see Figure 21). This deleted plasmid still 20 complemented the E. coli pantothenate kinase mutant. However, another deletion that removed DNA from the Swa I site in yacC through a Bst1107I site in the (already truncated) fisH gene, could not complement the E. coli pantothenate kinase mutant. From these results, it was concluded that the yacB open reading frame was responsible for the complementation activity. To confirm that yacB is a pantothenate kinase gene, the yacB ORF plus 112 base pairs of downstream flanking sequence was amplified by 25 PCR in two independent reactions and cloned downstream of a constitutive promote to give plasmids pAN341 and pAN342 (Figure 22). Both pAN341 and pAN342 complemented the defect in YH1 at 44°C, while a control plasmid, which has the same backbone, but expresses panBCD instead of yacB did not. This confirmed that the yacB 30 open reading frame was responsible for the complementation of YH1.

As such, a novel gene that encodes pantothenate kinase activity in *B. subtilis* has been discovered that is not related by homology to any previously known pantothenate kinase gene. This gene has been renamed *coaX*, as a second, alternative gene that encodes an enzyme that catalyzes the first step in the pathway from pantothenate to CoaA. Deletion of *coaX* by methods described above for deleting *coaA*, in conjunction

with reduction in the activity of the CoaA enzyme, provides a means to reduce pantothenate kinase activity to the desired level.

Several homologues of the *B. subtilis coaX* gene were identified by homology searching of various publically available databases using the published yacB (coaX) open reading frame sequence and predicted amino acid sequence (as set forth in SEQ ID NOs:84 and 85 respectively). In two cases (Mycobacterium tuberculosis and Streptomyces coelicolor) the homologous coaX genes are adjacent to, or almost adjacent to, pantothenate biosynthetic genes, consistent with these homologs having a role in pantothenate metabolism. The CoaX proteins show no homology to the CoaA family of pantothenate kinases, nor to the eukaryotic family of pantothenate kinases exemplified by PanK of Saccharomyces cerevisiae.

Alignment of the amino acid sequences of several bacterial CoaX homologs with the amino acid sequence predicted from translating the *B. subtilis yacB* ORF described in the published *B. subtilis* genome sequence revealed that the CoaX proteins from other bacteria contained additional amino acid residues at their carboxy-terminal ends. Moreover, these extensions beyond the end of the predicted amino acid for the *B. subtilis* gene product contained two relatively well conserved segments of sequence.

Translation of nucleotide sequences just downstream from the stop codon of the B. subtilis yacB ORF in a different reading frame revealed the existence of amino acid sequences very similar to the carboxy-terminal extensions of the other bacterial CoaX proteins. It is thus believed that an error exists in the published DNA sequence of the B. subtilis yacB ORF sequence that causes a frame shift leading to an artifactual downstream amino acid sequence and premature termination.

The PCR-generated sequences of *B. subtilis CoaX* in pAN341 and pAN342 (described above) contain enough downstream flanking sequence to encode the putative carboxy-terminal extension described above, which is consistent with the result that the clones were functional in the complementation assay. However when the 3' PCR primer was positioned to include only the shorter *yacB* ORF predicted from the published sequence, but not to include the putative carboxy-terminal extension, then the resulting plasmids, pAN329 and pAN330 (similar in structure to pAN341 and pAN342; *see* Figure 22), did not complement the defect in YH1. This result supports the notion that the published *yacB* coding sequence contains a frame-shift error, and that the carboxy-terminal end of CoaX is necessary for pantothenate kinase activity. The predicted correct nucleotide sequence for *B. subtilis coaX* is set forth as SEQ ID NO:19 and the translated amino acid sequence is set forth as SEQ ID NO:9. A multiple

35

sequence alignment of the CoaX amino acid sequences of B. subtilis and 11 homologues thereof is set forth in Figure 23.

EXAMPLE XV: Generation of mutant coaA genes encoding pantothenate kinase having reduced or temperature sensitive activities

This Example describes strategies for modifying the *coaA* gene (*i.e.*, by introducing point mutations) to reduce the activity of pantothenate kinase after *coaX* is deleted from the genome.

10 Cloning and sequencing of the temperature sensitive allele of the E. coli coaA gene.

Two E. coli strains, each exhibiting a different mutant CoaA phenotype, were obtained from the E. coli Genetic Stock Center. Strain DV62 contains the coaA15(Ts) allele, and DV79 contains the coaA16(Fr) mutation. DV62 is temperature sensitive at 43°C and produces a pantothenate kinase that is temperature sensitive. DV79 was obtained by reversion of DV62 to temperature resistance, and it produces a temperature stable, feedback resistant pantothenate kinase activity. Since the DNA sequences of these alleles are not available in the literature, the coaA genes from the two mutant strains were cloned by PCR and sequenced, in addition to a coaA gene from a strain that is wild type at the coal locus, MM294. The PCR primer at the 5' end was designed to include the start codon plus four bases upstream, and added an arbitrarily chosen 20 ribosome binding site (RBS). The three PCR generated fragments were each ligated between the XbaI and BamHI sites of pAN229 to give pAN284 (from coaA15(Ts)). pAN285 (from wild type coaA), and pAN286 (from coaA16(Fr)). pAN229 is a low copy E. coli vector that provides expression from the P₁₅ promoter and that can integrate 25 by single crossover at bpr in B. subtilis with tetracycline selection.

All three plasmids were transformed into the *E. coli* strain YH1 for complementation testing. All three plasmids complemented the temperature sensitive *coaA* mutation in *E. coli* YH1. It is presumed that the *coaA15(Ts)* gene in pAN284 is probably significantly overexpressed relative to the normal chromosomal gene, such that the overproduction compensates for the temperature sensitive defect. Complementation of a defect by overproduction is a well-documented phenomenon in *E. coli*.

The coal coding regions from pAN284, 285, and 286 were subcloned into pGEM7 to give pAN306, 307, and 308, respectively, for DNA sequencing. As expected, the DNA sequence of the insert in pAN307 (from wild type coal) matched the coal sequence from the $E.\ coli$ genome database (GenBankTM). The sequence from pAN306 contains a single base change that causes a S176L substitution (i.e., a Ser \rightarrow

Leu substitution in the amino acid sequence set forth as SEQ ID NO:2). Interestingly, the DNA sequence of the pAN308 insert, derived from the feedback resistant strain, was identical to that derived from its temperature sensitive parent (represented in pAN306). This is in accord with the genetic data that indicates that the reversion of the temperature sensitive mutation occurred at a second site unlinked to the coal gene.

The S176L mutation, predicted to cause the temperature sensitive defect in E. coli pantothenate kinase, changed a serine residue that is conserved in all known or suspected bacterial coaA encoded pantothenate kinases, including that of B. subtilis (see SEQ ID NO:3 and refer to alignment). Based on this, a serine to leucine change at the homologous residue in the B. subtilis pantothenate kinase is predicted to result in either a temperature sensitive enzyme or one which is less active. Accordingly, to produce a mutant B. subtilis coaA gene, this specific change was introduced into the B. subtilis coaA gene. The mutant version is installed in the chromosome of a B. subtilis strain deleted for coaX, for example, and the recombinant microorganism is checked for temperature sensitivity (e.g., reduced growth at 43°C). The mutation is then installed into a pantothenate overproducing strain, preferably a strain deleted for the above mentioned coaX gene by standard methods to give strains favorable for pantothenate production in B. subtilis, i.e., a strain that has reduced pantothenate kinase activity under typical fermentation conditions.

20

30

Additional coaA point mutations resulting in reduced pantothenate kinase activity

Of course it is expected that many other point mutations or combinations of more than one point mutation in *B. subtilis coaA* will also lead to reduced activity. Appropriate mutations can be generated by mutagenic polymerase chain reaction and *in vitro* recombination, and identified by screening for alleles that poorly complement the *E. coli coaA15(Ts)* mutant. An example of such a mutation of this type is a tyrosine to histidine substitution at amino acid 181 of B. *subtilis coaA*, generated by mutagenic polymerase chain reaction (see SEQ ID NO:3 and first line of the alignment of Figure 24).

Isolate pAN282A was derived from the middle-sized B. subtilis coaA open reading frame described in Example XII. pAN282A complemented the E. coli coaA15(Ts) mutant very poorly, but nonetheless at a level that was detectable above background. As was done for the E. coli coaA clones, the open reading frame from pAN282A was subcloned into pGEM7 to give pAN303. The DNA sequence of the insert in pAN303 showed a single base change that led to a tyrosine to histidine amino

30

acid change at the tyrosine corresponding to Y181 of SEQ ID NO:3. This tyrosine residue is conserved in all bacterial coaA genes/homologues present in GenBank (Figure 24). This tyrosine residue and the serine that is altered in the E. coli temperature sensitive pantothenate kinase described above are separated by only three amino acid residues in a region which is highly conserved in bacterial pantothenate kinases whereas the DNA sequence of a second isolate of the middle-sized open reading frame, from pAN282B, was identical to the wild type sequence from the B. subtilis genome sequencing project. The single base change found in pAN303 probably occurred during PCR amplification of the coaA gene. If this variant of coaA2 has sufficient residual biological activity in B. subtilis, it may be useful in the future for providing reduced pantothenate kinase activity.

A preferred plasmid that can serve as a basis for mutagenizing the coaA open reading frame is pAN294 (see e.g., Figure 25 and Example XII). Briefly, mutagenic PCR is performed using pAN294 as a template and variants of coaA having reduced pantothenate kinase activity are screened as described above. Alternatively, mutations such as the one isolated in pAN282A can be installed into pAN294. The desired mutation is then introduced into the chromosome of a B. subtilis strain by transformation with the appropriate pAN294 derivative and selected for chloramphenicol resistance at 5 mg/L. Among the resulting transformants will be isolates that contain the desired mutation.

In a similar fashion, mutations that reduce the activity of the CoaX enzyme can be generated and identified, and such mutations used for optimizing pantothenate production by reducing CoA production as described above.

25 EXAMPLE XVI: Deleting the second pantothenate kinase gene, coaX gene from B. subtilis

With the knowledge gained above concerning the existence and nature of coaX, one can create a deletion of the coaX open reading frame from the B. subtilis chromosome that will remove the encoded activity, and that will not adversely affect the expression of the genes downstream from coaX. In such a deleted strain, the coaA gene will be the only gene that encodes pantothenate kinase.

15

To delete the coaX gene from B. subtilis, plasmid pAN336 (SEQ ID NO:92), which contains upstream and downstream homology for double crossover, was constructed with a kanamycin resistance gene replacing most of the coaX ORF (Figure 26). Strain PY79 was transformed to kanamycin resistance by pAN336, and an isolate confirmed to have resulted from a double crossover by PCR was named PA876. As predicted, deletion of coaX by itself is not lethal for B. subtilis. Furthermore, chromosomal DNA from PA876 would not transform competent PA861 (PY79 ΔcoaA ::cat) to kanamycin resistance. These results indicate that it is the combination of ΔcoaA::cat and ΔcoaX :: kan that is lethal for B. subtilis, confirming that B. subtilis contains two unlinked genes that encode pantothenate kanase, coaA and coaX, and that either gene alone is capable of supplying sufficient pantothenate kinase for a normal rate of growth.

EXAMPLE XVII: Construction of a plasmid designed to allow directed mutagenesis of the *B. subtilis coaA* gene

In order to easily introduce mutated coaA genes into the B. subtilis chromosome, it was necessary to install an antibiotic resistance gene adjacent to the coaA gene. This was accomplished by joining together in the vector pGEM5 three DNA fragments: (1) a 3.4 kb DNA sequence containing 2.5 kb of genomic sequence upstream from coaA and the coal open reading frame(s); (2) a 1.1 kb DNA sequence containing a chloramphenicol resistance gene (cat); and (3) a 1.4 kb DNA sequence comprising a region downstream from the operon that contains coaA. The resulting plasmid, named pAN294, effectively replaces the open reading frame yqjT (the open reading frame just downstream from coaA) with the cat gene, with enough homology flanking both sides of the cat gene to allow double recombination into the B. subtilis chromosome (Figure 25). pAN294 was transformed into B. subtilis strain PY79, selecting for chloramphenicol resistance at 5 mg/l to give strains PA836 and PA837, which are presumably identical. PA836 and 837 were checked by diagnostic PCR to show that the cat gene had integrated by double crossover, as opposed to single crossover. PA836 and PA837 grow normally, leading to the conclusion that the open reading frame yqjT is not essential (i.e., the yqjT open reading frame could be deleted from strains PA836 and PA837 with no significant effect on growth or pantothenate production). Thus, variant alleles (i.e., mutations) of the coad gane can be introduced into pAN294 and the resulting plasmids can be used to introduce the variant alleles into the chromosome of, for example, a B. subtilis strain.

WO 01/21772

EXAMPLE XVIII: Generation of mutant *coaX* genes encoding pantothenate kinase having reduced or temperature sensitive activities

Mutant coaX genes are generated by introducing point mutations into the gene and testing the resulting mutants for the ability to complement the E. coli YH1 strain as described in Example XII. Preferred mutations in the coaX gene sequences are those that encode a substitution of a residue conserved among CoaX sequences from a variety of bacterial sources (e.g., a conserved residue set forth in Figure 23). Alternatively, random mutations in the coaX gene sequence are generated by mutagenic PCR and in vitro recombination and identified by screening for alleles that poorly complement the E. coli coaA15(Ts) mutant.

Mutants so generated (i.e., mutants having reduced coaX activity) can be further engineered such that the endogenous coaA gene is deleted (as described in Example XIII). CoaX reduced-activity mutants can also be further engineered to contain reduced-activity CoaA gene products as described in Example XV.

15

20

25

30

35

10

EXAMPLE XIX: Enhanced Production of Panto-Compounds Using Bacteria Having Deletions in One or More Pantothenate Biosynthetic Enzymes

If the desired panto-compound is not pantothenate, then an appropriate deletion of one or more of the pantothenate biosynthetic genes from a pantothenate overproducing strain will provide a strain that produces said desired panto-compound. In this example, the desired panto-compound is pantoate. Starting with, for example, strain PA236, PA313 or PA824 either one or both of the *panC* and *panD* genes is deleted. In another example, ketopantoate is the desired panto-compound. Starting with, for example strain PA244, PA245 or PA824 one, two or all of the *ilvC*, *panE1*, *panC* and *panD* genes are deleted from the starting strain. If β-alanine is the desired panto-compound, then *panB* and *panC* can be deleted, preferably in a fashion that leaves an in frame fusion of a small portion of the 5' end of *panB* with a small portion of the 3' end of *panC*, from the strain PA221, PA235, PA245, or PA313. In all of the abovementioned examples, the panto-compound producing strain will be a pantothenate auxotroph. Accordingly, the growth medium requires sufficient pantothenate for adequate growth. Vectors designed to overexpress *panD* as described above are then transformed into the above strains to further enhance β-alanine production.

The above-mentioned deletions are accomplished by methods well-known to those skilled in the art, for example, by insertion of an antibiotic resistance gene and removing sufficient sequence from the target gene(s) to inactivate said target gene(s).

Alternatively, removal of targeted sequences is accomplished without simultaneous introduction of an antibiotic resistance gene in said target gene and then introduced by congression (co-transformation with any other appropriate selectable DNA sequence) followed by screening for the loss of function of said target gene by replica plating.

Table 24: Strains (and corresponding phenotypes) for panto-compound production

Name	Pheno type	Drug resist.	panBCD locus	panE locus	ilvD locus	amyE locus	<i>bpr</i> locus	Parent
PA221	Trp-		P26panBCD			•		
PA222			P ₁₅ panBCD					RL-I
PA235			P26panBCD					
PA236			P ₂₆ panBCD	P ₂₆ panE1		1		PA221
PA327	Тгр-		P26panBCD	P26panE1				PA221
PA328	Trp-		P26panBCD	P26panE1				PA235
PA340	Trp-	Spc	P26panBCD	P26panE1		P26ilvBNC		PA327
PA342	Trp-	Spc	P26panBCD	P26panE1		P26ilvBNC		PA328
PA354	Trp-	Spc	P26panBCD	P26panE1	P26ilvD	P26ilvBNC		PA342
PA365	Trp-	Spc, Tet	P26panBCD	P26panE1	P26ilvD	P26ilvBNC	P26panD423	PA354
PA374	Trp-	Spc	P26panBCD	P26panE1	P26ilvD	P26ilvBNC		PA340
PA377	Тгр-	Spc, Tet	P26panBCD	P26panE1	P26ilvD	P26ilvBNC	P26panD423	PA374
PA401	Тгр-		P26panBCD				P26panD423	PA221
PA402	Trp-		P26panBCD				P26panD428	PA221
PA403	Trp-		P26panBCD				P26panD429	PA221
PA404	Trp-	Spc, Tet	P26panBCD	P26panE1		P26ilvBNC	P26panD423	PA340
PA405	Trp-	Spc, Tet	P26panBCD	P26panE1		P26ilvBNC	P26panD423	PA342
PA651	Тгр-	Spc	P26panBC*D	P26panE1	P26ilvD	P26ilvBNC		PA374
PA284	1	Spc, Tet	P26'panBCD	P26panE1	P26ilvD	P26ilvBNC	P26panD423	PA377

Equivalents Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed:

- 1. A method of producing a panto-compound comprising culturing a microorganism which overexpresses at least one *Bacillus* pantothenate biosynthetic enzyme under conditions such that the panto-compound is produced.
 - 2. The method of claim 1, wherein the microorganism overexpresses at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.
- 3. The method of claim 1 or 2, wherein the pantothenate biosynthetic enzyme is selected from the group consisting of ketopantoate hydroxymethyltransferase, pantothenate synthetase, aspartate-α-decarboxylase and ketopantoate reductase.
- 4. The method of any one of claims 1 to 3, wherein the microorganism overexpresses at least two pantothenate biosynthetic enzymes.
 - 5. The method of any one of claims 1 to 3, wherein the microorganism overexpresses at least three pantothenate biosynthetic enzymes.
- 20 6. The method of any one of claims 1 to 5, wherein the panto-compound is selected from the group consisting of pantothenate, pantoate, ketopantoate and β-alanine.
- 7. A method of producing a panto-compound comprising culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism under conditions such that the panto-compound is produced.
 - 8. The method of claim 7, wherein the panto-compound is pantothenate or pantoate.
 - 9. The method of claim 7 or 8, wherein the ketopantoate reductase is bacterial-derived.
- 10. The method of claim 7 or 8, wherein the ketopantoate reductase is derived from *Bacillus*.

PCT/US00/25993

012177242 I V

- 11. The method of claim 7 or 8, wherein the ketopantoate reductase is derived from *Bacillus subtilis*.
- The method of any one of claims 7 to 11, wherein the KPAR-O
 microorganism further overexpresses at least one pantothenate biosynthetic enzyme in addition to overexpressing ketopantoate reductase.
- 13. The method of claim 12, wherein the KPAR-O microorganism further overexpresses at least one of ketopantoate hydroxymethyltransferase, pantothenate
 osynthetase and aspartate-α-decarboxylase.
- 14. A method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism having a deregulated isoleucine-valine (ilv) pathway under conditions
 15 such that pantothenate is produced.
- 15. A method of producing at least 2 g/L pantothenate in a manner independent of aspartate or β-alanine feed comprising culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism under conditions such that
 20 pantothenate is produced.
- 16. A method of producing at least 2 g/L pantothenate in a manner independent of valine or α-ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced.
- 17. A method of producing at least 30 g/L pantothenate in a manner independent of aspartate or β-alanine feed comprising culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism under conditions such that
 30 pantothenate is produced.
- 18. A method of producing at least 30 g/L pantothenate in a manner independent of valine or α-ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced.

- 19. A β -alanine independent high yield production method for producing pantothenate comprising culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly high yield.
- 5 20. The method of any one of claims 14 to 19, wherein the microorganism overexpresses acetohydroxyacid synthetase or is transformed with a vector comprising an *ilvBN* nucleic acid sequence or an *alsS* sequence.
- 21. The method of any one of claims 14 to 19, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase or is transformed with a vector comprising an *ilvC* nucleic acid sequence.
- 22. The method of any one of claims 14 to 19, wherein the microorganism overexpresses dihydroxyacid dehydratase or is transformed with a vector comprising an *ilvD* nucleic acid sequence.
 - 23. The method of any one of claims 19 to 22, wherein the microorganism overexpresses aspartate- α -decarboxylase or is transformed with a vector comprising a *panD* nucleic acid sequence.
 - 24. The method of any one of claims 14 to 23, wherein the microorganism further has a deregulated pantothenate biosynthetic pathway.
- 25. The method of any one of claims 14 to 24, wherein the microorganism further has at least one mutant gene selected from the group consisting of a mutant avtA gene, a mutant ilvE gene, a mutant ansB gene and a mutant alsD gene.
- The method of claim 24, wherein the microorganism overexpresses any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate
 synthetase and aspartate-α-decarboxylase.
 - 27. The method of claim 24 or 26, wherein the microorganism is transformed with a vector comprising a *panBCD* nucleic acid sequence or a vector comprising a *panE1* nucleic acid sequence.

PCT/US00/25993

- 28. The method of any one of claims 14 to 16 and 19 to 27, wherein pantothenate is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 5 29. The method of claim 20, wherein the microorganism overexpresses acetohydroxyacid synthetase derived from *Bacillus* or is transformed with a vector comprising an *ilvBN* nucleic acid sequence or an *alsS* nucleic acid sequence derived from *Bacillus*.
- 10 30. The method of claim 21, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase derived from *Bacillus* or is transformed with a vector comprising an *ilvC* nucleic acid sequence derived from *Bacillus*.
- 31. The method of claim 22, wherein the microorganism overexpresses dihydroxyacid dehydratase derived from *Bacillus* or is transformed with a vector comprising av *ilvD* nucleic acid sequence derived from *Bacillus*.
- 32. The method of claim 23, wherein the microorganism overexpresses aspartate-α-decarboxylase derived from Bacillus or is transformed with a vector
 comprising a panD nucleic acid sequence derived from Bacillus.
 - 33. The method of claim 24 or 26, wherein the microorganism overexpresses any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate-α-decarboxylase derived from *Bacillus*.
 - 34. The method of claim 27, wherein the vector comprises a panBCD nucleic acid sequence or a panE1 nucleic acid sequence derived from Bacillus.
- 35. A method of producing a panto-compound comprising contacting a composition comprising at least one pantothenate biosynthesis pathway precursor or isoleucine-valine biosynthesis pathway precursor with at least one isolated *Bacillus* enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate-α-decarboxylase, under conditions such that the panto-compound is produced.

- 36. A method of producing β -alanine comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that β -alanine is produced.
- 5 37. The method of claim 36, wherein the AαD-O microorganism has a mutation in a nucleic acid sequence encoding a pantothenate biosynthetic enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase and pantothenate synthetase.
- 38. A method of producing β-alanine comprising contacting a composition comprising aspartate with an isolated *Bacillus* aspartate-α-decarboxylase enzyme under conditions such that β-alanine is produced.
- 39. A method for enhancing production of a panto-compound comprising culturing a mutant microorganism having a mutant *coaX* gene under conditions such that the panto-compound production is enhanced.
 - 40. The method of claim 39, wherein said recombinant microorganism has a mutant *coaA* gene.
 - 41. A method of producing a panto-compound comprising a pantothenate kinase mutant microorganism under conditions such that the panto-compound is produced at a significantly high yield.
- 25 42. The method of claim 41, wherein said mutant microorganism has a mutant coaA gene.
 - 43. The method of claim 41, wherein said mutant microorganism has a mutant *coaX* gene.
 - 44. The method of claim 41, where said mutant microorganism has a mutant coaA and coaX gene.
- 45. The method of any one of claims 39 to 44, wherein said panto-compound is selected from the group consisting of ketopantoate, pantoate or pantothenate.

- 46. The method of any one of claims 39 to 44, wherein said panto-compound is pantothenate.
- 47. The method of any one of claims 39 to 44, wherein said panto-compound is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 48. The method of any one of claims 39 to 44, wherein said recombinant microorganism further has a deregulated pantothenate biosynthetic pathway or further has a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.
 - 49. The method of claim any one of claims 39 to 44, wherein said recombinant microorganism further overexpresses panD and panE.
- 15 50. The method of any one of claims 39 to 44, wherein said recombinant microorganism further has at least one mutant gene selected from the group consisting of a mutant avtA gene, a mutant ilvE gene, a mutant ansB gene and a mutant alsD gene.
- 51. A method for enhancing production of a panto-compound comprising culturing a microorganism that has a deregulated pantothenate biosynthetic pathway and that also has a mutation that results in reduced pantothenate kinase activity under conditions such that the panto-compound production is enhanced.
- 52. A method for identifying compounds which modulate pantothenate
 25 kinase activity comprising contacting a recombinant cell expressing pantothenate kinase
 encoded by the coaX gene with a test compound and determining the ability of the test
 compound to modulate pantothenate kinase activity in said cell.
- 53. The method of claim 52, wherein said cell further comprises a mutant 30 coaA gene encoding a pantothenate kinase having reduced activity.
 - 54. The method of any one of claims 1 to 51, wherein the microorganism is Gram positive.
- 35 55. The method of any one of claims 1 to 51, wherein the microorganism is Gram negative.

PCT/US00/25993

25

30

- 56. The method of any one of claims 1 to 51, wherein the microorganism is a microorganism belonging to a genus selected from the group consisting of *Bacillus*, *Cornvebacterium*, *Lactobacillus*, *Lactococci* and *Streptomyces*.
- 5 57. The method of any one of claims 1 to 51 and 54 to 56, wherein the microorganism is of the genus *Bacillus*.
 - 58. The method of any one of claims 1 to 51 and 54 to 57, wherein the microorganism is *Bacillus subtilis*.

10
59. The method of any one of claims 1 to 13, 35,

- 59. The method of any one of claims 1 to 13, 35, 39 to 51 and 54 to 58, further comprising recovering the panto-compound.
- 60. The method of any one of claims 14 to 34 and 54 to 58, further comprising recovering the pantothenate.
 - 61. The method of any one of claims 1 to 14, 35, 39 to 46, 48 to 51 and 54 to 59, wherein the panto-compound is produced at a level greater than 2 g/L.
- 20 62. A recombinant microorganism which overexpresses at least one *Bacillus* pantothenate biosynthetic enzyme.
 - 63. The recombinant microorganism of claim 62, which overexpresses at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.
 - 64. The recombinant microorganism of claim 62 or 63, wherein the pantothenate biosynthetic enzyme is selected from the group consisting of ketopantoate hydroxymethyltransferase, pantothenate synthetase, aspartate-α-decarboxylase and ketopantoate reductase.
 - 65. The recombinant microorganism of any one of claims 62 to 64, wherein the pantothenate biosynthetic enzyme is ketopantoate reductase.
- 66. A recombinant microorganism which overexpresses aspartate-α 35 decarboxylase and has a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.

PCT/US00/25993

- 67. A recombinant microorganism having a mutant coaX gene, said mutant coaX gene encoding reduced pantothenate kinase activity in said microorganism.
- 68. The recombinant microorganism of claim 67 further having a mutant
 5 coaA gene, said mutant coaA gene encoding reduced pantothenate kinase activity in said microorganism.
- 69. A recombinant microorganism having a mutant coaX gene and optionally having a mutant coaA gene, said mutant microorganism having reduced pantothenate

 10 kinase activity as compared to a microorganism having wild-type coaA and coaX genes.
 - 70. A recombinant microorganism comprising a vector comprising an isolated *coaX* gene.
- 15 71. A recombinant microorganism that overproduces a panto-compound, the microorganism having a deregulated pantothenate biosynthetic pathway and having at least one mutation that results in a decrease in the capacity of the microorganism to synthesize Coenzyme A (CoA).
- The recombinant microorganism of claim 71, having at least one mutation that results in a reduced level of pantothenate kinase activity.
- 73. The recombinant microorganism of claim 72, having a mutation in a coaA gene, or homologue thereof, that results in a reduced level of CoaA enzyme 25 activity.
 - 74. The recombinant microorganism of claim 72, having a mutation in a coaX gene, or homologue thereof, that results in a reduced level of CoaX enzyme activity.
 - 75. The recombinant microorganism of claim 72, having a mutation in a coaA gene, or homologue thereof, and having a mutation in a coaX gene, or homologue thereof, the mutations resulting in reduced levels of CoaA enzyme activity and reduced CoaX enzyme activity.
 - 76. The recombinant microorganism of any one of claims 66 to 70 which further has a deregulated pantothenate biosynthetic pathway.

35

30

- 77. The recombinant microorganism of any one of claims 62 to 65 and 67 to 75, further having a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.
- 5 78. The recombinant microorganism of any one of claims 62 to 77, which is Gram positive.
- 79. The recombinant microorganism of claim 78 belonging to a genus selected from the group consisting of Bacillus, Cornyebacterium, Lactobacillus,
 10 Lactococci and Streptomyces.
 - 80. The recombinant microorganism of claim 79 belonging to the genus *Bacillus*.
- 15 81. The recombinant microorganism of claim 80 which is *Bacillus subtilis*.
- 82. A recombinant microorganism selected from the group consisting of PA221, PA235, PA236, PA313, PA410, PA402, PA403, PA411, PA412, PA413, PA303, PA327, PA328, PA401, PA340, PA342, PA404, PA405, PA374, PA354,
 20 PA365, PA377, PA651 and PA824.
 - 83. A recombinant vector for use in the production of panto-compounds comprising a nucleic acid sequence which encodes at least one *Bacillus* pantothenate biosynthetic enzyme operably linked to regulatory sequences.

84. The vector of claim 83, comprising a nucleic acid sequence which encodes at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.

- 85. The vector of claim 84, wherein the nucleic acid sequence encodes at least one of ketopantoate hydroxymethyltransferase, pantothenate synthetase, aspartate-α-decarboxylase and ketopantoate reductase.
 - 86. A recombinant vector comprising at least one nucleic acid sequence selected from the group consisting of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27,
 5 SEQ ID NO:29 and SEQ ID NO:59.

25

PCT/US00/25993

10

15

25


30

- 87. The vector of claim 84, wherein the nucleic acid sequence encodes ketopantoate reductase.
- 88. A vector comprising a mutant *coaX* gene, said mutant encoding a pantothenate kinase enzyme having reduced activity.
 - 89. A vector comprising an isolated coaX gene.
 - 90. A vector comprising an isolated Bacillus coaX gene.

91. A vector comprising an isolated Bacillus subtilis coaX gene.

- 92. The vector of any one of claims 86 and 89 to 91, which further comprises regulatory sequences.
- 93. The vector of any one of claims 83 to 85, 87 and 92, wherein the regulatory sequences comprise a constitutively active promoter.
- 94. The vector of claim 93, wherein the constitutively active promoter comprises P_{veg} (SEQ ID NO:41), P_{15} (SEQ ID NO:39) or P_{26} (SEQ ID NO:40) sequences.
 - 95. The vector of claim 83, wherein the regulatory sequences comprise at least one artificial ribosome binding site (RBS).
 - 96. The vector of claim 95, wherein the artificial RBS comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56 and SEQ ID NO:57.
 - 97. A vector selected from the group consisting of pAN004, pAN005, pAN006, pAN236, pAN423, pAN428, pAN429, pAN441, pAN442, pAN443, pAN251, pAN267, pAN256, pAN257, pAN263, pAN240, pAN294, pAN296, pAN336, pAN341 and pAN342.
 - 98. A recombinant microorganism comprising the vector of claim 86 or 93.

- 99. An isolated nucleic acid molecule which encodes at least one *Bacillus* pantothenate biosynthetic gene.
- 100. The isolated nucleic acid molecule of claim 99 which encodes at least one *Bacillus subtilis* pantothenate biosynthetic gene.
 - 101. The isolated nucleic acid molecule of claim 99 or 100 which encodes ketopantoate reductase.
- 10 102. An isolated *Bacillus* pantothenate biosynthetic enzyme polypeptide.
 - 103. An isolated *Bacillus subtilis* pantothenate biosynthetic enzyme polypeptide.
- 15 104. An isolated *Bacillus* ketopantoate reductase polypeptide.
 - 105. An isolated Bacillus subtilis ketopantoate reductase polypeptide.
 - 106. An isolated *Bacillus* aspartate-α-decarboxylase polypeptide.
- 20
 - 107. An isolated Bacillus subtilis aspartate-α-decarboxylase polypeptide.
 - 108. An isolated nucleic acid molecule comprising a mutant coaX gene.
- 25 109. An isolated nucleic acid molecule comprising a coaX gene.
 - 110. An isolated pantothenate kinase protein encoded by a coaX gene.



☱

Figure 2. Plasmid pAN240, containing sequences ligated upstream of the P_{26} panBCD cassette, equivalent to the integrated version in strain PA221.





Figure 3A Plasmid pAN004, containing the panBCD operon expressed from P26 and RBS1.



TURNOUS AUD 012177282 I >

Figure 35 Plasmid pAN006, containing the panBCD operon expressed from P26 and RBS2.




Figure 4 Plasmid pAN236, containing an integratable and amplifiable P26-RBS2-panE1 expression cassette.




Figure 5 Construction of pAN423





Figure 6 Construction of pAN426 and pAN427.



ו באבדדינית מוני מוכרמייי

Figure 7 Construction of pAN428 and pAN429.




Figure 8. Construction of pAN431.




Figure 9. Construction of pAN441.




Figure 10. Construction of p.4N440.




Figure [Structure of pAN251, a plasmid designed to integrate a single copy of P₂₆ panE1 at the panE1 locus by double crossover.




Figure 12 Structure of pAN267, a plasmid designed to stably integrate a P₂₆ ilvBNC cassette at the amyE locus.




Figure 13 Structure of pAN257, a clone of B. subtilis ilvD in a low copy vector.




Figure 14 Structure of pAN263, designed to stably integrate a single copy of P₂₆ ilvD at the ilvD locus.





Figure 15 Structure of pAN261, designed to disrupt the B. subtilis ilvD gene with the cat gene.



minnonn inn arasmani.

17/31

Figure 16




Figure 17 Structure of pAN296, designed to delete most of the B. subtilis coaA gene and substitute a chloramphenical resistance gene.




Figure (8 Structure of the B. subtilis chrumosome in the region of the coad gene. The scale is in base pairs and the significant open reading frames are shown by the open arrows.

5250 , I 28S III buiH 2000 I uts I isq. 4750 <u>}</u>[b, I XISB 2500 III BIO 4250 4000 JOES. I Hms8 3750 III briH ·yaïT Xps I Sacı 11 168 -coaA3 coaA2 900 300 300 coaA1 2750 2500 2250 11 169 2000 III buiH 1750 I dia 1500 1250 <u>8</u> I sqA 750 Smal 1 ems 250 II 3tz8 1445

Figure 19 Structure of pAN281, a plasmid for expressing B. subtilis coaA after integration at the bpr locus. pAN282 and pAN283 have similar structures.




FIG.20A

Alignments
u .
_
Ψ
E
2
ö
ă
7
٠,
Sequence
υ
C
O
3
Ē
χ.
χ.
"
٠.
a)
-
Q.
-
٠
Multiple
€
_
7
٠
_
~
3
3
3
AL &
TAL
STAL W
USTAL W
LUSTAL W
CLUSTAL W
CLUSTAL W

Sequence type explicitly set to Protein	: to Protein	
Sequence 1: spi09X795!M.leprae	.ae 312 aa	
	olor	
	is	
<u>.</u>		
;	316 аа	
SDIQ9X795IM.leprae	MPRLSEPSPYVEFDRKQWRALRMSTPLALTEEELIGLR	
spl053440 M.tuberculosis	MSRLSEPSPYVEFDRRQWRALRMSTPLALTEEELVGLR	
sp10867791S.coelicolor	MISPVPSIPRSAHRÜRPEATPIVOLIRFENSALRONIFLFLIABERGRUNG	
splP447931H.intluenzae	ALTACIACIONE LA CONTRACTOR DE LA CONTRAC	
SpiPi50441E.Coll	SWENE STORES TO THE SWENE SWEN	
sp/P54556/B.subtilis	5	
spiQ9X795 M.leprae	GLGEQI DLLEVEEVYLPLARLI HLQVAARQRLFAATAEFLGEPQONPGRP	
spi0534401M.tuberculosis	GLGEQIDLLEVEEVYLPLARLIHLQVAARQRLFAATAEFLGEPQQNPDRP	
sp10867791S.coelicolor	GLGDVIDLDEVRDIYLPLSRLLNLYVGATDGLRGALNTFLGEQGSQSG	
sp P44793 H.influenzae	GENEDLSLDEVSTIYLPLTRLINYYIDENLHRQTVLHRFLGRNNAK	
Sp P15044 E.col1	GINEDLSLEEVAEIYLPLSRLLNFYISSNLRRQAVLEOFLGTNGQR	
sp1P54556 B.subtilis	GLNDYLSVEEVETIYIPLVRLLHLHVKSAAERNKHVNVFLKHPHSAK	
	the second of th	
00000 M-300000	VPETTCVAGSVAVGKSTTARVLOALLABWDHHTRVDLVTTDGFLYPNAEL	
Spigovojim reprac	VPFTTCVACSVAVCKSTTARVLOALLARWDHHPRVDLVTTDGFLYPNAEL	
sp(033440 m.cubarcarosts	TPFVICVAGSVAVGKSTVARLLÖALLSRWPEHPRVELVTTDGFLLPTREL	
TOTOTTEND: CIGITATION OF	TELETICONNOCKSTORE TO STORE THE PRESENT THE SERVICE TO THE SERVICE	
spir44/931H.Intinenzae	INCHOR COURT LONG AND	
sp!P15044 E.col1	IPYIISIAGSVAVGKSTTAKVLQALLSKWPBARKVBLI 1 JUGT LAFINGVL	
spiP545561B.subtilis	IPFIIGIAGSVAVGKSTTARILOKLLSRLPDRPKVSLITTDGFLFFIAEL	
	一条,是中央中央的,是一个一个一个一个一个一个一个一个一个一个一个一个一个一个一个一个一个一个一个	

FIG. 20E

GRRULMHRKGFPESYNRRALMRFVTSVKSCADYACAPVYSHLRYDTIPGA QRRNLMHRKGFPESYNRRALMRFVTSVKSGSDYACAPVYSHLHYDIIPGA EARGLMSRKGFPESYDRRALTRFVADIKAGKAEVTAPVYSHLIYDIVPDQ KQDNLLQKKGFPVSYDT PKLIRFLADVKSGKSNVTAPIYSHLTYDIIPDK KERGLMKKKGFPESYDMHRLVKFVSDLKSGVPNVTAPVYSHLIYDVIPOG KKNNMMSRKGFPESYDVKALLEFLNDLKSGKDSVKAPVYSHLTYDREEGV ::::*********************************	KHVVRHPDILILEGLNVLQTGPTLMVSDLFDFSLYVDARIQD EQVVRHPDILILEGLNVLQTGPTLMVSDLFDFSLYVDARIED RLVVRRPDILILVEGLNVLQPALPGKDGRT-RVGLADYFDFSVYVDARTED FDVVDKPDILILEGLNVLQTGNNKTD-QTFVSDFVDFSIYVDAEEKL DKTVVQPDILILEGLNVLQSGMDYPHDPH-HVFVSDFVDFSIYVDAPEDL FEVVEQADIVILEGINVLQSPTLEDDRENPRIFVSDFFDFSIYVDAEESR .* : .*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:*:	IEQWYVSRFLAMRGTAFADPESHFHHYSALTDSKAIIAAREIWRSINRPN IEQWYVSRFLAMRTTAFADPESHFHHYAAFSDSQAVVAAREIWRTINRPN IERWYLNRFRKLRATAFQNPSSYFRKYTQVSEEEALDYARTTWRTINRPN LKEWYIKRFLKFRESAFNDPNSYFKHYASLSKEEAIATASKIWDEINGLN LQTWYINRFLKFREGAFTDPDSYFHNYAKLTKEEAIKTAMTLWKEINWLN IFTWYLERFRLLRETAFQNPDSYFHKYDLSDQEADEMAASIWESVNRPN IFTWYLERFRLLRETAFQNPDSYFHKFKDLSDQEADEMAASIWESVNRPN : **: ** : * * : * : * : * : * : * : *	LVENILPTRPRATLVLRKDADHSINRLRLRKL LVENILPTRPRATLVLRKDADHSINRLRLRKL LVENVAPTRGRATLVLRKGPDHKVQRLSLRKL LNQNILPTRERANLILKKGHNHQVELIKLRK- LKQNILPTRERASLILTKSANHAVEEVLRK- LYENILPTKFRSDLILRKGDGHKVEEVLVRRV * :*: *: *: *: *: *: *: *: *: *: *: *: *
sp Q9X795 M. leprae sp O53440 M. tuberculosis sp O86779 S. coelicolor sp P44793 H. influenzae sp P15044 E. coli sp P54556 B. subtilis	sp Q9X795 M. leprae sp O53440 M. tuberculosis sp O86779 S. coelicolor sp P44793 H. influenzae sp P54556 B. subtilis	spl09X795 M.leprae spl053440 M.tuberculosis spl086779 S.coelicolor splP4793 H.influenzae splP15044 E.coli	splQ9X7954M.leprae splO534401M.tuberculosis splO867791S.coelicolor splP447931H.influenzae splP150441E.coli


14000

() Juil4

I PS 1405 00CT il aniH-I ISN 1 4011158 12000 I UIM Sacı -Kpn I Il aniH-Swa I - Bst1107 I -8-III buiH-10000 III puiHi uiS-1 BMS III buiH-**8** Capasi 1835 Hinc it U aniH-8000 III PUIH-II 168-Sall-Sall 1835-Csp45 I · Csp45 I 900 Cap42 I 000 Il oniH-Bam HI - Bgl II 115d~ I ISd---C20421 2000 III puiji. Cap45 1 400 1962 I MIS. I isN ~ ! Hmsg. III buiH -900 II aniH I ISd r, sa L 2000 IMS. II aniH 1185 -<u>8</u>. II aniH-1 805 ...-

Figure 21

Figure 22 Structure of pAN341 and pAN342, two independent PCR- derived clones of yacB (renamed coaX).



CLUSTAL W (1.7) Multiple Sequence Alignments

rotein 258 aa Seq.8: sp O51477 B.burgdorferi 262 aa 212 aa Seq.9: sp P74045 Synechocystis 257 aa 246 aa Seq.10: sp O25533 H.pylori 223 aa 265 aa Seq.11: sp O67753 A.aeolicus 229 aa 267 aa Seq.12: sp Q9RX54 D.radiodurans 262 aa 263 aa Seq.13: WIT RCA0330 C.acetobutylicum 250 aa 273 aa Seq.14: WIT RRCO2473 R.capsulatus 258 aa	NKRAAFMLLETRSVLKVILVLDVGNTNTVLGVYHDGKLEYHWRIE NKRAAFMLLEFLRSVLKVILVLDVGNTNTVLGLYDGEDIVEHWRIS
Sequence type explicitly set to Protein Sequence format is Pearson Seq. 1: B.subtilis!Coax!SEQNO 9 258 as Seq.2: dbj!BAA21476.1 D.vulgaris 212 as Seq.3: gb!AAD35964.1 T.maritima 246 as Seq.4: pir!T36391 S.coelicolor 265 as Seq.5: sp!Q45338 B.pertussis 267 as Seq.6: sp Q06282 M.tuberculosis 272 as Seq.7: sp O83446 T.pallidum 273 as	B. subtilis Coax SEQIDNO_9 WIT RCA0330 C.acetobutylicum pir T3639 S.coelicolor splO06282 M.tuberculosis WIT RRC02473 R.capsulatus dbj BAA21476.1 D.vulgaris splO9RX54 D.radiodurans gb AAD35964.1 T.maritima splO9RX54 D.radiodurans gp O9RX54 D.radiodurans splO9RX51 R.pallidum splO51477 B.burgdorferi splO67753 A.aeolicus splO67753 A.aeolicus splO25533 H.pylori splO45338 B.pertussis

----LGIQKEIFYISVNEE

WLATLPRRP----Q----RALGVNVAGLARGEAIA

AKEDIKR-----

YNPKSAQLP------VLLGKVPLMLASVVPE

ILRSLFDHS----GLMFEQIDGIIISSVVPPIMFALER QVMNLFQQD----KLDPTLVEGVIISSVVPNIMYSLEH **LLQGLMGMHPLLGDELGDGIDGIAICATVPSVLHELRE** WLNTLMQLK----GLQGRISEAIISSTAPRVVFNLRV RLLEVLRHAG----LGPADVGACVASSVVPGVNPLIRR HLHPLLG-----DAMREIKGIGVASVVPTQNTVIER LIHALCERAG----VGRASLRDAFISSVVPVLTKTIAD FFEENFOFN----VN---K-VFISSVVPILNETFKN EEFPKLK------ALGISVKQSFSEKVRG

TIDGLIG-----EDSERLTGTAALSTVPSVLHEVRI

QLHGLFTLA----GAP-IPRAAVLSSVAPPVGENYAL

subtilis Coax SEQIDNO 9	TSRHKTEDEFGM
TIRCA03301 C.acetobutylicum	TDVLRSADEYGI
r T36391 S.coelicolor	TOSRRTADELAV
010062821M.tuberculosis	TESEVTADELAL
TIRRC024731R.capsulatus	TDHRRTADEYFV
oj BAA21476.11D.vulgaris	TDPGQTTDSIGL
OlQ9RX54 D.radiodurans	INREMLPODLAL
AAD35964.1 T.maritima	TGVFQTEDELFS
10834461T.pallidum	PDARKTQDEYSL
10514771B.burgdorferi	TNLMLRYDEVYS
(067753 A.aeolicus	DFLKLSHEEFLKI
P74045 Synechocystis	SGNAPLQTWVTD
10255331H.pylori	IHFAQNYQLFSS/
1Q453381B.pertussis	AFDNLDLDALGR

MIRKYFKINPLVVG-PG-IKTGINIKYDNPKEVGADRIVNAVAAHEIYK-VTRRYYGDVPAVLVEPG-VKTGVPILTDHPKEVGADRIINAVAAVELYG-MLDQYWPSVPHVLIEPG-VRTGIPLLVDNPKEVGADRIVNCLAAYDRFR-AVAQISGVQPVVFGPWAYEHLPVRIPEPVRAEIGTDLVANAVAAYVHFR-MCTKYFHIEPQIVG-PG-MKTGLNIKYDNPKEVGADRIVNAVAAIHLYG-LCNRYFDCRPYVVGKPG-CELPVAPRVDPGTTVGPDRLVNTVAGYDRHG-VI FSFFKI KPLFIGFOLNYOLTFNPYKSOKFLLGSDVFANLVAAI ENYS-QTEVWRVYQPKILTLKN---LPLVNLYP---SFGIDRALAGLGTGLTYG-NEKALLNCYPNAKNIAG--FFHLETDYVG---LGIDRQMACLA---VN--ATLRAGGCDIRWLRAQP-LAMGLRNGYRNPDQLGADRWACMVGVLARQPS ACERYL--YRKLLFAPGOIAIPLONRYERPAEVGADRLVAAYAARRLYP-ALKRHEMIDAFAVSAEN--LPDVTVELDTPGSVGADRLCNLFGAEKYLG-FSQKYFIIISPIWVKAKN---GCVKWNVKNPSEVGADRVANVVAFVKEYG-KIPKIK-----FLKKEN---FPIQVDYKTPETLGTDRVALAYSAKKFYG-

WIT!RCA03301|C.acetobut \overline{y} 11cum WIT!RRC02473|R.capsulatus db | BAA21476.11D.vulgaris sp|006282|M.tuberculosis gb|AAD35964.1|T.maritima B. subtilis | Coax | SEQIDNO pir | T36391 | S. coelicolor splQ9RX541D.radiodurans splP74045|Synechocystis spl051477 B. burgdorferi splQ453381B.pertussis spl083446|T.pallidum spi0677531A.aeolicus |025533|H.pylori MIJORIAN SEPTION OF SE gb

WIT!RCA03301|C.acetobutylicum

B. subtilis|Coax|SEQIDNO

dbjiBAA21476.11D.vulgaris WIT!RRC02473!R.capsulatus

gb|AAD35964.1|T.maritima

sp10514771B.burgdorferi splP74045|Synechocystis

spl0834461T.pallidum spi0677531A.aeolicus sp|025533|H.pylori sp|045338|B.pertussis

spl@9RX54lD.radiodurans

spi006282IM.tuberculosis

pir (1363911S.coelicolor

KA--AIVVDFGSSICVÖVVSAKGEFLGGAIAPGVQVSSDAAAARSAALRR GPRSLVSVDFGTATTFDCVEG-GAYLGGLICPGVLSSAGALSSRTAKLPR KN--VVVISAGTALVIDLVLE-GKFKGGFITLGLGKKLKILSDLAEGIPE NG---VVVDAGSAITIDLIKE-GKHLGGCILPGLAQYIHAYKKSAKILEQ NP--LIVVDFGTATTYCYIDENKQYMGGAIAPGITISTEALYSRAAKLPR RS--LIIIDFGTATTFCAVRENGDYLGGAICPGIKVSSEALFEKAAKLPR GP--AIVVÖFGTATTFDAVSARGEYIGGVIAPGIEISVEALGVKGAQLRK GD--LIVVOFGTATTFDVVAPDGAYIGGVIAPGVNLSLEALHMAAAAALPH **GLDYAVVVDFGTSTNFDVVGRGRRFLGGILATGAQVSADALFARAAKLPR** KN--GIIIDMGTATTVDLVVN-GSYEGGAILPGFFMMVHSLFRGTAKLPL SA--CVVVDCGTALTFTAVDGTGLIQGVAIAPGLRTAVQSLHTGTAQLPL FEN-VLVVDLGTACTIFAVSRODGILGGIINSGPLINFNSLLDNAYLIKK FP--CLVVDGGTALTITGFDQDKKLVGGAILPGLGLQLATLGDRLAALPK UHPPLLVASFGTATTLDTIGPONVFPGGLILPGPAMMRGALAYGTAHLPL

> WIT! RCA03301|C.acetobutylicum B.subtilis|Coax|SEQIDNO_9 WIT!RRC02473!R.capsulatus dbjlBAA21476.11D.vulgaris spi0062821M.tuberculosis gb|AAD35964.1|T.maritima pir | T36391|S. coelicolor spi@9RX54|D.radiodurans spi051477|B.burgdorferi spl P740451Synechocystis splQ453381B.pertussis spi083446|T.pallidum spi0677531A.aeolicus sp10255331H.pylori

VELIKPAY---AICKNTISSIQSGIVYRYLRQVKYLFEKLKENLPDGRRT IEVARPRS---VIGKNTVEAMÓSGIVYGFAGQVDGVVNRMARELADD--P VELARPRS----VVGKNTVECMQAGAVFGFAGLVDGLVGRIREDVSGFSVD IEITRPDN---IIGKNTVSAMQSGILFGYVGQVEGIVKRMKWQAKQDLK-ITLQAPET---AIGKNTVHALQSGLVFGYAEMVDGLLRRIRAELPGE---VEVKPADF---VVGKDTEENIRLGVVNGSVYALEGIIGRIKEVYGDLP--VDVTKPQG----VIGTNTVACIQSGVYWGYIGLVEGIVRQIRMERDRP--adglvady-----Pidthqaiasgiaaaqagaivrqwlagrqgrygqap-FPISTPNN----LLERTTSGSVNSGLFYQYKYLIEGVYRDIKQMYKKK-PFKALÖSL--EVLPKSTRDAVNYGMVLSVIACIQHLAK--NQK----LEMDQLTELPÖRWALDTPSAIFSGVVYGVLGALQSYLQDWQKLFPGAvplalpds---vlgkótthavóagvvrgtleviramiaócókelgcr-FFPEEVEI---FLGRSTRECVLGGAYRESTEFIKSTLKLWRKVFKRK-ISLEVEEDS-PVIGRSTTTSLNHGFIFGFAAMTEGVLAA----

FIG.23D

VIATGGLAPLIANESDCIDIVDPFLT1KGLELI DDVTVIATGGLAPWILGESSVIDEHEPWLTLMGLRLV HDVAIVATGGTAPLLLPELHTVDHYDQHLTLQGLRLV	
VIATGGLAPLI RTSLVLATGGLARLI DDVTVIATGGLAPWY HDVAIVATGHTAPLI	YERNRVGSV YERNVSRM FERNLEVQRGRLKTAR FDYNKGLGA WASRSEVR CFGD ARLVPTSLLPPATVSGSSGN GNSIDFKFVN LYLYHRI
B.subtilis Coax SEQIDNO_by WIT RCA03301 C.acetobutylicum pir T36391 S.coelicolor sp O06282 M.tuberculosis WIT RRC02473 R.capsulatus db IBAA21476.1 D.vulgaris sp Q9KX54 D.radiodurans gb AAD35964.1 T.maritima sp O81477 B.burgdorferi sp O51477 B.burgdorferi sp O67753 A.aeolicus sp O753 A.aeolicus sp O25533 H.pylori sp Q45338 B.pertussis	B. subtilis Coax SEQIDNO_9 WIT RCA03301 C.acetobutylicum pir T36391 S.coelicolor sp O06282 M.tuberculosis WIT RRC02473 R.capsulatus db IBA21476.1 D.vulgaris sp Q9RX54 D.radiodurans gb AD35964.1 T.maritima sp O81471 B.burgdorferi sp O5753 A.aeolicus sp O67753 A.aeolicus sp Q25533 H.pylori sp Q45338 B.pertussis

Figure 24 Alignment of a portion of the amino acid sequences of several known or suspected pantothenate kinases. The residues that are mutaled in E. coli coa A15(Ts) and B. subtills coa from plasmid pAN282A are indicated below and above the alignment, respectively. The coordinate given in the left margin for the B. subtills protein refers to the coa A1 open reading frame.

Majority	B. subtilis CoaA1 E. coli CoaA H. influenzae CoaA M. leprae CoaA M. tuberculosis CoaA S. coelecolor CoaA
K D N V T A P V Y S H L I Y D I I P G A	Y D R E E G V Y D V I P D G Y Y D T I P D K X Y D T I P G A Y Y D I V P D Q A Y D I V P D Q A
O	000000
ما	8 4 4 4 4
-	
-	
. .	
≻ =	<u> </u>
-	**************************************
اد -	
I	
S	10000000 D
>	
>	>> - >>>
ما	
⋖	44444
-	X F FOOF
>	>>><
Z	ωZZ>>ω
۵	DOOV
\mathbf{z}	K D S V K A P V Y S H K S N V T A P V Y S H K S N V T A P Y Y S H X D Y A C A P V Y S H X D Y A C A P V Y S H X A C A P V Y S H
	168 167 169 169 179

Figure 25 Structure of pAN294, a plasmid for integrating mutagenized B. subtilis coaA at its native locus.





Figure 26 Structure of pAN336, a plasmid designed to delete B. subtilis coaX from the chromosome and replace it with a kanamycin resistance gene.



WO 01/21772 PCT/US00/25993

-1-

SEQUENCE LISTING

<110> OMNIGENE BIOPRODUCTS

<120> METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS.

<130> BGI-141CPPC

<140>

<141>

<150> USSN 09/400,494

<151> 1999-09-21

<150> USSN 60/210,072

<151> 2000-06-07

<150> USSN 60/221,836

<151> 2000-07-28

<150> USSN 60/221,836

<151> 2000-08-24

<160> 94

<170> PatentIn Ver. 2.0

<210> 1

<211> 311

<212> PRT

<213> Haemophilus influenzae

<400> 1

Met Glu Phe Ser Thr Gln Gln Thr Pro Phe Leu Ser Phe Asn Arg Glu
1 5 10 15

Gln Trp Ala Glu Leu Arg Lys Ser Val Pro Leu Lys Leu Thr Glu Gln 20 25 30

Asp Leu Lys Pro Leu Leu Gly Phe Asn Glu Asp Leu Ser Leu Asp Glu 35 40 45

Val Ser Thr Ile Tyr Leu Pro Leu Thr Arg Leu Ile Asn Tyr Tyr Ile 50 55 60

Asp Glu Asn Leu His Arg Gln Thr Val Leu His Arg Phe Leu Gly Arg 65 70 75 80

Asn Asn Ala Lys Thr Pro Tyr Ile Ile Ser Ile Ala Gly Ser Val Ala 85 90 95

Val Gly Lys Ser Thr Ser Ala Arg Ile Leu Gln Ser Leu Leu Ser His 100 105 110

Trp Pro Thr Glu Arg Lys Val Asp Leu Ile Thr Thr Asp Gly Phe Leu 115 120 125 Tyr Pro Leu Asn Lys Leu Lys Gln Asp Asn Leu Leu Gln Lys Lys Gly 130 135 140

Phe Pro Val Ser Tyr Asp Thr Pro Lys Leu Ile Arg Phe Leu Ala Asp 145 150 155 160

Val Lys Ser Gly Lys Ser Asn Val Thr Ala Pro Ile Tyr Ser His Leu 165 170 175

Thr Tyr Asp Ile Ile Pro Asp Lys Phe Asp Val Val Asp Lys Pro Asp 180 185 190

Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Thr Gly Asn Asn Lys 195 200 205

Thr Asp Gln Thr Phe Val Ser Asp Phe Val Asp Phe Ser Ile Tyr Val 210 215 220

Asp Ala Glu Glu Lys Leu Leu Lys Glu Trp Tyr Ile Lys Arg Phe Leu 225 230 235 240

Lys Phe Arg Glu Ser Ala Phe Asn Asp Pro Asn Ser Tyr Phe Lys His 245 250 255

Tyr Ala Ser Leu Ser Lys Glu Glu Ala Ile Ala Thr Ala Ser Lys Ile 260 265 270

Trp Asp Glu Ile Asn Gly Leu Asn Leu Asn Gln Asn Ile Leu Pro Thr 275 280 285

Arg Glu Arg Ala Asn Leu Ile Leu Lys Lys Gly His Asn His Gln Val 290 295 300

Glu Leu Ile Lys Leu Arg Lys 305 310

<210> 2

<211> 316

<212> PRT

<213> Escherichia coli

<400> 2

Met Ser Ile Lys Glu Gln Thr Leu Met Thr Pro Tyr Leu Gln Phe Asp

Arg Asn Gln Trp Ala Ala Leu Arg Asp Ser Val Pro Met Thr Leu Ser 20 25 30

Glu Asp Glu Ile Ala Arg Leu Lys Gly Ile Asn Glu Asp Leu Ser Leu 35 40 45

Glu Glu Val Ala Glu Ile Tyr Leu Pro Leu Ser Arg Leu Leu Asn Phe 50 60

Tyr Ile Ser Ser Asn Leu Arg Arg Gln Ala Val Leu Glu Gln Phe Leu 65 70 75 80

WO 01/21772 PCT/US00/25993

- 3 -

Gly Thr Asn Gly Gln Arg Ile Pro Tyr Ile Ile Ser Ile Ala Gly Ser 85 90 95

Val Ala Val Gly Lys Ser Thr Thr Ala Arg Val Leu Gln Ala Leu Leu 100 105 110

Ser Arg Trp Pro Glu His Arg Arg Val Glu Leu Ile Thr Thr Asp Gly 115 120 125

Phe Leu His Pro Asn Gln Val Leu Lys Glu Arg Gly Leu Met Lys Lys 130 135 140

Lys Gly Phe Pro Glu Ser Tyr Asp Met His Arg Leu Val Lys Phe Val 145 150 155 160

Ser Asp Leu Lys Ser Gly Val Pro Asn Val Thr Ala Pro Val Tyr Ser 165 170 175

His Leu Ile Tyr Asp Val Ile Pro Asp Gly Asp Lys Thr Val Val Gln 180 185 190

Pro Asp Ile Leu Ile Leu Glu Gly Leu Asn Val Leu'Gln Ser Gly Met 195 200 205

Asp Tyr Pro His Asp Pro His His Val Phe Val Ser Asp Phe Val Asp 210 215 220

Phe Ser Ile Tyr Val Asp Ala Pro Glu Asp Leu Leu Gln Thr Trp Tyr 225 230 235

Ile Asn Arg Phe Leu Lys Phe Arg Glu Gly Ala Phe Thr Asp Pro Asp 245 250 255

Ser Tyr Phe His Asn Tyr Ala Lys Leu Thr Lys Glu Glu Ala Ile Lys

Thr Ala Met Thr Leu Trp Lys Glu Ile Asn Trp Leu Asn Leu Lys Gln 275 280 285

Asn Ile Leu Pro Thr Arg Glu Arg Ala Ser Leu Ile Leu Thr Lys Ser 290 295 300

Ala Asn His Ala Val Glu Glu Val Arg Leu Arg Lys 305 310 315

<210> 3

<211> 319

<212> PRT

<213> Bacillus subtilis

<400> 3

Met Lys Asn Lys Glu Leu Asn Leu His Thr Leu Tyr Thr Gln His Asn 1 5 10 15

Arg Glu Ser Trp Ser Gly Phe Gly Gly His Leu Ser Ile Ala Val Ser 20 25 30 Glu Glu Glu Ala Lys Ala Val Glu Gly Leu Asn Asp Tyr Leu Ser Val 35 40 45

Glu Glu Val Glu Thr Ile Tyr Ile Pro Leu Val Arg Leu Leu His Leu
50 55 60

His Val Lys Ser Ala Ala Glu Arg Asn Lys His Val Asn Val Phe Leu
65 70 75 80

Lys His Pro His Ser Ala Lys Ile Pro Phe Ile Ile Gly Ile Ala Gly 85 90 95

Ser Val Ala Val Gly Lys Ser Thr Thr Ala Arg Ile Leu Gln Lys Leu 100 105 110

Leu Ser Arg Leu Pro Asp Arg Pro Lys Val Ser Leu Ile Thr Thr Asp 115 120 125

Gly Phe Leu Phe Pro Thr Ala Glu Leu Lys Lys Lys Asn Met Met Ser 130 135 140

Arg Lys Gly Phe Pro Glu Ser Tyr Asp Val Lys Ala Leu Leu Glu Phe 145 150 155 160

Leu Asn Asp Leu Lys Ser Gly Lys Asp Ser Val Lys Ala Pro Val Tyr 165 170 175

Ser His Leu Thr Tyr Asp Arg Glu Glu Gly Val Phe Glu Val Val Glu 180 185 190

Gln Ala Asp Ile Val Ile Ile Glu Gly Ile Asn Val Leu Gln Ser Pro 195 200 205

Thr Leu Glu Asp Asp Arg Glu Asn Pro Arg Ile Phe Val Ser Asp Phe 210 220

Phe Asp Phe Ser Ile Tyr Val Asp Ala Glu Glu Ser Arg Ile Phe Thr 225 230 235 240

Trp Tyr Leu Glu Arg Phe Arg Leu Leu Arg Glu Thr Ala Phe Gln Asn 245 250 255

Pro Asp Ser Tyr Phe His Lys Phe Lys Asp Leu Ser Asp Gln Glu Ala 260 265 270

Asp Glu Met Ala Ala Ser Ile Trp Glu Ser Val Asn Arg Pro Asn Leu 275 280 285

Tyr Glu Asn Ile Leu Pro Thr Lys Phe Arg Ser Asp Leu Ile Leu Arg 290 295 300

Lys Gly Asp Gly His Lys Val Glu Glu Val Leu Val Arg Arg Val 305 310 315

<210> 4

0121772421

<211> 312

<212> PRT

<213> Mycobacterium leprae

<400> 4

Met Pro Arg Leu Ser Glu Pro Ser Pro Tyr Val Glu Phe Asp Arg Lys
1 5 10 15

Gln Trp Arg Ala Leu Arg Met Ser Thr Pro Leu Ala Leu Thr Glu Glu 20 25 30

Glu Leu Ile Gly Leu Arg Gly Leu Gly Glu Gln Ile Asp Leu Leu Glu
35 40 45

Val Glu Glu Val Tyr Leu Pro Leu Ala Arg Leu Ile His Leu Gln Val
50 55 60

Ala Ala Arg Gln Arg Leu Phe Ala Ala Thr Ala Glu Phe Leu Gly Glu 65 70 75 80

Pro Gln Gln Asn Pro Gly Arg Pro Val Pro Phe Ile Ile Gly Val Ala 85 90 95

Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala Arg Val Leu Gln Ala 100 105 110

Leu Leu Ala Arg Trp Asp His His Thr Arg Val Asp Leu Val Thr Thr 115 120 125

Asp Gly Phe Leu Tyr Pro Asn Ala Glu Leu Gly Arg Arg Asn Leu Met 130 135 140

His Arg Lys Gly Phe Pro Glu Ser Tyr Asn Arg Arg Ala Leu Met Arg 145 150 155 160

Phe Val Thr Ser Val Lys Ser Gly Ala Asp Tyr Ala Cys Ala Pro Val 165 170 175

Tyr Ser His Leu Arg Tyr Asp Thr Ile Pro Gly Ala Lys His Val Val 180 185 190

Arg His Pro Asp Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Thr 195 200 205

Gly Pro Thr Leu Met Val Ser Asp Leu Phe Asp Phe Ser Leu Tyr Val 210 . 215 220

Asp Ala Arg Ile Gln Asp Ile Glu Gln Trp Tyr Val Ser Arg Phe Leu 225 230 235 240

Ala Met Arg Gly Thr Ala Phe Ala Asp Pro Glu Ser His Phe His His 245 250 255

Tyr Ser Ala Leu Thr Asp Ser Lys Ala Ile Ile Ala Ala Arg Glu Ile 260 265 270

Trp Arg Ser Ile Asn Arg Pro Asn Leu Val Glu Asn Ile Leu Pro Thr 275 . 280 285

Arg Pro Arg Ala Thr Leu Val Leu Arg Lys Asp Ala Asp His Ser Ile

		205	300	•
290	•	295	300	•
Asn Arg Leu F 305	Arg Leu Arg 310	Lys Leu		
<210> 5 <211> 312 <212> PRT <213> Mycobae	cterium tub	erculosis		
<400> 5 Met Ser Arg :	Leu Ser Glu 5	Pro Ser P	Pro Tyr Val Glu	Phe Asp Arg Arg 15
Gln Trp Arg	Ala Leu Arç 20	Met Ser 1	hr Pro Leu Ala : 25	Leu Thr Glu Glu 30
Glu Leu Val	Gly Leu Arc	Gly Leu G	Gly Glu Gln Ile	Asp Leu Leu Glu 45
Val Glu Glu 50	Val Tyr Le	Pro Leu A	Ala Arg Leu Ile 60	His Leu Gln Val
Ala Ala Arg 65	Gln Arg Le		Ala Thr Ala Glu 75	Phe Leu Gly Glu 80
Pro Gln Gln	Asn Pro As	o Arg Pro	Val Pro Phe Ile 90	Ile Gly Val Ala 95
Gly Ser Val	Ala Val Gl 100		Thr Thr Ala Arg 105	Val Leu Gln Ala 110
Leu Leu Ala 115	Arg Trp As	p His His . 120	Pro Arg Val Asp	Leu Val Thr Thr 125
Asp Gly Phe 130	Leu Tyr Pr	o Asn Ala 135	Glu Leu Gln Arg 140	Arg Asn Leu Met
His Arg Lys 145	Gly Phe Pr		Tyr Asn Arg Arg 155	Ala Leu Met Arg 160
Phe Val Thr	Ser Val Ly 165	s Ser Gly	Ser Asp Tyr Ala 170	Cys Ala Pro Val 175
Tyr Ser His	Leu His Ty 180	r Asp Ile	Ile Pro Gly Ala 185	Glu Gln Val Val 190
Arg His Pro 195		u Ile Leu 200	Glu Gly Leu Asn	Val Leu-Gln Thr 205
Gly Pro Thr 210	Leu Met Va	l Ser Asp 215	Leu Phe Asp Phe 220	Ser Leu Tyr Val
Asp Ala Arg 225	Ile Glu As		-Gln Trp Tyr Val 235	Ser Arg Phe Leu 240

Ala Met Arg Thr Thr Ala Phe Ala Asp Pro Glu Ser His Phe His His

- 7 **-**

245 250 255 Tyr Ala Ala Phe Ser Asp Ser Gln Ala Val Val Ala Ala Arg Glu Ile Trp Arg Thr Ile Asn Arg Pro Asn Leu Val Glu Asn Ile Leu Pro Thr 280 Arg Pro Arg Ala Thr Leu Val Leu Arg Lys Asp Ala Asp His Ser Ile 295 Asn Arg Leu Arg Leu Arg Lys Leu <210> 6 <211> 329 <212> PRT <213> Streptomyces coelicolor Met Ile Ser Pro Val Pro Ser Ile Pro Arg Ser Ala'His Arg Gln Arg Pro Glu Ala Thr Pro Tyr Val Asp Leu Thr Arg Pro Glu Trp Ser Ala Leu Arg Asp Lys Thr Pro Leu Pro Leu Thr Ala Glu Glu Val Glu Lys Leu Arg Gly Leu Gly Asp Val Ile Asp Leu Asp Glu Val Arg Asp Ile Tyr Leu Pro Leu Ser Arg Leu Leu Asn Leu Tyr Val Gly Ala Thr Asp Gly Leu Arg Gly Ala Leu Asn Thr Phe Leu Gly Glu Gln Gly Ser Gln Ser Gly Thr Pro Phe Val Ile Gly Val Ala Gly Ser Val Ala Val Gly Lys Ser Thr Val Ala Arg Leu Leu Gln Ala Leu Leu Ser Arg Trp Pro 120 Glu His Pro Arg Val Glu Leu Val Thr Thr Asp Gly Phe Leu Leu Pro Thr Arg Glu Leu Glu Ala Arg Gly Leu Met Ser Arg Lys Gly Phe Pro Glu Ser Tyr Asp Arg Arg Ala Leu Thr Arg Phe Val Ala Asp Ile Lys Ala Gly Lys Ala Glu Val Thr Ala Pro Val Tyr Ser His Leu Ile Tyr

Asp Ile Val Pro Asp Gln Arg Leu Val Val Arg Arg Pro Asp Ile Leu

									-						1
		195			1		200					205			
Ile	Val 210	Glu	Gly	Leu	Asn	Val 215	Leu	Gln	Pro	Ala	Leu 220	Pro -	Gly	Lys	Asp
Gly 225	Arg	Thr	Arg	Val	Gly 230	Leu	Ala	Asp	Tyr	Phe 235	Asp	Phe	Ser	Val	Tyr 240
Val	Asp	Ala	Arg	Thr 245	Glu	Asp	Ile	Glu	Arg 250	Trp	Tyr	Leu	Asn ·	Arg 255	Phe
Arg	Lys	Leu	Arg 260	Ala	Thr	Ala	Phe	Gln 265	Asn	Pro	Ser	Ser	Tyr 270	Phe	Arg
Lys	Tyr	Thr 275	Gln	Val	Ser	Glu	Glu 280	Glu	Ala	Leu	Asp	Tyr 285	Ala	Arg	Thr
Thr	Trp 290	Arg	Thr	Ile	Asn	Lys 295		Asn	Leu	Val	Glu 300		Val	Ala	Pro
Thr 305	Arg	Gly	Arg	Ala	Thr 310	Leu	Val	Leu	Arg	L ys 315		Pro	Asp	His	Lys 320
Val	Gln	Arg	Leu	Ser 325	Leu	Arg	Lys	Leu							
<21 <21 <21		65 RT trep	tomy	ces	coel	icol	or								
			Thr	Ile S		Val	. Gly	/ Asn	Thr 10		s Thr	· Val	L L eu	'Gly 15	Leu
Phe	. Asp	Gly	/ Glu 20		Ile	· Val	. Glu	His 25		Ar	g İle	e Se	r Thr		Ser
Arg	, Arg	Th:		a Asp	·Glu	Lev	ı Ala 4(Let	ı Le	u Gli	n G1:		. Met	: Gly
Met	His 50		Leu	Let	ı Gly	Asr 55		ı Lev	ı Gly	y As	p Gly 6		e Asp	Gly	/ Ile
Ala 65		е Су:	s Ala	a Thi	r Val		Sez	r Val	l Le	u Hi 7		u Le	u Arç	g Gl	val 80
Thi	Arg	Ar	д Туі	с Ту: 8!		/ As	o Vai	l Pro	9 Ala		l Le	u Va	l -Glu	1 Pro 9!	o :Gly 5
Va.	L Lys	Th	r Gl		l Pro	Ile	e Le	10:		p Hi	s Pr	o Ły	s Glu 111		l Gly
Ala	a Asp	Ar 11		e Il	e Asr	n Ala	a Va.		a Al	a Va	1 G1	u Le 12		r Gl	y Gly

Pro Ala Ile Val Val Asp Phe Gly Thr Ala Thr Thr Phe Asp Ala Val

-9-

140 135 130 Ser Ala Arg Gly Glu Tyr Ile Gly Gly Val Ile Ala Pro Gly Ile Glu Ile Ser Val Glu Ala Leu Gly Val Lys Gly Ala Gln Leu Arg Lys Ile Glu Val Ala Arg Pro Arg Ser Val Ile Gly Lys Asn Thr Val Glu Ala Met Gln Ser Gly Ile Val Tyr Gly Phe Ala Gly Gln Val Asp Gly Val Val Asn Arg Met Ala Arg Glu Leu Ala Asp Asp Pro Asp Asp Val Thr Val Ile Ala Thr Gly Gly Leu Ala Pro Met Val Leu Gly Glu Ser Ser Val Ile Asp Glu His Glu Pro Trp Leu Thr Leu Met Gly Leu Arg Leu 250 Val Tyr Glu Arg Asn Val Ser Arg Met 260 <210> 8 <211> 272 <212> PRT <213> Mycobacterium tuberculosis <400> 8 Met Leu Leu Ala Ile Asp Val Arg Asn Thr His Thr Val Val Gly Leu Leu Ser Gly Met Lys Glu His Ala Lys Val Val Gln Gln Trp Arg Ile Arg Thr Glu Ser Glu Val Thr Ala Asp Glu Leu Ala Leu Thr Ile Asp Gly Leu Ile Gly Glu Asp Ser Glu Arg Leu Thr Gly Thr Ala Ala Leu Ser Thr Val Pro Ser Val Leu His Glu Val Arg Ile Met Leu Asp Gln Tyr Trp Pro Ser Val Pro His Val Leu Ile Glu Pro Gly Val Arg Thr Gly Ile Pro Leu Leu Val Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val Asn Cys Leu Ala Ala Tyr Asp Arg Phe Arg Lys Ala Ala Ile

Val Val Asp Phe Gly Ser Ser Ile Cys Val Asp Val Val Ser Ala Lys

	130				1	135					140	•			
Gly 145	Glu	Phe	Leu	Gly	Gly 150	Ala	Ile	Ala	₽ro	Gly 155	Val	≅Gln	Val	Ser	Ser 160
Asp	Ala	Ala	Ala	Ala 165	Arg	Ser	Ala	Ala	Leu 170	Arg	Arg	Val	-Glu	Leu 175	Ala
Arg	Pro	Arg	Ser 180	Val	Val	СĴЪ	Lys	Asn 185	Thr	Val	Glu	Cys	Met 1.90	:Gln	Ala
Gly	Ala	Val 195	Phe	СĴУ	Phe	Ala	Gly 200	Leu	Val	Asp	Gly	Leu 205	Val	Gly	Arg
Ile	Arg 210	Glu	Asp	Val	Ser	Gly 215	Phe	Ser	Val	Asp	His 220	Asp	Val	Ala	Ile
Val 225	Ala	Thr	Gly	His	Thr 230	Ala	Pro	Leu	Leu	Leu 235	Pro	G⊉u	Leu	His	Thr 240
Val	Asp	His	Tyr	Asp 245	Gln	His	Leu	Thr	Leu 250		Gly	Leu	Arg	Leu 255	
Phe	Glu	Arg	Asn 260	Leu	Glu	Val	Gln	Arg 265		Arg	Leu	Lys	Thr 270	Ala	Arq
					4										

Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 135 140

Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala 145 150 155 160

Leu Tyr Ser Arg Ala Ala Lys Leu Pro Arg Ile Glu Ile Thr Arg Pro
165 170 175

Asp Asn Ile Ile Gly Lys Asn Thr Val Ser Ala Met Gln Ser Gly Ile 180 185 190

Leu Phe Gly Tyr Val Gly Gln Val Glu Gly Ile Val Lys Arg Met Lys 195 200 205

Trp Gln Ala Lys Gln Asp Leu Lys Val Ile Ala Thr Gly Gly Leu Ala 210 215 220

Pro Leu Ile Ala Asn Glu Ser Asp Cys Ile Asp Ile Val Asp Pro Phe 225 230 235 240

Leu Thr Leu Lys Gly Leu Glu Leu Ile Tyr Glu Arg Asn Arg Val Gly · 245 250 255

Ser Val

<210> 10

<211> 262

<212> PRT

<213> Deinococcus radiopugnans

<400> 10

Met Pro Ala Phe Pro Leu Leu Ala Val Asp Ile Gly Asn Thr Thr 1 5 10 15

Val Leu Gly Leu Ala Asp Ala Ser Gly Ala Leu Thr His Thr Trp Arg
20 25 30

Ile Arg Thr Asn Arg Glu Met Leu Pro Asp Asp Leu Ala Leu Gln Leu 35 40 45

His Gly Leu Phe Thr Leu Ala Gly Ala Pro Ile Pro Arg Ala Ala Val 50 55 60

Leu Ser Ser Val Ala Pro Pro Val Gly Glu Asn Tyr Ala Leu Ala Leu 65 70 75 80

Lys Arg His Phe Met Ile Asp Ala Phe Ala Val Ser Ala Glu Asn Leu 85 90 95

Pro Asp Val Thr Val Glu Leu Asp Thr Pro Gly Ser Val Gly Ala Asp

Arg Leu Cys Asn Leu Phe Gly Ala Glu Lys Tyr Leu Gly Gly Leu Asp

		115					120				•	125			
Tyr	Ala 130	Val	Val	Val		Phe 135	-Gjy	Thr	Ser	Thr	Asn 140	Phe	Asp	Val	Val
Gly 145	Arg	Gly	Arg	Arg	Phe 150	Leu	G ly	∘Gly	Ile	Leu 155	Ala	Thr	Gly ,	Ala	Gln 160
Val	Ser	Ala	Asp	Ala 165	Leu	Phe	Ala	Arg	Ala 170	Ala	Lys	Leu	₽εο	Arg 175	Ile
Thr	Leu	Gln	Ala 180	Pro	Glu	Thr	Ala	Ile 185	Gly	Lys	Asn	Thr	Val 190	His	Ala
Leu	Gln	Ser 195	Gly	Leu	Val	₽he	Gly 200	Tyr	Ala	Glu	Met	Val 205	Asp	-Gly	Leu
Leu	Arg 210		Ile	Arg	Ala	Glu 215		Pro	Gly	Glu	Ala 220	Val	Ala	Val	Ala
Thr 225	Gly	Gly	Phe	Ser	Arg 230	Thr	Val	Gln	:Gly	Ile 235	•Суs	Ġln	-Glu	Ile	Asp 240
Tyr	Tyr	Asp	Glu	Thr 245	Leu	Thr	Leu	Arg	.Gly 250		Val	-Glu	Leu	Тер 255	Ala
Ser	Arg	Ser	Glu 260		Arg	•					•	•			
<21 <21	0> 1 1> 2 2> E 3> D	12 RT	.fovi	brio	vul	gari	.s ˙								
	0> 1 Thr		n His	: Phe	Leu	Lev	ı ₽h∈	Asp			Asn	Thr	Asn		Lys
1				5					10					15	
Ile	Gly	/ Ile	Ala 20		Glu	Thi	: Ala	Val 25		Thr	Ser	Tyr	30		Pro
Thr	: Asp) Pro		/ Glr	Thr	Thi	Asg 4(Ile	Gl)	/ Leu	Arc 45		Lec	ı 'Glu
Va]	L Let		g His	s Ala	Gly	Let 5		y Pro	Ala	Asg	Val 60		/ Ala	Cys	. Val
Ala 69	_	r Se	r Val	L Val	Pro 70		y Val	l Asr	Pro	Let 79		Arq	g Arg	Ala	Cys 80
Gl	ı Ar	у Ту	r Le	Tyr 85		Ly	s Le	ı Lev	2he		a Pro	Gly	y Asp) Ile 99	e Ala
116	e Pro	o Le	u Ası		n Arg	Ту	r Gl	u Arq		o Ala	a Glu	ı Va	l Gly 110		a Asp

Arg Leu Val Ala Ala Tyr Ala Ala Arg Arg Leu Tyr Pro Gly Pro Arg

115 120 125 Ser Leu Val Ser Val Asp Phe Gly Thr Ala Thr Thr Phe Asp Cys Val Glu Gly Gly Ala Tyr Leu Gly Gly Leu Ile Cys Pro Gly Val Leu Ser 155 Ser Ala Gly Ala Leu Ser Ser Arg Thr Ala Lys Leu Pro Arg Ile Ser Leu Glu Val Glu Glu Asp Ser Pro Val Ile Gly Arg Ser Thr Thr Ser Leu Asn His Gly Phe Ile Phe Gly Phe Ala Ala Met Thr Glu Gly Val Leu Ala Ala 210 <210> 12 <211> 246 <212> PRT <213> Thermotoga maritima <400> 12 Met Tyr Leu Leu Val Asp Val Gly Asn Thr His Ser Val Phe Ser Ile 10 Thr Glu Asp Gly Lys Thr Phe Arg Arg Trp Arg Leu Ser Thr Gly Val Phe Gln Thr Glu Asp Glu Leu Phe Ser His Leu His Pro Leu Leu Gly Asp Ala Met Arg Glu Ile Lys Gly Ile Gly Val Ala Ser Val Val Pro 50 Thr Gln Asn Thr Val Ile Glu Arg Phe Ser Gln Lys Tyr Phe His Ile Ser Pro Ile Trp Val Lys Ala Lys Asn Gly Cys Val Lys Trp Asn Val 85 90 Lys Asn Pro Ser Glu Val Gly Ala Asp Arg Val Ala Asn Val Val Ala Phe Val Lys Glu Tyr Gly Lys Asn Gly Ile Ile Asp Met Gly Thr 115 120 125 Ala Thr Thr Val Asp Leu Val Val Asn Gly Ser Tyr Glu Gly Gly Ala 135 Ile Leu Pro Gly Phe Phe Met Met Val His Ser Leu Phe Arg Gly Thr 145 155 160

Ala Lys Leu Pro Leu Val Glu Val Lys Pro Ala Asp Phe Val Val Gly

									•						
				165					170		•			175	
Lys	Asp	Thr	Glu 180	Glu	Asn	Ile	Arg	Leu 185	Gly	Val	Val	Asn	-Gly 190	Ser	Val
Tyr	Ala	Leu 195	Glu	Gly	Ile	Ile	Gly 200	Arg	Ile	Lys	-Glu	Val 205	Tyr	- Gly	Asp
Leu	Pro 210	Val	Val	Leu	Thr	Gly 215	Gly	Gl:n	Ser	Lys	11e 220	Val	Lys	Asp	Met
Ile 225	Lys	His	Glu	Ile	Phe 230	Asp	Glu	Asp	Leu	Thr 235	Ile	Lys	Gly	Val	Tyr 240
His	Phe	Суѕ	Phe	Gly 245	Asp										
<21 <21	0> 1 1> 2 2> P 3> T	73 RT	onema	pal	lidu	m									
<40 Met		3 Leu	ı Ile	Asp		Gly	Asn	Ser	His		. Val	. Phe	e Gly	Ile 15	Gln
Gly	Glu	ı Ası	n Gly 20		Arg	Val	. Cys	Va]		g Glu	ı Lev	Phe	Arg 30		Ala
Pro	Asp	Ala 3		, Lys	Thr	Glr	Asp 40		а Туз	: Sea	Leu	1 Let 45	ı Ile 5	His	Ala
Let	ı Cys 50		ı Arç	g Ala	Gly	Va :		/ Arc	g Ala	a 'Se	r Lei 60		g Asp	Ala	₽h∈
I16	_	r Se	r Val	l Val	Pro 70		l Lev	ı Th	r Ly:	s Th:		e Ala	a Asp	Ala	Val 80
Ala	a Gl	n Il	e Se	r Gly 85		. Gl	n Pro	o Va	1 Va.		e Gl	y Pr	o Trį	99 99	ту: Б
G1	u Hi	s Le	u Pr 10		l Arç	j Il	e Pro	o €1 10		o Va	l Ar	g Al	a Glu 111	ı Ile	e Gl
Th	r As	р Le 11	_	l Ala	a Asr	n Al	a Va 12		a Al	а Ту	r Va	1 Hi 12	s Pho 5	e Aro	g Se
Al	a Cy 13		l Va	l Va	l Ası	Cy 13		y Th	r Al	a Le	u Th 14		e Th	r Ala	a Va
As 14	_	y Th	r Gl	y Le	u Ile 15		n Gl	y Va	l Al	a Il 15		a Pr	o Gl	y Łe	u Ar
Th	r Al	a Va	1 G1	n Se 16		u Hi	s Th	r Gl	y Th		a Gl	n Le	u Pr	o Le 17	

Pro Leu Ala Leu Pro Asp Ser Val Leu Gly Lys Asp Thr Thr His Ala

- 15 -180 190 185 Val Gln Ala Gly Val Val Arg Gly Thr Leu Phe Val Ile Arg Ala Met 200 Ile Ala Gln Cys Gln Lys Glu Leu Gly Cys Arg Cys Ala Ala Val Ile Thr Gly Gly Leu Ser Arg Leu Phe Ser Ser Glu Val Asp Phe Pro Pro Ile Asp Ala Gln Leu Thr Leu Ser Gly Leu Ala His Ile Ala Arg Leu Val Pro Thr Ser Leu Leu Pro Pro Ala Thr Val Ser Gly Ser Ser Gly 265 Asn <210> 14 <211> 262 <212> PRT <213> Borrelia burgdorferi Met Asn Lys Pro Leu Leu Ser Glu Leu Ile Ile Asp Ile Gly Asn Thr Ser Ile Ala Phe Ala Leu Phe Lys Asp Asn Gln Val Asn Leu Phe Ile 20 25 Lys Met Lys Thr Asn Leu Met Leu Arg Tyr Asp Glu Val Tyr Ser Phe Phe Glu Glu Asn Phe Asp Phe Asn Val Asn Lys Val Phe Ile Ser Ser 55 Val Val Pro Ile Leu Asn Glu Thr Phe Lys Asn Val Ile Phe Ser Phe Phe Lys Ile Lys Pro Leu Phe Ile Gly Phe Asp Leu Asn Tyr Asp Leu Thr Phe Asn Pro Tyr Lys Ser Asp Lys Phe Leu Leu Gly Ser Asp Val

Thr Phe Asn Pro Tyr Lys Ser Asp Lys Phe Leu Leu Gly Ser Asp Val 100

Phe Ala Asn Leu Val Ala Ala Ile Glu Asn Tyr Ser Phe Glu Asn Val 115

Leu Val Val Asp Leu Gly Thr Ala Cys Thr Ile Phe Ala Val Ser Arg 130

Gln Asp Gly Ile Leu Gly Gly Ile Ile Asn Ser Gly Pro Leu Ile Asn 160

Phe Asn Ser Leu Leu Asp Asn Ala Tyr Leu Ile Lys Lys Phe Pro Ile

165	170	ı	175

Ser Thr Pro Asn Asn Leu Leu Glu Arg Thr Thr Ser Gly Ser Val Asn 180 185 190

Ser Gly Leu Phe Tyr Gln Tyr Lys Tyr Leu Ile Glu Gly Val Tyr Arg 195 200 205

Asp Ile Lys Gln Met Tyr Lys Lys Phe Asn Leu Ile Ile Thr Gly 210 215 220

Gly Asn Ala Asp Leu Ile Leu Ser Leu Ile Glu Ile Glu Phe Ile Phe 225 230 235 240

Asn Ile His Leu Thr Val Glu Gly Val Arg Ile Leu Gly Asn Ser Ile 245 250 255

Asp Phe Lys Phe Val Asn 260

<210> 15

<211> 229

<212> PRT

<213> Aquifex aeolicus

<400> 15

Met Arg Phe Leu Thr Val Asp Val Gly Asn Ser Ser Val Asp Ile Ala 1 5 10 15

Leu Trp Glu Gly Lys Lys Val Lys Asp Phe Leu Lys Leu Ser His Glu 20 25 30

Glu Phe Leu Lys Glu Glu Phe Pro Lys Leu Lys Ala Leu Gly I $\frac{1}{2}$ e Ser .

Val Lys Gln Ser Phe Ser Glu Lys Val Arg Gly Lys Ile Pro Lys Ile 50 55 60

Lys Phe Leu Lys Lys Glu Asn Phe Pro Ile Gln Val Asp Tyr Lys Thr 65 70 75 80

Pro Glu Thr Leu Gly Thr Asp Arg Val Ala Leu Ala Tyr Ser Ala Lys 85 90 95

Lys Phe Tyr Gly Lys Asn Val Val Val Ile Ser Ala Gly Thr Ala Leu 100° 105 110

Val Ile Asp Leu Val Leu Glu Gly Lys Phe Lys Gly Gly Phe Ile Thr 115 120 125

Leu Gly Leu Gly Lys Lys Leu Lys Ile Leu Ser Asp Leu Ala Glu Gly 130 135 140

Ile Pro Glu Phe Phe Pro Glu Glu Val Glu Ile Phe Leu Gly Arg Ser 145 150 155 160

Thr Arg Glu Cys Val Leu Gly Gly Ala Tyr Arg Glu Ser Thr Glu Phe

165

170

175

Ile Lys Ser Thr Leu Lys Leu Trp Arg Lys Val Phe Lys Arg Lys Phe 180 185 190

Lys Val Val Ile Thr Gly Gly Glu Gly Lys Tyr Phe Ser Lys Phe Gly 195 200 205

Ile Tyr Asp Pro Leu Leu Val His Arg Gly Met Arg Ash Leu Leu Tyr 210 215 220

Leu Tyr His Arg Ile 225

<210> 16

<211> 257

<212> PRT

<213> Synechocystis sp.

<400> 16

Met Glu Thr Ser Lys Pro Gly Cys Gly Leu Ala Leu Asp Asn Asp Lys
1 5 10 15

Gln Lys Pro Trp Leu Gly Leu Met Ile Gly Asn Ser Arg Leu His Trp 20 25 30

Ala Tyr Cys Ser Giy Asn Ala Pro Leu Gln Thr Trp Val Thr Asp Tyr 35 40 45

Asn Pro Lys Ser Ala Gln Leu Pro Val Leu Leu Gly Lys Val Pro Leu 50 55 60

Met Leu Ala Ser Val Val Pro Glu Gln Thr Glu Val Trp Arg Val Tyr 65 70 75 80

Gln Pro Lys Ile Leu Thr Leu Lys Asn Leu Pro Leu Val Asn Leu Tyr 85 90 95

Pro Ser Phe Gly Ile Asp Arg Ala Leu Ala Gly Leu Gly Thr Gly Leu 100 105 110

Thr Tyr Gly Phe Pro Cys Leu Val Val Asp Gly Gly Thr Ala Leu Thr 115 120 125

Ile Thr Gly Phe Asp Gln Asp Lys Lys Leu Val Gly Gly Ala Ile Leu 130 135 140

Pro Gly Leu Gly Leu Gln Leu Ala Thr Leu Gly Asp Arg Leu Ala Ala 145 150 155 160

Leu Pro Lys Leu Glu Met Asp Gln Leu Thr Glu Leu Pro Asp Arg Trp 165 170 175

Ala Leu Asp Thr Pro Ser Ala Ile Phe Ser Gly Val Val Tyr Gly Val
180 185 190

Leu Gly Ala Leu Gln Ser Tyr Leu Gln Asp Trp Gln Lys Leu Phe Pro

		. 05					200				,	205			
		195					200		_				_		-
Gly A	Ala 210	Ala	Met	Val	Iłe	Thr 215	Gly	Gly	Asp	∙Gly	Lys 220	He	Leu	His	grà
Phe 1 225	Leu	Lys	Glu	His	Ser 230	Pro	Asn	Leu	Ser	Val 235	Ala	Trp	Asp	Asp	Asn 240
Leu	Ile	Phe	Leu	Gly 245	Met	Ala	Ala	Ile	His 250	His	Gly	Asp	Arg	₽ro 255	Ile
Cys ·												•			
<210 <211 <212 <213	> 22 > PR	!3 !T	obact	er j	pyloi	ri						ı			
<400 Met 1			Arg	Gln 5	Ser	Phe	Thr	Asp	Leu 10		Asn	Leu	Val	Leu 15	Cys
Asp	Ile	Gly	Asn 20	Thr	Arg	Ile		₽he -25		•Gln	Asn	Tyr	Gln 30	Leu	Phe
Ser	Ser	Ala 35		Glu	Asp	Leu	Lys 40		Leu	Gly	Ile	Gln 45		Glu	Ile
Phe	Tyr 50	Ile	Ser	Val	Asn	Glu 55		Asn	Glu	Lys	Ala 60		Leu	Asn	'Cys
Tyr 65	Pro	Asn	Ala	Lys	Asn 70		Ala	ı-G1y	Phe	Phe 75		Łeu	-Glu	The	Asp 80
Tyr	Val	Gly	Leu	Gly 85		Asp	Arg	g Gl r	Met 90		Cys	Leu	Ala	Va]	Asn
Asn	Gly	Val	Val 100		Asp	Ala	ė ė̃l7	/ Ser 105		ı īle	e Thr	Ile	Asp 110		ı Ile
Lys	Glu	Gly 115		His	Leu	Gly	/ Gly		: Ile	e Let	ı Pro	Gly 125		Ala	a ∍Gln
Tyr	Ile 130		s Ala	Туг	Lys	Lys 139		r Ala	Ly:	s Ile	2 Leu 140		Gln	Pro	o Phe
Lys 145	Ala	Leu	ı Asp	Ser	Leu 150		ı Val	l Le	ı Pro	Ly:		Thi	Arg	j Asį	p Ala 160
Val	Asn	Туз	c Gly	Met 165		. Le	ı Se:	r Vai	l I16		a Cys	s Ile	e Glr	17:	s Leu 5
Ala	Lys	Ası	n Glr 180		; Ile	Ty	r Le	u Cy:		y -G 1	y As _t	o Ala	190		r Łeu

Ser Ala Phe Leu Pro His Ser Val Cys Lys Glu Arg Leu Val Phe Asp

- 19 -

195 200 205

Gly Met Glu Ile Ala Leu Lys Lys Ala Gly Ile Leu Glu Cys Lys 210 220

<210> 18
<211> 267

<211> 267 <212> PRT <213> Bordetella pertussis

Phe Asp Pro Asp Ala Pro Gln Ala Ala Arg Glu Pro Ala Pro Val Ala 20 25 30

Phe Asp Asn Leu Asp Leu Asp Ala Leu Gly Arg Trp Leu Ala Thr Leu 35 40 45

Pro Arg Arg Pro Gln Arg Ala Leu Gly Val Asn Val Ala Gly Leu Ala 50 55 60

Arg Gly Glu Ala Ile Ala Ala Thr Leu Arg Ala Gly Gly Cys Asp Ile 65 70 75 80

Arg Trp Leu Arg Ala Gln Pro Leu Ala Met Gly Leu Arg Asn Gly Tyr
85 90 95

Arg Asn Pro Asp Gln Leu Gly Ala Asp Arg Trp Ala Cys Met Val Gly
100 105 110

Val Leu Ala Arg Gln Pro Ser Val His Pro Pro Leu Leu Val Ala Ser 115 120 125

Phe Gly Thr Ala Thr Thr Leu Asp Thr Ile Gly Pro Asp Asn Val Phe 130 135 140

Pro Gly Gly Leu Ile Leu Pro Gly Pro Ala Met Met Arg Gly Ala Leu 145 150 155 160

Ala Tyr Gly Thr Ala His Leu Pro Leu Ala Asp Gly Leu Val Ala Asp 165 170 175

Tyr Pro Ile Asp Thr His Gln Ala Ile Ala Ser Gly Ile Ala Ala 180 185 190

Gln Ala Gly Ala Ile Val Arg Gln Trp Leu Ala Gly Arg Gln Arg Tyr 195 200 205

Gly Gln Ala Pro Glu Ile Tyr Val Ala Gly Gly Gly Trp Pro Glu Val 210 215 220

Arg Gln Glu Ala Glu Arg Leu Leu Ala Val Thr Gly Ala Ala Phe Gly 225 230 235 240

Ala Thr Pro Gln Pro Thr Tyr Leu Asp Ser Pro Val Leu Asp Gly Leu

245 , 250 255

Ala Ala Leu Ala Ala Gln Gly Ala Pro Thr Ala 260 265

<210> 19 <211> 777 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(774) <400> 19 ttg tta ctg gtt atc gat gtg ggg aac acc aat act gta ctt ggt gta 48 Leu Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val 10 tat cat gat gga aaa tta gaa tat cac tgg cgt ata gaa aca agc agg 96 Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg cat aaa aca gaa gat gag ttt ggg atg att ttg cgc tcc tta ttt gat His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp cac too ggg off atg tit gaa cag ata gat ggc att att att tog toa 192 His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser gta gtg ccg cca atc atg ttt gcg tta gaa aga atg tgc aca aaa tac 240 Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr 288 ttt cat atc gag cct caa att gtt ggt cca ggt atg aaa acc ggt tta Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu aat ata aaa tat gac aat ccg aaa gaa gta ggg gca gac aga atc gta 336 Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val 100 aat gct gtc gct gcg ata cac ttg tac ggc aat cca tta att gtt gtc 384 Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 120 115 432. gat ttc gga acc gcc aca acg tac tgc tat att gat gaa aac aaa caa Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 135 130 480 tac atg ggc ggg gcg att gcc cct ggg att aca att tcg aca gag gcg Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala 155 145 150 ctt tac tcg cgt gca gca aag ctt cct cgt atc gaa atc acc cgg ccc Leu Tyr Ser Arg Ala Ala Lys Leu Pro Arg Ile Glu Ile Thr Arg Pro

ennein -wn - 012177282 L s

- 21 -

				165					170					175		
gac Asp	aat Asn	att Ile	atc Ile 180	gga Gly	aaa Lys	aac Asn	act Thr	gtt Val 185	agc Ser	gcg Ala	åtg Met	caa Gln	tct Ser 190	gga Gly	att Ile	576
tta Leu	ttt Phe	ggc Gly 195	tat Tyr	gtc Val	ggc Gly	caa Gln	gtg Val 200	gaa Glu	gga Gly	atc Ile	gtt Val	aag Lys 205	cga Arg	atg Met	aaa Lys	624
tgg Trp	cag Gln 210	gca Ala	aaa Lys	cag Gln	gac Asp	ctc Leu 215	aag Lys	gtc Val	att Ile	gcg Ala	aca Thr 220	gga Gly	ggc Gly	ctg Leu	gcg Ala	672
ccg Pro 225	ctc Leu	att Ile	gcg Ala	aac Asn	gaa Glu 230	tca Ser	gat Asp	tgt Cys	ata Ile	gac Asp 235	atc Ile	gtt Val	gat Asp	cca Pro	ttc Phe 240	¹ 72(
tta Leu	acc Thr	cta Leu	aaa Lys	ggg Gly 245	ctg Leu	gaa Glu	ttg Leu	att Ile	tat Tyr 250	gaa Glu	aga Arg	aac Asn	cgc Arg	gta Val 255	gga Gly	768
-	gta Val	tag														777
<213 <213	0> 20 1> 90 2> Di 3> Ba	06 NA	lus :	subt:	ilis		ı									
	0> l> CI 2> (1		(957))												,
<400)> 20)														
gtg Met 1	aaa Lys	aat Asn	aaa Lys	gaa Glu 5	ctt Leu	aac Asn	cta Leu	cat His	act Thr 10	tta Leu	tat Tyr	aca Thr	cag Gln	cac His 15	aat Asn	48
cgg Arg	gag Glu	tct Ser	tgg Trp 20	tct Ser	ggt Gly	ttt Phe	ggg Gly	ggg Gly 25	cat His	ttg Leu	tcg Ser	att Ile	gct Ala 30	Val	tct Ser	96
gaa Glu	gaa Glu	gag Glu 35	gca Ala	aaa Lys	gct Ala	gtg Val	gaa Glu 40	gga Gly	ttg Leu	aat Asn	gat Asp	tat Tyr 45	cta Leu	tct Ser	gtt Val	144
gaa Glu	gaa Glu 50	gtg Val	gag Glu	acg Thr	atc Ile	tat Tyr 55	att Ile	ccg Pro	ctt Leu	gtt Val	cgc Arg 60	ttg Leu	ctt Leu	cat His	tta Leu	192
cat His 65	gtc Val	aag Lys	tct Ser	gcg Ala	gct Ala 70	gaa Glu	cgc Arg	aat Asn	aag Lys	cat His 75	gtc Val	aat Asn	gtt Val	ttt Phe	ttg Leu 80	240
aag Lys	cac His	cca Pro	cat His	tca Ser	gcc Ala	aaa Lys	att Ile	ccg Pro	ttt Phe	att Ile	atc Ile	ggc Gly	att Ile	gcc Ala	ggc Gly	288

DEICHOOLIN SEIN NESTTORG I S

				85					90					95		
agt Ser	gtc Val	gca Ala	gtc Val 100	gga Gly	aaa Lys	agċ Ser	acg Thr	acg Thr 105	gcg Ala	cgg Arg	atc Ile	ttg Leu	cag Gln 110	aag Lys	ctg Leu	336
ctt Leu	tcg Ser	cgt Arg 115	ttg Leu	cct Pro	gac Asp	cgt Arg	cca Pro 120	aaa Lys	gtg Val	agc Ser	ctt Leu	ato Ile 125	acg Thr	aca Thr	gat Asp	384
ggt Gly	ttt Phe 130	tta Leu	ttt Phe	cct Pro	act Thr	gcc Ala 135	gag Glu	ctg Leu	aaa Lys	aag Lys	aaa Lys 140	aat Asn	atg Met	atg Met	tca Ser	432
aga Arg 145	aaa Lys	gga Gly	ttt Phe	cct Pro	gaa Glu 150	agc Ser	tat Tyr	gat Asp	gta Val	aag Lys 155	gcg Ala	ctg Leu	-ct-c Leu	gaa Glu	ttt Phe 160	480
ttg Leu	aat Asn	gac Asp	tta Leu	aaa Lys 165	tca Ser	gga Gly	aag Lys	gac Asp	agc Ser 170	gta Val	aag Lys	gcc Ala	ecg Pro	gtg Val 175	tat Tyr	528
tcc Ser	cat His	cta Leu	acc Thr 180	tat	gac Asp	cgc Arg	gag Glu	gaa Glu 185	Gly	gtg Val	ttc Phe	gag •Glu	gtt Val 190	Val	gaa Glu	576
cag Gln	gcg Ala	gat Asp 195	Ile	gtg Val	att Ile	att Ile	gaa Glu 200	Gly	att	aat Asn	gtt Val	ctt Leu 205	Gln	teg	Pro	624
acc Thr	ttg Leu 210	Glu	gat Asp	gac Asp	cgg Arg	gaa Glu 215	Asn	ccg Pro	,-cgt Arg	att Ile	ttt Phe 220	. Val	⊹tcc Ser	gat Asp	ttc Phe	672
ttt Phe 225	Asp	ttt Phe	t.co	att Ile	tat Tyr 230	. Val	gat Asp	gcg Ala	g gag a Glu	gaa Glu 235	Ser	cgç Arç	g att g Ile	tto Phe	act Thr 240	720
tgg Trp	tat Tyr	tta Leu	a gaq ı Glu	g cgt u Arg 245	Phe	cgc Arg	ctg Lev	ctt Lei	cgg Arg 250	Glι	aca Thi	a gct c Ala	ttt a Phe	caa e Gl: 25	a aat n Asn 5	768
cct	gat Asp	t tca o Sei	tat Ty: 26	r Phe	cat His	t aaa s Lys	ttt Phe	aaa Lys 265	s Asp	tto Lev	g tick i Sei	c gat r Asi	-caq -Gl: -276	n Gl	g gct u Ala	814
gac Asp	gaq Gl	g ate u Me 27	t Al	a gco a Ala	tc Sei	g att r Ile	tg Tr _{ 280	Gl	g agt u Sei	gte Va	c aad l Asi	c cge n Are 28:	g Pro	g aa o As	t tta n Leu	86
tat Ty:	ga: Gl: 29	u As	t at n Il	t tto e Leo	g cca u Pro	a act o Thi 29!	Ly:	a tto s Pho	c ago	g tca g Sea	a ga: r As; 30	p Le	c at	t tt e L e	g cgt u Arg	91
aaq Ly:	s G1	a ga y As	c gg p Gl	g cat y Hi:	t aa s Ly 31	s Va	gae l Gl	g ga	a gto u Va.	g tt l Le 31	u Va	a ag l Ar	g ag g Ar	g gt g Va	a tga l	96

<212 <212	0> 2: L> 88 2> Di 3> Ba	82 NA	lus :	subt	ilis						•					
	0> L> CI 2> (:		(879))		-							,			,
)> 2:															
ttg Met 1	tcg Ser	att Ile	gct Ala	gta Val 5	tct Ser	gaa Glu	gaa Glu	gag Glu	gca Ala 10	aaa Lys	gct Ala	gtg Val	gaa Glu	gga Gly 15	ttg Leu	48
aat Asn	gat Asp	tat Tyr	cta Leu 20	tct Ser	gtt Val	gaa Glu	gaa Glu	gtg Val 25	gag Glu	acg Thr	atc Ile	tat Tyr	att Ile 30	ccg Pro	ctt Leu	96
gtt Val	cgc Arg	ttg Leu 35	ctt Leu	cat His	tta Leu	cat His	gtc Val 40	aag Lys	tct Ser	gcg Ala	gct Ala	gaa Glu ′45	cgc Arg	aat Asn	aag Lys	144
cat His	gtc Val 50	aat Asn	gtt Val	ttt Phe	ttg Leu	aag Lys 55	cac His	cca Pro	cat His	tca Ser	gcc Ala 60	aaa Lys	att Ile	ccg Pro	ttt Phe	192
att Ile 65	atc Ile	ggc Gly	att Ile	gcc Ala	ggc Gly 70	agt Ser	gtc Val	gca Ala	gtc Val	gga Gly 75	aaa Lys	a'gc Ser	acg Thr	acg Thr	gcg Ala 80	240
cgg Arg	atc Ile	ttg Leu	cag Gln	aag Lys 85	ctg Leu	ctt Leu	tcg Ser	cgt Arg	ttg Leu 90	cct Pro	gac Asp	cgt Arg	cca Pro	aaa Lys 95	gtg Val	288
agc Ser	ctt Leu	atc Ile	acg Thr 100	aca Thr	gat Asp	ggt Gly	ttt Phe	tta Leu 105	ttt Phe	cct Pro	act Thr	gcc Ala	gag Glu 110	ctg Leu	aaa Lys	, 336
aag Lys	aaa Lys	aat Asn 115	atg Met	atg Met	tca Ser	aga Arg	aaa Lys 120	gga Gly	ttt Phe	cct Pro	gaa Glu	agc Ser 125	tat Tyr	gat Asp	gta Val	384
aag Lys	gcg Ala 130	ctg Leu	ctc Leu	gaa Glu	ttt Phe	ttg Leu 135	aat Asn	gac Asp	tta Leu	aaa Lys	tca Ser 140	gga Gly	aag Lys	gac Asp	agc Ser	432
gta Val 145	aag Lys	gcc Ala	ccg Pro	gtg Val	tat Tyr 150	tcc Ser	cat His	cta Leu	acc Thr	tat Tyr 155	gac Asp	cgc Arg	gag Gl _. u	gaa Glu	ggt Gly 160	480
gtg Val	ttc Phe	gag Glu	gtt Val	gta Val 165	gaa Glu	cag Gln	gcg Ala	gat Asp	att Ile 170	gtg Val	att Ile	att Ile	gaa Glu	ggc Gly 175	att Ile	528
aat Asn	gtt Val	ctt Leu	cag Gln 180	tcg Ser	ccc Pro	acc Thr	ttg Leu	gag Glu 185	gat Asp	gac Asp	cgg Arg	gaa Glu	aac Asn	ccg Pro	cgt Arg	576

att Ile	ttt Phe	gtt Val 195	tcc Ser	gat Asp	ttc Phe	ttt Phe	gat Asp 200	ttt Phe	tcg Ser	att Ile	tat Tyr	gtg Val 205	gat . Asp	gcg Ala	gag Glu	624
gaa Glu	agc Ser 210	cgg Arg	att Ile	ttc Phe	act Thr	tgg Trp 215	tat Tyr	tta Leu	gag: Glu	cgt Arg	ttt Phe 220	cgc Arg∙	ctg Leu	ctt Leu	egg Arg	672
gaa Glu 225	aca Thr	gct Ala	ttt Phe	caa Gln	aat Asn 230	cct Pro	gat Asp	tca Ser	tat Tyr	ttt Phe 235	cat His	aaa Lys	ttt Phe	aaa Lys	gac Asp 240	720
ttg Leu	tcc Ser	gat Asp	cag Gln	gag Glu 245	gct Ala	gac Asp	gag Glu	Met	gca Ala 250	gcc Ala	tcg Ser	att Ile	tgg Trp	gag Glu 255	agt Ser	768
gtc Val	aac Asn	cgg Arg	ccg Pro 260	Asn	tta Leu	tat Tyr	gaa Glu	aat Asn 265	att Ile	ttg Leu	cca Pro	ac't Thr	aaa Lys 270	Phe	agg Arg	816
tca Ser	gat Asp	ctc Leu 275	att Ile	ttg Leu	cgt Arg	aag Lys	gga Gly 280	gac Asp	Gly	cat His	aag Łys	gtc Val 285	gag Glu	gaa Glu	gtg Val	864
	gta Val 290	Arg			tga											882 ,
<21 <21	.0> 2 .1> 8 .2> 0 .3> E	146 NA	lus	subt	ilis											
<21 <21 <21 <22	.1> 8 .2> C .3> E	MA NA Bacil			ilis											
<21 <21 <21 <22 <22 <22 <40	1> 8 12> 0 13> E 20> 21> 0 22> 0	346 DNA Bacil CDS (1)	(843 a tto	3) gaat u Asr	: qat	tat	cta Leu	ı tot ı Sei	: gtt : Val	L Gl	a gaa u Glu	a gto	g gad	g ac u Th l	g atc r Ile 5	48
<21 <21 <22 <22 <22 <40 gtq	1> 8 12> 0 13> E 20> 21> 0 22> 0 00> 2 g gaa t Glu	MAGENTAL COST (1)	(843) a tto / Let	g aat 1 Asr 5 t gtt	gat n Asp	tato Tyr	r Leu	Sei	Val	l Gli D a cai	u Gli t gt:	u Val c aad	i Gli	u Th 1 t gc r Al	r Ile	4 8
<21 <21 <21 <22 <22 <22 <22 <40 gtg	1> 8 2> 0 13> E 20> 21> 0 22> 0 00> 2 10	A46 DNA Bacil CDS (1) 22 a gga a Gly t ccc	(843) a tto y Leo g ctt cttaac t aac n Ly:	g aat 1 Asr 5 t gtt 1 Val	gat n Asp o c ego L Aro	tato Tyr tto Lei	t gti	cat His 2: ttt	t tte	l Glu D a can u His	t gtos Va	u Val	tc s Se 3	u Th 1 t gc r Al 0	r lle 5 g gct	
<21 <21 <21 <22 <22 <22 <22 <22 <40 gtc Met Ty:	1> 8 2> 0 13> E 20> 21> 0 22> 0 22> 0 11 1	t ccce Pro	(843) a tto y Let 20 t aao	g aat 1 Asr 2 gtt 1 Val 0 g cat 1 s Hi:	gat n Asp c cgc L Arc t gtc s Val	tato Tyr tto Lev Lev Asi	g ctt g ctt g ctt t gtt n Val 40 c att	cat His 2: ttt	t tte	e can du Hi: g aad u Ly	t gto s Va. g cao s Hi	c aac c ccc s Pr 4 c gc l Al	g tc s Se 3 a ca o Hi	t gc r Al 0 t tc s Se	r lle 5 g gct a Ala a gcc	96

65					70					75					80	
						atc Ile										288
gcc Ala	gag Glu	ctg Leu	aaa Lys 100	aag Lys	aaa Lys	aat Asn	atg Met	atg Met 105	tca Ser	aga Arg	aaa Lys	gga Gly	ttt Phe 110	cct Pro	gaa Glu	336
agc Ser	tat Tyr	gat Asp 115	gta Val	aag Lys	gcg Ala	ctg Leu	ctc Leu 120	gaa Glu	ttt Phe	ttg Leu	aat Asn	gac Asp 125	tta Leu	aaa Lys	tca Ser	384
gga Gly	aag Lys 130	gac Asp	agc Ser	gta Val	aag Lys	gcc Ala 135	ccg Pro	gtg Val	tat Tyr	tcc Ser	cat His 140	cta Leu	acc Thr	tat Tyr	gac Asp	432
cgc Arg 145	gag Glu	gaa Glu	ggt Gly	gtg Val	ttc Phe 150	gag Glu	gtt Val	gta Val	gaa Glu	cag Gln 155	gcg Ala	gat Asp	att Ile	gtg Val	att Ile 160	480
att Ile	gaa Glu	ggc Gly	att Ile	aat Asn 165	gtt Val	ctt Leu	cag Gln	tcg Ser	ccc Pro 170	acc Thr	ttg Leu	gag Glu	gat Asp	gac Asp 175	cgg Arg	528
gaa Glu	aac Asn	ccg Pro	cgt Arg 180	att Ile	ttt Phe	gtt Val	tcc Ser	gat Asp 185	ttc Phe	ttt Phe	gat Asp	ttt Phe	tcg Ser 190	att Ile	tat Tyr	576
gtg Val	gat Asp	gcg Ala 195	gag Glu	gaa Glu	agc Ser	cgg Arg	att Ile 200	ttc Phe	act Thr	tgg Trp	tat Tyr	tta Leu 205	gag Glu	cgt Arg	ttt Phe	624
cgc Arg	ctg Leu 210	ctt Leu	cgg Arg	gaa Glu	aca Thr	gct Ala 215	ttt Phe	caa Gln	aat Asn	cct Pro	gat Asp 220	tca Ser	tat Tyr	ttt Phe	cat His	672
aaa Lys 225	ttt Phe	aaa Lys	gac Asp	ttg Leu	tcc Ser 230	gat Asp	cag Gln	gag Glu	gct Ala	gac Asp 235	gag Glu	atg Met	gca Ala	gcc Ala	tcg Ser 240	720
att Ile	tgg Trp	gag Glu	agt Ser	gtc Val 245	aac Asn	cgg Arg	ccg Pro	aat Asn	tta Leu 250	tat Tyr	gaa Glu	aat Asn	att Ile	ttg Leu 255	cca Pro	768
act Thr	aaa Lys	ttc Phe	agg Arg 260	tca Ser	gat Asp	ctc Leu	att Ile	ttg Leu 265	cgt Arg	aag Lys	gga Gly	gac Asp	ggg Gly 270	cat His	aag Lys	816
						agg Arg			tga							846

<210> 23 <211> 831 <212> DNA

<213> Bacillus	s subtilis				
<220> <221> CDS <222> (1)(8	31)			,	•
<400> 23 atg aaa aca a Met Lys Thr L 1	aa ctg gat t ys Leu Asp l 5	ttt cta aaa Phe Leu Lys	atg aag gag t Met Lys Glu S 10	cct gaa gaa ccg Ser Glu Glu Pro 15	48
Ile Val Met L	etg acc gct (Seu Thr Ala (20	tat gat tat Tyr Asp Tyr 25	ccg gca gct a	aaa ctt gct gaa Lys Leu Ala Glu 30	96
caa gcg gga g Gln Ala Gly V 35	gtt gac atg Val Asp Met	att tta gtc Ile Leu Val 40	ggt gat tca d Gly Asp Ser	ctt gga atg gtc Leu Gly Met Val 45	144
gtc ctc ggc c Val Leu Gly L 50	ctt gat tca Leu Asp Ser	act gtc ggt Thr Val Gly 55	gtg aca gtt Val Thr Val 60	gcg gac atg atc Ala Asp Met Ile	192
cat cat aca a His His Thr I 65	aaa gcc gtt Lys Ala Val 70	Lys Arg Gly	gcg ccg aat Ala Pro Asn 75	acc ttt att gtg Thr Phe Ile Val 80	240
aca gat atg o	ccg ttt atg Pro Phe Met 85	tct tat cac Ser Tyr His	ctg tct aag Leu Ser Lys 90	gaa gat acg ctg Glu Asp Thr Leu 95	288
Lys Asn Ala A	gcg gct atc Ala Ala Ile 100	gtt cag gaa Val Gln Glu 105	Ser Gly Ala	gac gca ctg aag Asp Ala Leu Lys 110	336
ctt gag ggc (Leu Glu Gly (115	gga gaa ggc Gly Glu Gly	gtg ttt gaa Val Phe Glu 120	tcc att cgc Ser Ile Arg	gca ttg acg ctt Ala Leu Thr Leu 125	384
gga ggc att o Gly Gly Ile 1 130	cca gta gtc Pro Val Val	agt cac tta Ser His Leu 135	ggt ttg aca Gly Leu Thr 140	ecg cag tea gte Pro Gln Ser Val	432
ggc gta ctg (Gly Val Leu (145	ggc ggc tat Gly Gly Tyr 150	aaa gta cag Lys Val Gln	ggc aaa gac Gly Lys Asp 155	gaa caa agc gcc Glu Gln Ser Ala 160	
aaa aaa tta Lys Lys Leu	ata gaa gac Ile Glu Asp 165	agt ata aaa Ser Ile Lys	tgc gaa gaa Cys Glu Glu 170	gca gga gct atg Ala Gly Ala Met 175	528
Met Leu Val	ctg gaa tgt Leu Glu Cys 180	gtg ccg gca Val Pro Ala 185	Glu Leu Thr	gcc aaa att gcc Ala Lys Ile Ala 190	: 57 [.] 6
gag acg cta Glu Thr Leu 195	agc ata ccg Ser Ile Pro	gtc att gga Val Ile Gly 200	a atc ggg gct / Ile Gly Ala	ggt gtg aaa gcg Gly Val Lys Ala 205	j 624 1

01/2177262 I S

gac Asp	gga Gly 210	caa Gln	gtt Val	ctc Leu	gtt Val	tat Tyr 215	cat His	gat Asp	att Ile	atc Ile	ggc Gly 220	cac His	ggt Gly	gtt Val	gag Glu	672
aga Arg 225	aca Thr	cct Pro	aaa Lys	ttt Phe	gta Val 230	aag Lys	caa Gln	tat Tyr	acg Thr	cgc Arg 235	att Ile	gat Asp	gaa Glu	acc Thr	atc Ile 240	720
gaa Glu	aca Thr	gca Ala	atc Ile	agc Ser 245	gga Gly	tat Tyr	gtt Val	cag Gln	gat Asp 250	gta Val	aga Arg	cat His	cgt Arg	gct Ala 255	ttc Phe	768
cct Pro	gaa Glu	caa Gln	aag Lys 260	cat His	tcc Ser	ttt Phe	caa Gln	atg Met 265	aac Asn	cag Gln	aca Thr	gtg Val	ctt Leu 270	gac Asp	ggc Gly	816
	tac Tyr										÷					831
<213 <213	0> 24 1> 2 2> P1 3> Ba	7 7 RT	lus :	subt:	ilis					•		ı				
	0> 2											•				
Met 1	Lys	Thr	Lys	Leu 5	Asp	Phe	Leu	Lys	Met 10	Lys	Glu	Ser	Glu	Glu 15	Pro	
Ile	Val	Met	Leu 20	Thr	Ala	Tyr	Asp	Tyr 25	Pro	Ala	Ala	Lys	Leu 30	Ala	Glu	•
Gln	Ala	Gly 35	Val	Asp	Met	Ile	Leu 40	Val	Gly	Asp	Ser	Leu 45	Gly	Met	Val	
Val	Leu 50	Gly	Leu	Asp	Ser	Thr 55	Val	Gly	Val	Thr	Val 60	Ala	Asp	Met	Ile	
His 65	His	Thr	Lys	Ala	Val 70	Lys	Arg	Gly	Ala	Pro 75	Asn	Thr	Phe	Ile	Val 80	
Thr	Asp	Met	Pro	Phe 85	Met	Ser	Tyr	His	Leu 90		Lys	Glu	Asp	Thr 95	Leu	
Lys	Asn	Ala	Ala 100	Ala	Ile	Val.	Gln	Glu 105	Ser	Gly	Ala	Asp	Ala 110	Leu	Lys	
Leu	Glu	Gly 115	Gly	Glu	Gly	Val	Phe 120	Glu	Ser	Ile	Arg	Ala 125	Leu	Thr	Leu	
Gly	Gly 130	Ile	Pro	Val	Val	Ser 135	His	Leu	Gly	Leu	Thr 140	Pro	Gln	Ser	Val	
Gly 145	Val	Leu	Gly	Gly	Tyr 150	Lys	Val	Gln	Gly	Lys 155	Asp	Glu	Gln	Ser	Ala 160	

.......

Lys	Lys	Leu	Ile	Glu 165	Asp	Ser	Ile	L ys	Cys 170	Glu	Glu	Ala	Gly	Ala 175	Met	
Met	Leu	Val	Leu 180	Glu	Суѕ	Val	Pro	Ala 185	Glu	Leu	Thr	Ala	Lys 190	Ile	Ala	
Glu	Thr	Leu 195	Ser	Ile	Pro	Val	11e 200	Gly	Ile	Gly	Ala	Gly 205	Vaļ	Lys	Ala	
Asp	Gly 210	Gln	Val	Leu	Val	Tyr 215	His	Asp	Ile	Ile	Gly 220	His	€ly	Val	Glu	
Arg 225	Thr	Pro	Lys	Phe	Val 230	Lys	-Gln	Tyr	Thr	Arg 235	Ile	Asp	-Glu	Thr	11e 240	•
Glu	Thr	Ala	Ile	Ser 245	Gly	Tyr	Val	Gln	Asp 250		Arg	His	Arg	Ala 255	₽he	
Pro	Glu	Gln	Lys 260	His	Ser	Phe	Gln	Met 265		Gln	Thr	Val '	Leu 270	Asp	Gly	
Leu	Tyr	Gly 275		Lys												
<21 <21	0> 2 1> 8 2> D 3> B	58	lus	subt	ilis											
	1> 0	DS (1)	(858	1)			·									
ato Met	00> 2 g aga : Arq L	cad	g att	act Thr	Asp	att Ile	tca Ser	caç Glr	g cto Lei 10	Lys	a gaa s Glu	a gcc a Ala	ata Ile	a aaa E Lys 15	a caa s Gln	48
tae Ty:	c cat	t toa	a gaq Glu 20	ı Gly	aaq Lys	tca Ser	ato Ile	gg; G1; 2:	y Phe	gtt Val	CCC L Pro	g acq o Thi	ato Met	Gl;	g ttt y Phe	96
ct: Le	g cat u Hi:	t gad s Glu 3	ı Gl	g cat y His	tta Le	a acc	tta Lev 40	ı Al	a gad a Asi	c aaa o Lys	a gca s Ala	a aga a Arq 4	g Gl	a gaa n Gl	a aac u Asn	144
ga As	c gc p Ala 5	a Va	t ati	t atq e Mei	g agt	att r Ile 5	e Phe	gte Va	g aas 1 Ass	t cci	t gca o Ala 6	a Glı	a tto n Pho	c gg e Gl	c .cct y Pro	192
aa As 6	n Gl	a ga u As	t tt p Ph	t gaa e Gl	a gca u Ala 70	а Ту	t cc	g cg o Ar	c ga g As	t ate p Ile 7	e Gl	g cg u Ar	g ga g As	t gc p Al	a gct a Ala 80	1
ct Le	t gc u Al	a ga a .Gl	a aa u As	c gco n Ala	a Gl	a gte y Va	c ga l As _l	t at p Il	t ct e Le 9	u Ph	t ac e Th	g cc r Pr	a ga o As	t gc p Al 9	t cat a His 5	288

gat Asp	atg Met	tat Tyr	ccc Pro 100	ggt Gly	gaa Glu	aag Lys	aat Asn	gtc Val 105	acg Thr	att Ile	cat His	gta Val	gaa Glu 110	aga Arg	cgc Arg	336
aca Thr	gac Asp	gtg Val 115	tta Leu	tgc Cys	ggg Gly	cgc Arg	tca Ser 120	aga Arg	gaa Glu	gga Gly	cat His	ttt Phe 125	gac Asp	ggg Gly	gtc Val	384
gcg Ala	atc Ile 130	gta Val	ctg Leu	acg Thr	aag Lys	ctt Leu 135	ttc Phe	aat Asn	cta Leu	gtc Val	aag Lys 140	ccg Pro	act Thr	cgt Arg	gcc Ala	432
tat Tyr 145	ttc Phe	ggt Gly	tta Leu	aaa Lys	gat Asp 150	gcg Ala	cag Gln	cag Gln	gta Val	gct Ala 155	gtt Val	gtt Val	gat Asp	ggg Gly	tta Leu 160	480
atc Ile	agc Ser	gac Asp	ttc Phe	ttc Phe 165	atg Met	gat Asp	att Ile	gaa Glu	ttg Leu 170	gtt Val	cct Pro	gtc Val	gat Asp	acg Thr 175	gtc Val	528
aga Arg	gag Glu	gaa Glu	gac Asp 180	ggc Gly	tta Leu	gcc Ala	aaa Lys	agc Ser 185	tct Ser	cgc Arg	aat Asn	gta Val	tac Tyr 190	tta Leu	aca Thr	576
gct Ala	gag Glu	gaa Glu 195	aga Arg	aaa Lys	gaa Glu	gcg Ala	cct Pro 200	aag Lys	ctg Leu	tat Tyr	cgg Arg	gcc Ala 205	ctt Leu	caa Gln	aca Thr	624
agt Ser	gcg Ala 210	gaa Glu	ctt Leu	gtc Val	caa Gln	gcc Ala 215	ggt Gly	gaa Glu	aga Arg	gat Asp	cct Pro 220	gaa Glu	gcg Ala	gtg Val	ata Ile	672
aaa Lys 225	gct Ala	gca Ala	aaa Lys	gat Asp	atc Ile 230	att Ile	gaa Glu	acg Thr	act Thr	agc Ser 235	gga Gly	acc Thr	ata Ile	gac Asp	tat Tyr 240	720
gta Val	gag Glu	ctt Leu	tat Tyr	tcc Ser 245	tat Tyr	ccg Pro	gaa Glu	ctc Leu	gag Glu 250	cct Pro	gtg Val	aat Asn	gaa Glu	att Ile 255	gct Ala	768
gga Gly	aag Lys	atg Met	att Ile 260	ctc Leu	gct Ala	gtt Val	gca Ala	gtt Val 265	gct Ala	ttt Phe	tca Ser	aaa Lys	gcg Ala 270	cgt Arg	tta Leu	816
ata Ile	gat Asp	aat Asn 275	atc Ile	att Ile	att Ile	gat Asp	att Ile 280	cga Arg	gaa Glu	atg Met	gag Glu	aga Arg 285	ata Ile			858
<212)> 26 l> 28 ?> PR	16 RT	ne e	usht i	lia											

<213> Bacillus subtilis

<400> 26

Met Arg Gln Ile Thr Asp Ile Ser Gln Leu Lys Glu Ala Ile Lys Gln 1 5 10 15

Tyr His Ser Glu Gly Lys Ser Ile Gly Phe Val Pro Thr Met Gly Phe Leu His Glu Gly His Leu Thr Leu Ala Asp Lys Ala Arg Gln Glu Asn Asp Ala Val Ile Met Ser Ile Phe Val Asn Pro Ala Gln Phe Gly Pro 55 Așn Glu Asp Phe Glu Ala Tyr Pro Arg Asp Ile Glu Arg Asp Ala Ala Leu Ala Glu Asn Ala Gly Val Asp Ile Leu Phe Thr Pro Asp Ala His Asp Met Tyr Pro Gly Glu Lys Asn Val Thr Ile His Val Glu Arg Arg Thr Asp Val Leu Cys Gly Arg Ser Arg Glu Gly His Phe Asp Gly Val Ala Ile Val Leu Thr Lys Leu Phe Asn Leu Val Lys Pro Thr Arg Ala Tyr Phe Gly Leu Lys Asp Ala Gln Gln Val Ala Val Val Asp Gly Leu 155 Ile Ser Asp Phe Phe Met Asp Ile Glu Leu Val Pro Val Asp Thr Val 170 165 Arg Glu Glu Asp Gly Leu Ala Lys Ser Ser Arg Asn Val Tyr Leu Thr 185 Ala Glu Glu Arg Lys Glu Ala Pro Lys Leu Tyr Arg Ala Leu Gln Thr 195 Ser Ala Glu Leu Val Gln Ala Gly Glu Arg Asp Pro Glu Ala Val Ile 215 Lys Ala Ala Lys Asp Ile Ile Glu Thr Thr Ser Gly Thr Ile Asp Tyr 225 Val Glu Leu Tyr Ser Tyr Pro Glu Leu Glu Pro Val Asn Glu Ile Ala Gly Lys Met Ile Leu Ala Val Ala Val Ala Phe Ser Lys Ala Arg Leu Ile Asp Asn Ile Ile Ile Asp Ile Arg Glu Met Glu Arg Ile

280

<210> 27 <211> 381 <212> DNA

275

<213> Bacillus subtilis

	L> CI		(381)		ı										•	
<400)> 27	7											•			
atg Met l	tat Tyr	cga Arg	aca Thr	atg Met 5	atg Met	agc Ser	ggc Gly	aaa Lys	ctt Leu 10	cac His	agg Arg	gca Ala	act Thr	gtt Val 15	acg Thr	,48
gaa Glu	gca Ala	aac Asn	ctg Leu 20	aac Asn	tat Tyr	gtg Val	gga Gly	agc Ser 25	att Ile	aca Thr	att Ile	gat Asp	gaa Glu 30	gat Asp	ctc Leu	96
att Ile	gat Asp	gct Ala 35	gtg Val	gga Gly	atg Met	ctt Leu	cct Pro 40	aat Asn	gaa Glu	aaa Lys	gta Val	caa Gln 45	att Ile	gtg Val	aat Asn	144
aat Asn	aat Asn 50	aat Asn	gga Gly	gca Ala	cgt Arg	ctt Leu 55	gaa Glu	acg Thr	tat Tyr	att Ile	att Ile 60	cct Pro	ggt Gly	aaa Lys	cgg Arg	192
gga Gly 65	agc Ser	ggc Gly	gtc Val	ata Ile	tgc Cys 70	tta Leu	aac Asn	ggt Gly	gca Ala	gcc Ala 75	gca Ala	cgc Arg	ctt Leu	gtg Val	cag Gln 80	240
gaa Glu	gga Gly	gat Asp	aag Lys	gtc Val 85	att Ile	att Ile	att Ile	tcc Ser	tac Tyr 90	aaa Lys	atg Met	atg Met	tct Ser	gat Asp 95	caa Gln	288
gaa Glu	gcg Ala	gca Ala	agc Ser 100	cat His	gag Glu	ccg Pro	aaa Lys	gtg Val 105	gct Ala	gtt Val	ctg Leu	aat Asn	gat Asp 110	caa Gln	aac Asn	336
aaa Lys	att Ile	gaa Glu 115	caa Gln	atg Met	ctg Leu	ggg Gly	aac Asn 120	gaa Glu	cca Pro	gcc Ala	cgt Arg	aca Thr 125	att Ile	ttg Leu		381
<212 <212	0> 28 l> 12 2> Pi 3> Ba	27 RT	lus :	subt:	ilis											
<400)> 28	3														
Met 1	Tyr	Arg	Thr	Met 5	Met	Ser	Gly	Lys	Leu 10	His	Arg	Ala	Thr	Val 15	Thr	
Glu	Ala	Asn	Leu 20	Asn	Tyr	Val	Gly	Ser 25	Ile	Thr	Ile	Asp	Glu 30	Asp	Leu	
Ile	Asp	Ala 35	Val	Gly	Met	Leu	Pro 40	Asn	Glu	Lys	Val	Gln 45	Ile	Val	Asn	
Asn	Asn 50	Asn	Gly	Ala	Arg	Leu 55	Glu	Thr	Tyr	Ile	Ile 60	Pro	Gly	Lys	Arg	
Gly 65	Ser	Gly	Val	Ile	Cys 70	Leu	Asn	Gly	Ala	Ala 75	Ala	Arg	Leu	Val	Gln 80	

												•					
	Glu	Gly	Asp	Lys	Val 85	Ile :	lle :	Ile s	Ser '	Tyr 90	Lys -	Met	Met '	Ser	Asp (95	Sln	
1	Glu	Ala	Ala	Ser 100	His	Glu -	Pro⊰	Lys '	Val :	Ala	Val	Leu	Asn	Asp 110	Gln i	Asn	,
	Lys	Ile	Glu 115	Gl n	Met	Leu	Gly	Asn (120	Glu	Pro.	Ala	Arg	Thr 125	I l'e	Leu		
	<210 <211 <212 <213	> 8 !> DI	94 NA	lus :	subti	.lis											•
	<220 <221 <222	L> C	DS 1)	(894)												
,	<400 atg Met 1	aaa	att	gga Gly	att Ile 5	atc Ile	ggc Gly	gga Gly	ggc Gly	tcc Ser 10	gtt Val	ggt Gly	ctt Leu	tta Leu	tgc Cys 15	gcc Ala	48
1	tat 'Tyr	tat Tyr	ttg Leu	tca Ser 20	Leu	tat Tyr	cac His	gac Asp	gtg Val 25	act Thr	gtt Val	gtg Val	acg Thr	agg Arg 30	egg Arg	caa Gln	96 , ,
	gaa Glu	cag Gln	gct Ala 35	Ala	gcc Ala	att Ile	-cag Gln	tct Ser 40	gaa Glu	gga Gly	atc Ile	cgg Arg	ctt Leu 45	Тут	aaa Lys	ggc ∙Gly	1-44
	G1 y	gaç Glu 50	ı Glu	tto Phe	agg Arg	gct Ala	gat Asp 55	tgc Cys	agt Ser	gcg Ala	gac Asp	acg Thr	Ser	ato Ile	aat Asn	tcg Ser	192
	gac Asp 65	Phe	gad Asp	cto Lev	g ctt 1 Leu	gtc Val 70	Val	aca Thr	gtg Val	aag Lys	caq Gln 75	His	cag Gln	ctt Leu	caa Gln	tct Ser 80	240
	gtt Val	tt1	t to	g tog r Sei	g ctt Leu 85	Glu	.cga Arg	atc Ile	ggg Gly	aag Lys 90	Thr	aat Asr	ata 11e	tta Lei	ttt Phe 95	Leu	288
	caa Glr	aa Ası	c ggd n Gl	c ato y Mei	t Gly	cat His	atc Ile	cac His	gac Asp 105	Leu	a aaa a Lys	a gad s Asp	tgq Tr	g cac His	c gtt s Val	-ggc -Gly	336
	cat His	tc Se	c at r Il	e Ty	t gtt r Val	· : gga : Gly	ato / Ile	gtt Val 120	Glu	g cad	c gga s Gly	a gci y Ala	t gta a Val 12	l Ar	a aaa g Lys	tcg Ser	384
	gat Ası	ac Th	r Al	t gt a Va	t gat l Asp	cat His	aca Thr	Gly	cta Lei	a ggt i Gly	t gco y Ala	g ata a Ila 14	e Ly:	a tg s Tæ	g ago p Ser	gcg Ala	432
	tte	ga :	c ga	t gc	t gaa	a cca	gad	cgg	j ete	g aa	c at	c tt	g tt	t ∙ca	g ·cat	aac	480

Phe 145	Asp	Asp	Ala	Glu	Pro 150	Asp	Arg	Leu	Asn	Ile 155	Leu	Phe	Gln	His	Asn [,] 160	
cat His	tcg Ser	gat Asp	ttt Phe	ccg Pro 165	att Ile	tat Tyr	tat Tyr	gag Glu	acg Thr 170	gat Asp	tgg Trp	tac Tyr	cgt Arg	ctg Leu 175	ctg Leu	528 ,
acg Thr	ggc Gly	aag Lys	ctg Leu 180	att Ile	gta Val	aat Asn	gcg Ala	tgt Cys 185	att Ile	aat Asn	cct Pro	tta Leu	act Thr 190	gcg Ala	tta Leu	576
ttg Leu	caa Gln	gtg Val 195	aaa Lys	aat Asn	gga Gly	gaa Glu	ctg Leu 200	ctg Leu	aca Thr	acg Thr	cca Pro	gct Ala 205	tat Tyr	ctg Leu	gct Ala	624
ttt Phe	atg Met 210	aag Lys	ctg Leu	gta Val	ttt Phe	cag Gln 215	gag Glu	gca Ala	tgc Cys	cgc Arg	att Ile 220	tta Leu	aaa Lys	ctt Leu	gaa Glu	672
aat Asn 225	gaa Glu	gaa Glu	aag Lys	gct Ala	tgg Trp 230	gag Glu	cgg Arg	gtt Val	cag Gln	gcc Ala 235	gtt Val	tgt Cys	ggg Gly	caa Gln	acg Thr 240	720
aaa Lys	gag Glu	aat Asn	cgt Arg	tca Ser 245	tca Ser	atg Met	ctg Leu	gtt Val	gac Asp 250	gtc Val	att Ile	gga Gly	ggc Gly	cgg Arg 255	cag Gln	768
acg Thr	gaa Glu	gct Ala	gac Asp 260	gcc Ala	att Ile	atc Ile	gga Gly	tac Tyr 265	tta Leu	ttg Leu	aag Lys	gaa Glu	gca Ala 270	agt Ser	ctt Leu	816
caa Gln	ggt Gly	ctt Leu 275	gat Asp	gcc Ala	gtc Val	cac His	cta Leu 280	gag Glu	ttt Phe	tta Leu	tat Tyr	ggc Gly 285	agc Ser	atc Ile	aaa Lys	864
		gag Glu														894
<213 <213	0> 30 l> 29 2> PE 3> Ba	8	lus s	subti	ilis			•						•		
<400)> 30)														
1				5					Ser 10					15		
Tyr	Tyr	Leu	Ser 20	Leu	Tyr	His	Asp	Val 25	Thr	Val	Val	Thr	Arg 30	Arg	Gln	
Glu	Gln	Ala 35	Ala	Ala	Ile	Gln	Ser 40	Glu	Gly	Ile	Arg	Leu 45	Tyr	Lys	Gly	

Gly Glu Glu Phe Arg Ala Asp Cys Ser Ala Asp Thr Ser Ile Asn Ser 50 55 60

CHICOCOID, 1810

PCT/US00/25993

Asp Phe Asp Leu Leu Val Val Thr Val Lys Gln His Gln Leu Gln Ser 75 65 Val Phe Ser Ser Leu Glu Arg Ile Gly Lys Thr Asn Ile Leu Phe Leu Gln Asn Gly Met Gly His Ile His Asp Leu Lys Asp Trp His Val Gly 110 105 His Ser Ile Tyr Val Gly Ile Val Glu His Gly Ala Val Arg Lys Ser Asp Thr Ala Val Asp His Thr Gly Leu Gly Ala Ile Lys Trp Ser Ala Phe Asp Asp Ala Glu Pro Asp Arg Leu Asn Ile Leu Phe Gln His Asn 145 His Ser Asp Phe Pro Ile Tyr Tyr Glu Thr Asp Trp Tyr Arg Leu Leu 170 Thr Gly Lys Leu Ile Val Asn Ala Cys Ile Asn Pro Leu Thr Ala Leu Leu Gln Val Lys Asn Gly Glu Leu Leu Thr Thr Pro Ala Tyr Leu Ala 205 . 200 Phe Met Lys Leu Val Phe Gln Glu Ala Cys Arg Ile Leu Lys Leu Glu 215 Asn Glu Glu Lys Ala Trp Glu Arg Val Gln Ala Val Cys Gly Gln Thr Lys Glu Asn Arg Ser Ser Met Leu Val Asp Val Ile Gly Gly Arg Gln 250 Thr Glu Ala Asp Ala Ile Ile Gly Tyr Leu Leu Lys Glu Ala Ser Leu 265

Gln Gly Leu Asp Ala Val His Leu Glu Phe Leu Tyr Gly Ser Ile Lys 280

Ala Leu Glu Arg Asn Thr Asn Lys Val Phe 295

<210> 31 <211> 1725 <212> DNA <213> Bacillus subtilis

<220> <221> CDS <222> (1)..(1722)

<400> 31 atg ggg act aat gta cag gtg gat tca gca tct gcc gaa tgt aca cag Met Gly Thr Asn Val Gln Val Asp Ser Ala Ser Ala Glu Cys Thr Gln

1				5					10					15	·	
				Ala										gag Glu		96
gta Val	gaa Glu	atg Met 35	atc Ile	ttc Phe	ggt Gly	tat Tyr	ccg Pro 40	ggc Gly	ggg Gly	gct Ala	gtg Val	ctt Leu 45	ccg Pro	att Ile	tac Tyr	144
														cac His		192
														gga Gly		240
														ctt Leu 95		288
														gtc Val		336
aca Thr	ggg Gly	cag Gln 115	gta Val	gca Ala	acc Thr	tct Ser	gta Val 120	atc Ile	ggg Gly	agc Ser	gat Asp	gca Ala 125	ttt Phe	cag Gln	gaa Glu	384
gca Ala	gac Asp 130	att Ile	tta Leu	ggg Gly	att Ile	acg Thr 135	atg Met	cca Pro	gta Val	aca Thr	aaa Lys 140	cac His	agc Ser	tac Tyr	cag Gln	432
gtt Val 145	cgc Arg	cag Gln	ccg Pro	gaa Glu	gat Asp 150	Leu	ccg Pro	cgc Arg	atc Ile	att Ile 155	Lys	gaa Glu	gcg Ala	ttc Phe	cat His 160	480
att Ile	gca Ala	aca Thr	act Thr	gga Gly 165	Arg	ccc Pro	gga Gly	cct Pro	gta Val 170	Leu	att Ile	gat Asp	att	ecg Pro 175	Lys	528
gat Asp	gta Val	gca Ala	aca Thr 180	Ile	gaa Glu	gga Gly	gaa Glu	ttc Phe 185	Ser	tac Tyr	gat Asp	cat His	gag Glu 190	atg Met	aat Asn	576
			Tyr					Glu					Glr	atc lle	cgc Arg	624
		Val					Ser					val			gcg Ala	672
	Ala					Gly					ı Glu				tat Tyr 240	720

gct Ala	gaa Glu	cag Gln	cag Gln	caa Gln 245	atç Ile	cct Pro	gtg Val	gca Ala	cac His 250	acc Thr	ctt Leu	ttg Leu	GJ Ä ddd	ctc Leu 255	gga Gly	768
ggc Gly	ttc Phe	ccg Pro	gct Ala 260	·gac Asp	cat His	ccg Pro	ctt Leu	ttc Phe 265	cta Leu	∙ggg ∙ggg	atg Met	Ala	gga Gly 270	atg Met	cac His	816
ggt Gly	act Thr	tat Tyr 275	aca Thr	gcc Ala	aat Asn	atg Met	gcc Ala 280	ctt Leu	cat His	gaa Glu	tgt Cys	gat Asp 285	cta Leu	tta Leu	atc Ile	864
agt Ser	atc Ile 290	ggc Gly	gcc Ala	cgt Arg	ttt Phe	gat Asp 295	-gac Asp	cgt Arg	gtc Val	aca Thr	gga Gly 300	aac Asn	c.tg Leu	aaa Lys	cac His	912
ttt Phe 305	gcc Ala	aga Arg	aac Asn	gca Ala	aag Lys 310	ata Ile	gcc Ala	cac His	atc Ile	gat Asp 315	att Ile	gat A s p	-cca Pro	gct Ala	gaa Glu 320	960
atc Ile	gga Gly	aaa Lys	atc Ile	atg Met 325	Lys	aca Thr	cag Gln	att Ile	.cct Pro 330	gta Val	gtc Val	gga Gly	gac Asp	agc Ser 335	Lys	1008
att Ile	gtc Val	ctg Leu	cag Gln 340	Glu	ctg Leu	atc	aaa Lys	caa Gln 345	Asp	ggc -Gly	aaa Lys	-caa -Gln	agc Ser 350	Asp	tca Ser	1056
agc Ser	gaa Glu	tgg Trp 355	Lys	aaa Lys	cag Gln	ctc Leu	gca Ala 360	Glu	tgg Trp	aaa Lys	gaa Glu	gag Glu 365	Tyr	eco Pro	ctc Leu	1104
tgg Trp	tat Tyr 370	Va]	gat L Asp	aat Asn	gaa Glu	gaa Glu 379	•Glu	ggt Gly	ttt Phe	aaa Lys	ect Pro 380	Gla	aaa Lys	tte Lei	att Ile	1152
gaa Glu 385	туг	att : Ile	cat His	caa Glr	ttt Phe 390	Thr	aaa Lys	ı gga : Gly	gaç Glu	g gcc 1 Ala 395	Ile	gto Val	gca L Ala	acq Thi	gat Asp 400	1200
gta Val	a ggo L Gly	caq Gli	g cat n His	caa s Glr 405	Met	tge Tr	tca Sei	gcç Ala	Glr 410	n Phe	tate Tyl	c eco	g tto Phe	€ -Caa € 'Gli 41'	a aaa n Lys 5	1248
gca Ala	a gat a Asp	aaa b Ly:	a tgo s Tr _i 420	p Val	c acq L Thi	tca Sea	a ggo	gga y Gly 425	y Le	t gga u Gly	a acq	g ato r Met	g gga t -Gl: 430	y Pho	c ggt e Gly	1296
ct: Le	t cc	g gco o Ala 43	a Al	g ato a Ile	e Gly	gca Ala	a cad a Gli 440	n Lei	g gco	c gaa a Gl	a aaa u Ly:	a gad s Ası 441	o Ala	t ac	t gtt r Val	1344
gt• Va	c gc 1 Ala 45	a Va	t gt l Va	c gga 1 Gl	a gad y Asi	gg G1: 45	y Gl	a tte y Pho	c ca e Gl	a ate	g ace t Th	r Le	t ca u Gl	a⊸ga n Gl	a ctc u Leu	1392
ga As	t gt p Va	t at l Il	t cg e Ar	c ga g Gl	a tta u Lei	a aa u As	t ct: n Le	t cc	g gt o Va	c aa l Ly	g gt s Va	a gto 1 Va	g at	t tt e Le	a aat u Asn	1440

465	470	475	480
aac gct tgt ctc gga Asn Ala Cys Leu Gly 489	Met Val Arg Gln	tgg cag gaa att ttc Trp Gln Glu Ile Phe 490	tat gaa 1488 Tyr Glu 495
gaa cgt tat tca gaa Glu Arg Tyr Ser Glu 500	tct aaa ttc gct Ser Lys Phe Ala 505	tct cag cct gac ttc Ser Gln Pro Asp Phe 510	gtc aaa 1536 Val Lys
ttg tcc gaa gca tad Leu Ser Glu Ala Tys 515	ggc att aaa ggc Gly Ile Lys Gly 520	atc aga att tca tca Ile Arg Ile Ser Ser 525	gaa gcg 1584 Glu Ala
gaa gca aag gaa aag Glu Ala Lys Glu Lys 530	g ctg gaa gag gca E Leu Glu Glu Ala 535	tta aca tca aga gaa Leu Thr Ser Arg Glu 540	cct gtt 1632 Pro Val
gtc att gas gtg cgg Val Ile Asp Val Arg 545	gtt gcc agc gaa Val Ala Ser Glu 550	gaa aaa gta ttc ccg Glu Lys Val Phe Pro 555	atg gtg 1680 Met Val 560
gct ccg ggg aaa ggg Ala Pro Gly Lys Gly 56	/ Leu His Glu Met	gtg ggg gtg aaa cct Val Gly Val Lys Pro 570	tga 1725
<210> 32 <211> 574 <212> PRT <213> Bacillus subs	ilis	,	
<400> 32 Met Gly Thr Asn Val 1	. Gln Val Asp Ser	Ala Ser Ala Glu Cys 10	Thr Gln
Thr Met Ser Gly Ala 20	Leu Met Leu Ile 25	Glu Ser Leu Lys Lys 30	Glu Lys
Val Glu Met Ile Pho 35	e Gly Tyr Pro Gly 40	Gly Ala Val Leu Pro 45	Ile Tyr
Asp Lys Leu Tyr Ası 50	Ser Gly Leu Val 55	His Ile Leu Pro Arg	His Glu
Gln Gly Ala Ile His	Ala Ala Glu Gly 70	Tyr Ala Arg Val Ser 75	Gly Lys 80
Pro Gly Val Val Ile 85		Pro Gly Ala Thr Asn	Leu Val
		30	,,
Thr Gly Leu Ala Asp 100	Ala Met Ile Asp 105	Ser Leu Pro Leu Val	
100	105	Ser Leu Pro Leu Val	Val Phe

	130					135					140				
Val 145	Arg	Gln	Pro	Glu	Asp 150	Leu	Pro	Arg	Ile	Ile 155	Lys	Glu	Ala	Phe	His 160
lle	Ala	Thr	Thr	Gly 165	Arg	Pro	Gly	Pro	Val 170	Leu	Ile	Asp	Ile	Pro 175	Lys
Asp	Val	Ala	Thr 180	Ile	Glu	Gly	Glu	Phe 185	Ser	Tyr	Asp	His	Glu 190	Met	Asn
Leu	Pro	Gly 195	Tyr	Gln	Pro	Thr	Thr 200	Glu	Pro	Asn	Tyr	Leu 205	Gln	Ile	Arg
Lys	Leu 210	Val	Glu	Ala	Val	Ser 215	Ser	Ala	Lys	Lys	Pro 220	Val	Ile	Leu	Ala
Gly 225	Ala	Gly	Val	Leu	His 230	Gly	Lys	Ala	Ser	Glu 235	Glu	Leu	Lys	Asn	Tyr 240
Ala	Glu	Gln	Gln	Gln 245	Ile	Pro	Val	Ala	His 250	Thr	Leu	Lėu	Gly	Leu 255	Gly
Gly	Phe	Pro	Ala 260	Asp	His	Pro	Leu	Phe 265	Leu	Gly	Met	Äla	Gly 270	Met	His
Gly	Thr	Tyr 275	Thr	Ala	Asn	Met	Ala 280	' Leu	His	Glu	Cys	Asp 285	Leu	Leu	Ile
Ser	11e 290	Gly	Ala	Arg	Phe	Asp 295	Asp	Arg	Val	Thr	Gly 300	Asn	Leu	Lys	His
Phe 305	Ala	Arg	Asn	Ala	Lys 310	Ile	Ala	His	Ile	Asp 315	Ile	Asp	Pro	Ala	G11 320
Ile	Gly	Lys	Ile	Met 325	Lys	Thr	Gln	Ile	Pro 330	Val	Val	Gly	Asp	Ser 335	_
Ile	Val	Leu	Gln 340	Glu	Leu	Ile	Lys	Gln 345	Asp	Gly	Łys	Gl·n	Ser 350	Asp	'Se
Ser	Glu	355	Lys	Lys	Gln	Leu	Ala 3 6 0		Trp	Lys	Glu	Glu 365	Tyr	Pro	Le
Trp	Tyr 370		Asp	Asn	Glu	Glu 375	Glu	Gly	Phe	Lys	Pro 380		Lys	Leu	116
Glu 385	_	Ile	His	Gln	Phe 390		Lys	Gly	Glu	Ala 395		Val	Ala	Thr	As ₁
Val	Gly	Gln	His	Gln 405		Trp	Ser	Ala	Gln 410		Tyr	Pro	₽he	Gln 415	
Ala	Asp	Lys	Trp 420		Thr	Ser	Gly	Gly 425		Gly	Thr	Met	Gly 430		Gl
Leu	Pro	Ala 435		Ile	Gly	Ala	Gln 440		Ala	Glu	Lys	Asp	Ala	Thr	Va:

	A1a 450	Val	Val	Gly	Asp	Gly 455	Gly	Phe	Gln	Met	Thr 460	Leu	Gln	Glu	Leu	
Asp 465	Val	Ile	Arg	Glụ	Leu 470	Asn	Leu	Pro	Val	Lys 475	Val	Val	Ile	Leu	Asn 480	
Asn	Ala	Cys	Leu	Gly 485	Met	Val	Arg	Gln	Trp 490	Gln	Glu	Ile	Phe	Tyr 495	Glu	
Glu	Arg	Tyr	Ser 500	Glu	Ser	Lys	Phe	Ala 505	Ser	Gln	Pro	Asp	Phe 510	Val	Lys	
Leu	Ser	Glu 515	Ala	Tyr	Gly	Ile	Lys 520	Gly	Ile	Arg	Ile	Ser 525	Ser	Glu	Ala	
Glu	Ala 530	Lys	Glu	Lys	Leu	Glu 535	Glu	Ala	Leu	Thr	Ser 540	Arg	Glu	Pro	Val	
Val 545	Ile	Asp	Val	Arg	Val 550	Ala	Ser	Glu	Glu	Lys 555	Val	Phe '	Pro	Met	Val 560	
Ala	Pro	Gly	Lys	Gly 565	Leu	His	Glu	Met	Val 570	Gly	Val	Lys	Pro			
<21 <21	0> 3: 1> 5: 2> Di 3> Ba	25 NA	lus s	subti	ilis	ι.										
	0> 1> CI 2> (1		(522))			٠									
<22 <22 <40	1> CI 2> (1 0> 31	L)					٠									
<22 <22 <40 ttg	1> CI 2> (1	l) 3 aga	att	atc												48
<22 <22 <40 ttg Met 1	1> CI 2> (1 0> 31 aaa	aga Arg	att Ile	atc Ile 5 ggt	Thr	Leu	Thr	Val , aaa	Val 10 agg	Asn	Arg tac	Ser	Gly att	Val 15 gaa	Leu	48 96
<22 <22 <40 ttg Met 1 aac Asn	1> CI 2> (3 0> 33 aaa Lys	aga Arg atc Ile	att Ile acc Thr 20	atc Ile 5 ggt Gly	Thr cta Leu aca	ttc Phe gaa	Thr aca Thr	val aaa Lys 25	Val 10 agg Arg	Asn cat His	tac Tyr	Ser aac Asn	Gly att Ile 30 atc	Val 15 gaa Glu acc	agc Ser	
<22 <22 <40 ttg Met 1 aac Asn att Ile	1> CI 2> (1 0> 31 aaa Lys cgg Arg	aga Arg atc Ile gtt Val 35	att Ile acc Thr 20 gga Gly	atc Ile 5 ggt Gly cac His	Thr cta Leu aca Thr	ttc Phe gaa Glu	Thr aca Thr aca Thr 40	val aaa Lys 25 gcc Ala	Val 10 agg Arg ggc Gly	Asn cat His gtt Val	tac Tyr tcc Ser	aac Asn aga Arg 45	att Ile 30 atc Ile	Val 15 gaa Glu acc Thr	agc Ser ttc Phe	96
<22 <20 ttg Met 1 aac Asn att Ile gtc Val	1 > CI 2 > (: 0 > 3: aaa Lys cgg Arg aca Thr yal 50 aac Asn	aga Arg atc Ile gtt Val 35 Cat His	att Ile acc Thr 20 gga Gly gtt Val	atc Ile 5 ggt Gly cac His gaa Glu	Thr cta Leu aca Thr ggt Gly	ttc Phe gaa Glu gaa Glu 55	Thr aca Thr aca Thr 40 aat Asn	val aaaa Lys 25 gcc Ala gat Asp	Val 10 agg Arg ggc Gly gtt Val	Asn cat His gtt Val gaa Glu aca	tac Tyr tcc Ser cag Gln 60	aac Asn aga Arg 45 tta Leu	att Ile 30 atc Ile acg Thr	Val 15 gaa Glu acc Thr aaa Lys	agc Ser ttc Phe cag Gln	96 144

85	90		95
		•	

tca aca aga aca gag att aat gga atc ata gaa ccg ttt aga gcc tct Ser Thr Arg Thr Glu Ile Asn Gly Ile Ile Glu Pro Phe Arg Ala Ser 1.05 gtc gtt gat gtc agc aga gac agc atc gtt gtt cag gtg aca ggt gaa 384 Val Val Asp Val Ser Arg Asp Ser Ile Val Val Gln Val Thr Gly Glu 120 tct aac aaa att gaa gcg ctt att gag tta tta aaa cct tat ggc att 432 Ser Asn Lys Ile Glu Ala Leu Ile Glu Leu Leu Lys Pro Tyr Gly Ile 135 480 aaa gaa atc gcg aga aca ggt aca acg gct ttt gcg agg gga acc agc Lys Glu Ile Ala Arg Thr Gly Thr Thr Ala Phe Ala Arg Gly Thr Ser 150 155 525 aaa agg cgt cat cca ata aaa caa tat cta ttg tat aaa aca taa Lys Arg Arg His Pro Ile Lys Gln Tyr Leu Leu Tyr Lys Thr 170

<210> 34 <211> 174 <212> PRT <213> Bacillus subtilis

<400> 34
Met Lys Arg Ile Ile Thr Leu Thr Val Val Asn Arg Ser Gly Val Leu
1 5 10 15

Asn Arg Ile Thr Gly Leu Phe Thr Lys Arg His Tyr Asn Ile Glu Ser 20 25 30

Ile Thr Val Gly His Thr Glu Thr Ala Gly Val Ser Arg Ile Thr Phe 35 40 45

Val Val His Val Glu Gly Glu Asn Asp Val Glu Gln Leu Thr Lys Gln
50 55 60

Leu Asn Lys Gln Ile Asp Val Leu Lys Val Thr Asp Ile Thr Asn Gln 65 70 75 80

Ser Ile Val Gln Arg Glu Leu Ala Leu Ile Lys Val Val Ser Ala Pro 85 90 95

Ser Thr Arg Thr Glu Ile Asn Gly Ile Ile Glu Pro Phe Arg Ala Ser 100 105 110

Val Val Asp Val Ser Arg Asp Ser Ile Val Val Gln Val Thr Gly Glu 115 120 125

Ser Asn Lys Ile Glu Ala Leu Ile Glu Leu Leu Lys Pro Tyr Gly Ile 130 135 140

Lys Glu Ile Ala Arg Thr Gly Thr Thr Ala Phe Ala Arg Gly Thr Ser 145 150 155 160 Lys Arg Arg His Pro Ile Lys Gln Tyr Leu Leu Tyr Lys Thr 165 170

<210> 35 <211> 1029 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(1026) <400> 35 atg gta aaa gta tat tat aac ggt gat atc aaa gag aac gta ttg gct Met Val Lys Val Tyr Tyr Asn Gly Asp Ile Lys Glu Asn Val Leu Ala gga aaa aca gta gcg gtt atc ggg tac ggt tcg caa ggc cac gca cat 96 Gly Lys Thr Val Ala Val Ile Gly Tyr Gly Ser Gln Gly His Ala His 25 gcc ctg aac ctt aaa gaa agc gga gta gac gtg atc gtc ggt gtt aga Ala Leu Asn Leu Lys Glu Ser Gly Val Asp Val Ile Val Gly Val Arg 35 caa gga aaa tot tto act caa goo caa gaa gac gga cat aaa gta ttt 192 Gln Gly Lys Ser Phe Thr Gln Ala Gln Glu Asp Gly His Lys Val Phe tca gta aaa gaa gcg gca gcc caa gcc gaa atc atc atg gtt ctg ctt Ser Val Lys Glu Ala Ala Ala Gin Ala Glu Ile Ile Met Val Leu Leu ccg gat gag cag caa aaa gta tac gaa gct gaa atc aaa gat gaa 288 Pro Asp Glu Gln Gln Lys Val Tyr Glu Ala Glu Ile Lys Asp Glu 85 ttg aca gca gga aaa tca tta gta ttc gct cat gga ttt aac gtg cat 336 Leu Thr Ala Gly Lys Ser Leu Val Phe Ala His Gly Phe Asn Val His 100 105 ttc cat caa att gtt cct ccg gcg gat gta gat gta ttc tta gtg gcc 384 Phe His Gln Ile Val Pro Pro Ala Asp Val Asp Val Phe Leu Val Ala 115 120 125 cct aaa ggc ccg gga cac ttg gta aga aga aca tat gag caa gga gct 432 Pro Lys Gly Pro Gly His Leu Val Arg Arg Thr Tyr Glu Gln Gly Ala 130 135 140 ggc gta cct gca ttg ttc gca atc tat caa gat gtg act gga gaa gca 480 Gly Val Pro Ala Leu Phe Ala Ile Tyr Gln Asp Val Thr Gly Glu Ala 145 150 aga gac aaa gcc ctc gct tat gct aaa gga atc ggc ggc gca aga gcg Arg Asp Lys Ala Leu Ala Tyr Ala Lys Gly Ile Gly Gly Ala Arg Ala 165 170

DAICOCCIO: -MIC. 012177282 1 -

ggc Gly	gta Val	tta Leu	gaa Glu 180	acg Thr	aca Thr	ttt Phe	aaa Lys	gaa Glu 185	gaa Glu	aca Thr	gaa Glu	aca Thr	gat Asp 190	ttg Leu	ttc Phe	576
ggt Gly	gag Glu	caa Gln 195	gca Ala	gtt Val	ctt Leu	tgc Cys	ggc Gly 200	gga Gly	tta Leu	agc Ser	gcg Ala	ctt Leu 205	gtc Val	aaa Lys	gcc Ala	624
gga Gly	ttt Phe 210	gaa Glu	acc Thr	tta Leu	act Thr	gaa Glu 215	gca Ala	ggt Gly	tat Tyr	Gln	cct Pro 220	gaa Glu	ctt Leu	gca Ala	tac Tyr	672
ttc Phe 225	gag Glu	tgt Cys	ctt Leu	cat His	gag Glu 230	ctg Leu	aaa Lys	tta Leu	atc Ile	gta Val 235	gac Asp	ctt Leu	atg Met	tac Tyr	gaa Glu 240	720
gaa Glu	gga Gly	ctt Leu	gca Ala	gga Gly 245	atg Met	aga Arg	tat Tyr	tca Ser	atc Ile 250	tct Ser	gac Asp	aca Thr	gca Ala	cag Gln 255	tgg Trp	768
gga Gly	gat Asp	ttc Phe	gta Val 260	tca Ser	ggc Gly	ect Pro	ege Arg	gtt Val 265	gtg Val	gac Asp	gcc Ala	aaa Lys	gta Val 270	aaa Lys	gaa Glu	816
tct Ser	atg Met	aaa Lys 275	Glu	gta Val	tta Leu	aaa Lys	gat Asp 280	Ile	caa Gln	aac Asn	ggt Gly	aca Thr 285	ttc Phe	gca Ala	aaa Lys	854
gag Glu	tgg Trp 290	Ile	gtc Val	gaa Glu	aac Asn	caa Gln 295	Val	aac Asn	cgt Arg	cct Pro	cgt Arg 300	Phe	aac Asn	gct Ala	atc Ile	912
aat Asn 305	Ala	ago Ser	gag Glu	aac Asn	gaa Glu 310	His	caa Gln	atc Ile	gaa Glu	gta Val 315	. Val	gga Gly	aga Arg	aag Lys	Ctt Leu 320	960
cgt Arg	gaa Glu	ato Met	atg : Met	Pro 325	Phe	gtg Val	aaa Lys	caa Gln	ggc Gly 330	Lys	aaq Lys	aaq Lys	gaa Glu	gcg Ala 335	gtg Val	1008
_		_		g caa Glm			1									1029
<21 <21	.0> 3 .1> 3 .2> E	842 PRT	llus	subt	:ilis	5										
			s Vai		Tyi	c Ası	n Gl	y Asp) Ile 10		s Gl	u Ası	n Vai	L Le	u Ala 5	
Gly	y Lys	s Th	r Va:		a Val	l Ile	e Gl	y Ty: 25		y Se:	r Gl	n Gl	y Hi:		a His	

Ala Leu Asn Leu Lys Glu Ser Gly Val Asp Val Ile Val Gly Val Arg 40 Gln Gly Lys Ser Phe Thr Gln Ala Gln Glu Asp Gly His Lys Val Phe Ser Val Lys Glu Ala Ala Ala Gln Ala Glu Ile Ile Met Val Leu Leu Pro Asp Glu Gln Gln Lys Val Tyr Glu Ala Glu Ile Lys Asp Glu Leu Thr Ala Gly Lys Ser Leu Val Phe Ala His Gly Phe Asn Val His 105 Phe His Gln Ile Val Pro Pro Ala Asp Val Asp Val Phe Leu Val Ala 120 Pro Lys Gly Pro Gly His Leu Val Arg Arg Thr Tyr Glu Gln Gly Ala 135 Gly Val Pro Ala Leu Phe Ala Ile Tyr Gln Asp Val Thr Gly Glu Ala Arg Asp Lys Ala Leu Ala Tyr Ala Lys Gly Ile Gly Gly Ala Arg Ala Gly Val Leu Glu Thr Thr Phe Lys Glu Glu Thr Glu Thr Asp Leu Phe Gly Glu Gln Ala Val Leu Cys Gly Gly Leu Ser Ala Leu Val Lys Ala Gly Phe Glu Thr Leu Thr Glu Ala Gly Tyr Gln Pro Glu Leu Ala Tyr Phe Glu Cys Leu His Glu Leu Lys Leu Ile Val Asp Leu Met Tyr Glu Glu Gly Leu Ala Gly Met Arg Tyr Ser Ile Ser Asp Thr Ala Gln Trp 250 Gly Asp Phe Val Ser Gly Pro Arg Val Val Asp Ala Lys Val Lys Glu Ser Met Lys Glu Val Leu Lys Asp Ile Gln Asn Gly Thr Phe Ala Lys 280 Glu Trp Ile Val Glu Asn Gln Val Asn Arg Pro Arg Phe Asn Ala Ile 295 Asn Ala Ser Glu Asn Glu His Gln Ile Glu Val Val Gly Arg Lys Leu 310 Arg Glu Met Met Pro Phe Val Lys Gln Gly Lys Lys Glu Ala Val 325

Val Ser Val Ala Gln Asn

012177242 1 -

- 44 -

340

<211 <212	3> 37 > 16 2> DN 3> Ba	574 IA	.us s	subt i	lis								•			•
<220> <221> CDS <222> (1)(1674)																
<400)> 37	1									•					
_	gca Ala	-		-	-		-						-	-	_	48
	cac His															96
	aag Lys															144
	cat His 50															192
	gaa Giu															240
	ggc Gly															288
	gaa Glu															336
	gac Asp															384
	ctt Leu 130															432
	gga Gly															480
	tcc. Ser									Tyr						528
aac	gaa	aac	gag	ctt	caa	gaa	cta	gag	cag	ttc	gga	ŧgc	сса	acg	tgc	576

8NSDOCIO: <WO 0121772A2 1 >

Asn	Glu	Asn	Glu 180	Leu	Gln	Glu	Leu	Glu 185	Gln	Phe	Gly	Cys	Pro 190	Thr	Cys	
							acg Thr 200									624
							ccg Pro									672
							gtg Val									720
_			-		_		aaa Lys		-	_		-		-		768
gcg Ala	att Ile	gat Asp	aac Asn 260	gcg Ala	ttt Phe	gca Ala	ctc Leu	gat Asp 265	atg Met	gcg Ala	ctc Leu	gga Gly	ggt Gly 270	tct Ser	aca Thr	816
							gcc Ala 280									864
							gaa Glu									912
-	_	_			-	_	gat Asp				-	-			-	960
							ctg Leu									1008
							act Thr									1056
							aag Lys 360									1104
-					_	-	gga Gly			_	_					1152
							att Ile									1200
							gct Ala									1248

gcg Ala	ctt Leu	gac Asp	ggc Gly 420	att Ile	atc Ile	'aac Asn	cga Arg	aaa Lys 425	gta Val	aaa Lys	gaa Glu	ggc Gly	gac Asp 430	gtt Val	gtc Val	1296
atc Ile	atc Ile	aga Arg 435	tac Tyr	gaa Glu	ggg Gly	cca Pro	aaa Lys 440	ggc Gly	gga Gly	cct Pro	ggc Gly	atg Met 445	Pro	gaa Glu	atg Met	1344
ctg Leu	gcg Ala 450	cca Pro	aca Thr	tcc Ser	caa Gln	atc Ile 455	gtt Val	gga Gly	atg Met	gga Gly	ctc Leu 460	ggg Gly	510 CCá	aaa Lys	gtg Val	1392
gca Ala 465	ttg Leu	att Ile	acg Thr	Asp	gga Gly 470	cgt Arg	ttt Phe	tcc Ser	gga Gly	gcc Ala 475	tcc Ser	cgt Arg	ggc Gly	ctc Leu	tca Ser 480	1440
												ecg, Pro				1488
												aaa Lys				1536
												aaa Lys 525				1584
							Lys					gca Ala				1632
aaa Lys 545	Leu	gtg Val	aca Thr	agt Ser	gcc Ala 550	Asn	acc Thr	ggc Gly	ggt	att Ile 555	Met	aaa Lys	atc Ile			1674
<21 <21	0> 3 1> 5 2> P 3> B	58 RT	lus	subt	ilis							٠				
	0> 3 Ala		. Leu	Arg	Ser	Asn	Met	Ile			Gly	, Ile	Asp		Ala	
1 Pro		Arg	, Ser	5 Leu		Arg	Ala	Ala	10 Gly		. Lys	s Glu	Glu	15 Asp	Phe	
G) v	Lve	Pro	20		. מו	Val	Cve	25 : Asr		- Tvr	- T.).e	Asr	30 Tle		. Pro	
_		35	•				40)				45	•			
Gly	/ His		. His	Leu	Glr	55		e Gly	/ Lys	: Ile	• Val		Glu	n Ala	Ile	
Arg 65		a Ala	Gly	/ Gly	/ Va] 7(Phe	e Glu	Phe	Asr 75		: Ile	Gly	/ Val	Asp 80	

Asp	Gly	Ile	Ala	Met 85	Gly	His	Ile	Gly	Met 90	Arg	Tyŗ	Ser	Leu	Pro 95	Ser
Arg	Glu	Ile	Ile 100	Ala	Asp	Ser	Val	Glu 105	Thr	Val	Val	Ser	Ala 110	His	Trp
Phe	Asp	Gly 115	Met	Val	Cys	Ile	Pro 120	Asn	Cys	Asp	Lys	Ile 125	Thr	Pro	Gly
Met	Leu 130	Met	Ala	Ala	Met	Arg 135	Ile	Asn	Ile	₽ro	Thr 140	Ile	Phe	Val	Ser
Gly 145	Gly	Pro	Met	Ala	Ala 150	Gly	Arg	Thr	Ser	Tyr 155	Gly	Arg	Lys	Ile	Ser 160
Leu	Ser	Ser	Val	Phe 165	Glu	Gly	Val	Gly	Ala 170	Tyr	Gln	Ala	Gly	Lys 175	Ile
Asn	Glu	Asn	Glu 180	Leu	Gln	Glu	Leu	Glu 185	Gln	Phe	Gly	Cys	Pro 190	Thr	Cys
Gly	Ser	Cys 195	Ser	Gly	Met	Phe	Thr 200	Ala	Asn	Ser	Met	Asn 205	Cys	Leu	Ser
Glu	Ala 210	Leu	Gly	Leu	Ala	Leu 215	Pro	Gly	Asn	Gly	Thr 220	Ile	Leu	Ala	Thr
Ser 225	Pro	Glu	Arg	Lys	Glu 230	Phe	Val	Arg	Lys	Ser 235	Ala	Ala	Gln	Leu	Met 240
Glu	Thr	Ile	Arg	Lys 245	Asp	Ile	Lys	Pro	Arg 250	Asp	Ile	Val	Thr	Val 255	Lys
Ala	Ile	Asp	Asn 260	Ala	Phe	Ala	Leu	Asp 265	Met	Ala	Leu	Gly	Gly 270	Ser	Thr
Asn	Thr	Val 275	Leu	His	Thr	Leu	Ala 280	Leu	Ala	Asn	Glu	Ala 285	Gly	Val	Glu
Tyr	Ser 290	Leu	Glu	Arg	Ile	Asn 295	Glu	Val	Ala	Glu	Arg 300	Val	Pro	His	Leu
Ala 305	Lys	Leu	Ala	Pro	Ala 310	Ser	Asp	Val	Phe	Ile 315	Glu	Asp	Leu	His	Glu 320
Ala	Gly ·	Gly	Val	Ser 325	Ala	Ala	Leu	Asn	Glu 330		Ser	Łys	Lys	Glu 335	Gly
Ala	Leu	His	Leu 340	Asp	Ala	Leu	Thr	Val 345		Gly	Lys	Thr	Leu 350	Gly	Glu
Thr	Ile	Ala 355	Gly	His	Glu	Val	Lys 360	Asp	Tyr	Asp	Val	11e 365		Pro	Leu
Asp	Gln 370	Pro	Phe	Thr	Glu	Lys 375	Gly	Gly	Leu	Ala	Val 380	Leu	Phe	Gly	Asn

Leu Ala Pro Asp Gly Ala Ile Ile Lys Thr Gly Gly Val Gln Asn Gly 395 400 390 ' Ile Thr Arg His Glu Gly Pro Ala Val Val Phe Asp Ser Gln Asp Glu 410 Ala Leu Asp Gly Ile Ile Asn Arg Lys Val Lys Glu Gly Asp Val Val 420 Ile Ile Arg Tyr Glu Gly Pro Lys Gly Gly Pro Gly Met Pro Glu Met Leu Ala Pro Thr Ser Gln Ile Val Gly Met Gly Leu Gly Pro Lys Val 455 Ala Leu Ile Thr Asp Gly Arg Phe Ser Gly Ala Ser Arg Gly Leu Ser Ile Gly His Val Ser Pro Glu Ala Ala Glu Gly Gly Pro Leu Ala Phe Val Glu Asn Gly Asp His Ile Ile Val Asp Ile Glu Lys Arg Ile Leu 505 Asp Val Gln Val Pro Glu Glu Glu Trp Glu Lys Arg Lys Ala Asn Trp 520 515 Lys Gly Phe Glu Pro Lys Val Lys Thr Gly Tyr Leu Ala Arg Tyr Ser Lys Leu Val Thr Ser Ala Asn Thr Gly Gly Ile Met Lys Ile 555 545 <210> 39 <211> 194 <212> DNA <213> Artificial Sequence <223> Description of Artificial Sequence:promoter sequence <220> <221> -35 signal <222> (136)..(141) <220> <221> -10 signal <222> (159)..(164) gctattgacg acagctatgg ttcactgtcc accaaccaaa actgtgctca gtaccgccaa 60 tatttctccc ttgaggggta caaagaggtg tccctagaag agatccacgc tgtgtaaaaa 120

```
gcaaccccgc ctgt
                                                                   194
<210> 40
<211> 163
<212> DNA
'<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence:promoter
      sequence
<220>
<221> -35 signal
<222> (113)..(118)
<220>
<221> -10_signal
<222> (136)..(141)
gcctacctag cttccaagaa agatatccta acagcacaag agcggaaaga tgttttgttc 60
tacatccaga acaacctctg ctaaaattcc tgaaaaattt tgcaaaaagt tgttgacttt 120
atctacaagg tgtggtataa taatcttaac aacagcagga cgc
                                                                   163
<210> 41
<211> 127
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence:promoter
      sequence
<220>
<221> -35 signal
<222> (34)..(39)
<220>
<221> -10 signal
<222> (58)..(63)
<220>
<221> -35_signal
<222> (75)..(80)
<220>
<221> -10 signal
<222> (98)..(103)
<400> 41
gaggaatcat agaattttgt caaaataatt ttattgacaa cgtcttatta acqttgatat 60
aatttaaatt ttatttgaca aaaatgggct cgtgttgtac aataaatgta gtgaggtgga 120
```

tgcaatg	1.27
<210> 42 <211> 24 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 42 taaacatgag gaggagaaaa catg	24
<210> 43 <211> 28 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	·
<400> 43 attcgagaaa tggagagaat ataatatg	. 28
<210> 44 <211> 13 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 44 agaaaggagg tga	13
<210> 45 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 45 ttaagaaagg aggtgannnn atg <210> 46	.23
<210 46 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome	

binding site

<400> 46 ttagaaagga ggtgannnnn atg	23
<210> 47 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	,
<400> 47 agaaaggagg tgannnnnn atg	23
<210> 48 <211> 22 <212> DNA <213> Artificial Sequence	23
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 48 agaaaggagg tgannnnna tg	22
<210> 49 <211> 25 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	ı
<400> 49 ccctctagaa ggaggagaaa acatg	25
<210> 50 <211> 24 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 50 ccctctagag gaggagaaaa catg	24
<210> 51 <211> 23 <212> DNA	

<213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 51 ttagaaagga ggatttaaat atg	, 23
<210> 52 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 52 ttagaaagga ggtttaatta atg	. 23
<pre><210> 53 <211> 23 <212> DNA <213> Artificial Sequence</pre>	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 53 ttagaaagga ggtgatttaa atg	-23
<210> 54 <211> 23 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	-
<400> 54 ttagaaagga ggtgtttaaa atg	-23
<210> 55 <211> 28 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 55 attcgagaaa ggaggtgaat ataatatg	28
<210> 56	

<211> 27 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 56 attcgagaaa ggaggtgaat aataatg 27	
<210> 57 <211> 28 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 57 attcgtagaa aggaggtgaa ttaatatg 28	
<210> 58 <211> 3291 <212> DNA <213> Bacıllus subtilis	
<400> 58 atggggacta atgtacaggt ggattcagca tctgccgaat gtacacagac gatgagcgga 60	
gcattaatgc tgattgaatc attaaaaaaa gagaaagtag aaatgatctt cggttatccg 120	0
ggcggggctg tgcttccgat ttacgataag ctatacaatt cagggttggt acatatcctt 18	0
ccccgtcacg aacaaggagc aattcatgca gcggagggat acgcaagggt ctccggaaaa 24	0
ccgggtgtcg tcattgccac gtcagggccg ggagcgacaa accttgttac aggccttgct 30	0
gatgccatga ttgattcatt gccgttagtc gtctttacag ggcaggtagc aacctctgta 36	0
atcgggagcg atgcatttca ggaagcagac attttaggga ttacgatgcc agtaacaaaa 424	0
cacagetace aggitegeca geeggaagat etgeegegea teattaaaga agegiteeat 48	0
attgcaacaa ctggaagacc cggacctgta ttgattgata ttccgaaaga tgtagcaaca 54	0
attgaaggag aattcagcta cgatcatgag atgaatctcc cgggatacca gccgacaaca 60	0
gagccgaatt atttgcagat ccgcaagctt gtggaagccg tgagcagtgc gaaaaaaccg 66	0
gtgatcctgg cgggtgcggg cgtactgcac ggaaaagcgt cagaagaatt aaaaaattat 72	0
getgaacage ageaaateee tgtggcacae accettttgg ggeteggagg etteeegget 78	0

gaccatccgc ttttcctagg gatggcggga atgcacggta cttatacagc caatatggcc 840

DESCRIPTION 013177383 | 1

cttcatgaat gtgatctatt aatcagtatc ggcgcccgtt ttgatgaccg tgtcacagga 900 aacctgaaac actttgccag aaacgcaaag atagcccaca tcgatattga tccagctgaa 960 atcggaaaaa tcatgaaaac acagatteet gtagteggag acageaaaat tgteetgeag 1020 gagetgatea aacaagaegg caaacaaage gatteaageg aatggaaaaa acagetegea 1080 gaatggaaag aagagtatcc gctctggtat gtagataatg aagaagaagg ttttaaacct 1140 cagaaattga ttgaatatat tcatcaattt acaaaaggag aggccattgt cgcaacggat 1200 gtaggccagc atcaaatgtg gtcagcgcaa ttttatccgt tccaaaaagc agataaatgg 1260 gtcacgtcag gcggacttgg aacgatggga ttcggtcttc cggcggcgat cggcgcacag 1320 ctggccgaaa aagatgctac tgttgtcgcg gttgtcggag acggcggatt ccaaatgacg 1380 cttcaagaac tcgatgttat tcgcgaatta aatcttccgg tcaaggtagt gattttaaat 1440 aacgettgte teggaatggt cagacagtgg caggaaattt tetatgaaga aegttattea 1500 gaatctaaat togottotoa gootgactto gtoaaattgt oogaagcata oggoattaaa 1560 ggcatcagaa tttcatcaga agcggaagca aaggaaaagc tggaagaggc attaacatca 1620 agagaacctg ttgtcattga cgtgcgggtt gccagcgaag aaaaagtatt cccgatggtg 1680 gctccgggga aagggctgca tgaaatggtg ggggtgaaac cttgaaaaga attatcacat 1740 tgactgtggt gaaccgctcc ggggtgttaa accggatcac cggtctattc acaaaaaggc 1800 attacaacat tgaaagcatt acagttggac acacagaaac agccggcgtt tccagaatca 1860 ccttcgtcgt tcatgttgaa ggtgaaaatg atgttgaaca gttaacgaaa cagctcaaca 1920 aacagattga tgtgctgaaa gtcacagaca tcacaaatca atcgattgtc cagagggagc 1980 tggccttaat caaggttgtc tccgcacctt caacaagaac agagattaat ggaatcatag 2040 aaccgtttag agcctctgtc gttgatgtca gcagagacag catcgttgtt caggtgacag 2100 gtgaatctaa caaaattgaa gcgcttattg agttattaaa accttatggc attaaagaaa 2160 tcgcgagaac aggtacaacg gcttttgcga ggggaaccag caaaaggcgt catccaataa 2220 aacaatatct attgtataaa acataacaag ggagagattg aaatggtaaa agtatattat 2280 aacggtgata tcaaagagaa cgtattggct ggaaaaacag tagcggttat cgggtacggt 2340 togcaaggoo acgcacatgo cotgaacott aaagaaagog gagtagacgt gatogtoggt 2400 gttagacaag gaaaatcttt cactcaagcc caagaagacg gacataaagt attttcagta 2460 aaagaagcgg cagcccaagc cgaaatcatc atggttctgc ttccggatga gcagcagcaa 2520 aaagtatacg aagctgaaat caaagatgaa ttgacagcag gaaaatcatt agtattcgct 2580

337

catggattta acgtgcattt ccatcaaatt gttcctccgg cggatgtaga tgtattctta 2640 gtggccccta aaggcccggg acacttggta agaagaacat atgagcaagg agctggcgta 2700 cctgcattgt tcgcaatcta tcaagatgtg actggagaag caagagacaa agccctcgct 2760 tatgctaaag gaatcggcgg cgcaagagcg ggcgtattag aaacgacatt taaagaagaa 2820 acagaaacag atttgttcgg tgagcaagca gttctttgcg gcggattaag cgcgcttgtc 2880 aaagccggat ttgaaacctt aactgaagca ggttatcage ctgaacttgc atacttcgag 2940 tgtcttcatg agctgaaatt aatcgtagac cttatgtacg aagaaggact tgcaggaatg 3000 agatattcaa tctctgacac agcacagtgg ggagatttcg tatcaggccc tcgcgttgtg 3060 gacgccaaag taaaagaatc tatgaaagaa gtattaaaag atatccaaaa cggtacattc 3120 gcaaaagagt ggatcgtcga aaaccaagta aaccgtcctc gtttcaacgc tatcaatgca 3180 agcgagaacg aacatcaaat cgaagtagtg ggaagaaagc ttcgtgaaat gatgccgttt 3240 gtgaaacaag gcaagaagaa ggaagcggtg gtctccgttg cgcaaaatta a 3291

```
<210> 59
<211> 2363
<212> DNA
<213> Bacillus subtilis
<220>
<221> CDS
<222> (242)..(1072)
<220>
<221> CDS
<222> (1077)..(1934)
<220>
<221> CDS
<222> (1939)..(2319)
<400> 59
ttggtacaag cccgttgatt ttggtatact tccattgggc agtatcqcct qcqaactqca 60
cctattatta aaatagatag acattgcagc agtctgcctt gatccaaaaa aqqactqqqa 120
cagagggatg aaactcgccg aactttagaa agtgaagaat ccttctcgtt gtaacggaag 180
gtttttttggc ttgcagaaga aaacggcaga tcatctcctc taaacatgag gaggagaaaa 240
c atg aaa aca aaa ctg gat ttt cta aaa atg aag gag tct gaa gaa ccg 289
  Met Lys Thr Lys Leu Asp Phe Leu Lys Met Lys Glu Ser Glu Glu Pro
                                       10
```

att gtc atg ctg acc gct tat gat tat ccg gca gct aaa ctt gct gaa

Ile Val Met Leu Thr Ala Tyr Asp Tyr Pro Ala Ala Lys Leu Ala Glu

DEICEOCIE: AMO 010177080 I >

			20					25					30			
aa iln	gcg Ala	gga Gly 35	gtt Val	gac Asp	atg Met	att Ile	tta Leu 40	gtc Val	ggt (Gly)	gat Asp	tca Ser	ctt Leu 45	gga Gly	atg Met	gtc Val	385
jtc /al	ctc Leu 50	ggc Gly	ctt Leu	gat Asp	tca Ser	act Thr 55	gtc Val	ggt Gly	gtg Val	aca Thr	gtt Val 60	gcg Ala	gac Asp	atg Met	atc Ile	433 .
cat His 65	cat His	aca Thr	aaa Lys	gcc Ala	gtt Val 70	aaa Lys	agg Arg	ggt Gly	gcg Ala	ccg Pro 75	aat Asn	acc Thr	ttt Phe	att Ile	gtg Val 80	481
aca Thr	gat Asp	atg Met	ccg Pro	ttt Phe 85	atg Met	tct Ser	tat Tyr	cac His	ctg Leu 90	tct Ser	aag Lys	gaa Glu	gat Asp	acg Thr 95	ctg Leu	529
aaa Lys	aat Asn	gca Ala	gcg Ala 100	gct Ala	atc Ile	gtt Val	cag Gln	gaa Glu 105	agc Ser	gga Gly	gct Ala	-gac Asp	gca Ala 110	ctg Leu	aag Lys	577
ctt Leu	gag Glu	ggc Gly 115	Gly	gaa Glu	ggc Gly	gtg Val	ttt Phe 120	gaa Glu	tcc Ser	att Ile	cgc Arg	gca Ala 125	ttg Leu	acg Thr	ctt Leu	625
gga Gly	ggc Gly 130	Ile	cca Pro	gta Val	gtc Val	agt Ser 135	cac His	tta Leu	ggt Gly	ttg Leu	aca Thr 140	ccg Pro	-cag -Gln	tca Ser	gtc	673
ggc Gly 145	gta Val	ctg Leu	ggc Gly	ggc Gly	tat Tyr 150	Lys	gta Val	cag Gln	ggc	aaa Lys 155	Asp	gaa Glu	caa Gln	ago Ser	gcc Ala 160	7.21
aaa Lys	aaa Lys	tta Lev	ata ı Ile	gaa Glu 165	Asp	agt Ser	ata	aaa Lys	tgc ∙Cys 170	gaa Glu	gaa Glu	gca Ala	G17	a gct / Ala 175	atg Met	7,69
atg Met	ctt Lev	gto Val	cto Lev 180	ı Glı	tgt Cys	gtg Val	Pro	g gca Ala 185	'Glu	etc Leu	aca Thr	gcc Ala	: aaa Lys 190	5 I.F.	gcc Ala	817
gaç	aco Thi	g cta Lei 19	u Se	c ata	a cco	g gtc Val	: att	Gly	ato / Ile	ggç Gly	g gct ; Ala	ggt Gly 205	y Va.	g aaa l Ly:	a gcg s Ala	865
gad Asp	gga Gly 210	y Gl	a gti n Va	t cte	c gtt u Val	tat L Tyr 215	: His	t gat s Asp	: att o Ile	ato Ile	ggc = Gly 220	/ His	c gg s Gl	t gt y Va	t gag 1 Glu	913
aga Arc	g Th	a cc r Pr	t aa o Ly	a tt s Ph	t gta e Vai 230	l Lys	g cas s Gli	a tat n Ty	t acq	23	g Ile	gat e As _l	t ga p Gl	a ac u Th	c atc r Ile 240	-96 1
ga Gl	a ac	a ġc r Al	a at a Il	c ag e Se 24	r Gl	a ta y Ty	t gt r Va	t cad	g gat n Ası 250	o Va	a aga	a ca g Ki:	t cg s Ar	t gc g Al 25	t ttc a Phe 5	1009

cct Pro	gaa Glu	caa Gln	aag Lys 260	cat His	tcc Ser	ttt Phe	caa Gln	atg Met 265	aac Asn	cag Gln	aca Thr	gtg Val	ctt Leu 270	gac Asp	ggc , Gly	1057
ttg Leu	tac Tyr	ggg Gly 275	gga Gly	aaa Lýs	taaq	g ato Med	g aga	a cad g Gl: 280	ı Ile	act Thi	t gat	t att	t tca e Sea 28!	r Gli	g ctg n Leu	1106
aaa Lys	gaa Glu	gcc Ala 290	ata Ile	aaa Lys	caa Gln	tac Tyr	cat His 295	tca Ser	gag Glu	ggc Gly	aag Lys	tca Ser 300	atc Ile	gga Gly	ttt Phe	1154
gtt Val	ccg Pro 305	acg Thr	atg Met	ggg Gly	ttt Phe	ctg Leu 310	cat His	gag Glu	ggg Gly	cat His	tta Leu 315	acc Thr	tta Leu	gca Ala	gac Asp	1202
aaa Lys 320	gca Ala	aga Arg	caa Gln	gaa Glu	aac Asn 325	gac Asp	gcc Ala	gtt Val	att Ile	atg Met 330	agt Ser	att Ile	ttt Phe	gtg Val	aat Asn 335	1250
cct Pro	gca Ala	caa Gln	ttc Phe	ggc Gly 340	cct Pro	aat Asn	gaa Glu	gat Asp	ttt Phe 345	gaa Glu	gca Ala	tat Tyr	ccg Pro	cgc Arg 350	gat Asp	1298
att Ile	gag Glu	cgg Arg	gat Asp 355	gca Ala	gct Ala	ctt Leu	gca Ala	gaa Glu 360	aac Asn	gcc Ala	gga Gly	gtc Val	gat Asp 365	att Ile	ctt Leu	1346
ttt Phe	acg Thr	cca Pro 370	gat Asp	gct Ala	cat His	gat Asp	atg Met 375	tat Tyr	ccc Pro	ggt Gly	gaa Glu	aag Lys 380	aat Asn	gtc Val	acg Thr	1394
att Ile	cat His 385	gta Val	gaa Glu	aga Arg	cgc Arg	aca Thr 390	gac Asp	gtg Val	tta Leu	tgc Cys	ggg Gly 395	cgc Arg	tca Ser	aga Arg	gaa Glu	1442
gga Gly 400	cat His	ttt Phe	gac Asp	ggg Gly	gtc Val 405	gcg Ala	atc Ile	gta Val	ctg Leu	acg Thr 410	aag Lys	ctt Leu	ttc Phe	aat Asn	cta Leu 415	1490
gtc Val	aag Lys	ccg Pro	act Thr	cgt Arg 420	gcc Ala	tat Tyr	ttc Phe	ggt Gly	tta Leu 425	aaa Lys	gat Asp	gcg Ala	cag Gln	cag Gln 430	gta Val	1538
gct Ala	gtt Val	gtt Val	gat Asp 435	ggg Gly	tta Leu	atc Ile	agc Ser	gac Asp 440	ttc Phe	ttc Phe	atg Met	gat Asp	att Ile 445	gaa Glu	ttg Leu	1586
gtt Val	cct Pro	gtc Val 450	gat Asp	acg Thr	gtc Val	aga Arg	gag Glu 455	Glu	gac Asp	ggc Gly	tta Leu	gcc Ala 460	aaa Lys	agc Ser	tct Ser	1634
cgc Arg	aat Asn 465	gta Val	tac Tyr	tta Leu	aca Thr	gct Ala 470	gag Glu	gaa Glu	aga Arg	aaa Lys	gaa Glu 475	gcg Ala	cct Pro	aag Lys	ctg Leu	1682
tat Tyr	cgg Arg	gcc Ala	ctt Leu	caa Gln	aca Thr	agt Ser	gcg Ala	gaa Glu	ctt Leu	gtc Val	caa Gln	gcc Ala	ggt Gly	gaa Glu	aga Arg	1730

480	485	490	495
gat cct gaa gcg Asp Pro Glu Ala	gtg ata aaa gct go Val Ile Lys Ala A 500	ca aaa gat atc att gaa la Lys Asp Ile Ile Glu 505	acg act 1778 Thr Thr 510
agc gga acc ata Ser Gly Thr Ile 515	Asp Tyr Val Glu L	tt tat tcc tat ccg gaa eu Tyr Ser Tyr Pro Glu, 20 525	ctc gag 1826 Leu Glu
cct gtg aat gaa Pro Val Asn Glu 530	att gct gga aag a Ile Ala Gly Lys M 535	tg att ctc gct gtt gca et Ile Leu Ala Val Ala 540	gtt-gct 1874 Val Ala
ttt tca aaa gcg Phe Ser Lys Ala 545	cgt tta ata gat a Arg Leu Ile Asp A 550	at atc att att gat att sn Ile Ile Ile Asp Ile 555	cga gaa 1922 Arg Glu
atg gag aga ata Met Glu Arg Ile 560	taat atg tat cga Met Tyr Arg 565	aca atg atg agc ggc aaa Thr Met Met Ser Gly Lys 570	ctt cac 1971 Leu His
agg gca act gtt Arg Ala Thr Val 575	acg gaa gca aac c Thr Glu Ala Asn L 580	etg aac tat gtg gga agc Leu Asn Tyr Val Gly Ser 585	att aca 2019 Ile Thr . 590
att gat gaa gat Ile Asp Glu Asp	ctc att gat gct'g Leu Ile Asp Ala V 595	gtg gga atg ott oot aat Val Gly Met Leu Pro Asn 600	gaa aaa 2067 Glu Lys 605
gta caa att gtg Val Gln Ile Val 610	. Asn Asn Asn Asn G	gga gca cgt ctt gaa acg Gly Ala Arg Leu Glu Thr 615	tat att 2115 Tyr Ile
att cct ggt aaa Ile Pro Gly Lys 625	a cgg gga agc ggc g s Arg Gly Ser Gly V 630	gtc ata tgc tta aac ggt Val Ile Cys Leu Asn Gly 635	gca gcc 2163 Ala Ala
gca cgc ctt gtc Ala Arg Leu Va 640	g cag gaa gga gat a l Gln Glu Gly Asp l 645	aag gtc att att att tcc Lys Val Ile Ile Ser 650	tac aaa 2211 Tyr Lys
atg atg tct ga Met Met Ser As 655	t caa gaa gcg gca a p Gln Glu Ala Ala : 660	agc cat gag ccg aaa gtg Ser His Glu Pro Lys Val 665	gct gtt 2259 Ala Val 670
ctg aat gat ca Leu Asn Asp Gl	a aac aaa att gaa d n Asn Lys Ile Glu (675	caa atg ctg ggg aac gaa Gln Met Leu Gly Asn Glu 680	cca gcc 2307 Pro Ala 685
cgt aca att tt Arg Thr Ile Le 69	u	cttta tcgggggttt tctttt	aaga tttt 2363

<210> 60 <211> 293 <212> PRT <213> Bacillus subtilis

<400> 60

Met Ser Ile Ala Val Ser Glu Glu Glu Ala Lys Ala Val Glu Gly Leu
1 5 10 15

Asn Asp Tyr Leu Ser Val Glu Glu Val Glu Thr Ile Tyr Ile Pro Leu 20 25 30

Val Arg Leu Leu His Leu His Val Lys Ser Ala Ala Glu Arg Asn Lys 35 40 45

His Val Asn Val Phe Leu Lys His Pro His Ser Ala Lys Ile Pro Phe 50 55 60

Ile Ile Gly Ile Ala Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala 65 70 75 80

Årg Ile Leu Gln Lys Leu Leu Ser Arg Leu Pro Asp Arg Pro Lys Val 85 90 95

Ser Leu Ile Thr Thr Asp Gly Phe Leu Phe Pro Thr Ala Glu Leu Lys
100 105 110

Lys Lys Asn Met Met Ser Arg Lys Gly Phe Pro Glu Ser Tyr Asp Val 115 120 125

Lys Ala Leu Leu Glu Phe Leu Asn Asp Leu Lys Ser Gly Lys Asp Ser 130 135 140

Val Lys Ala Pro Val Tyr Ser His Leu Thr Tyr Asp Arg Glu Glu Gly 145 150 155 160

Val Phe Glu Val Glu Gln Ala Asp Ile Val Ile Ile Glu Gly Ile 165 170 175

Asn Val Leu Gln Ser Pro Thr Leu Glu Asp Asp Arg Glu Asn Pro Arg 180 185 190

Ile Phe Val Ser Asp Phe Phe Asp Phe Ser Ile Tyr Val Asp Ala Glu 195 200 205

Glu Ser Arg Ile Phe Thr Trp Tyr Leu Glu Arg Phe Arg Leu Leu Arg 210 215 220

Glu Thr Ala Phe Gln Asn Pro Asp Ser Tyr Phe His Lys Phe Lys Asp 225 230 235 240

Leu Ser Asp Gln Glu Ala Asp Glu Met Ala Ala Ser Ile Trp Glu Ser 245 250 255

Val Asn Arg Pro Asn Leu Tyr Glu Asn Ile Leu Pro Thr Lys Phe Arg 260 265 270

Ser Asp Leu Ile Leu Arg Lys Gly Asp Gly His Lys Val Glu Glu Val 275 280 285

Leu Val Arg Arg Val

290

<210> 61 <211> 281 <212> PRT <213> Bacillus subtilis

<400> 61
Met Glu Gly Leu Asn Asp Tyr Leu Ser Val Glu Glu Val Glu Thr Ile
1 5 10 15

Tyr Ile Pro Leu Val Arg Leu Leu His Leu His Val Lys Ser Ala Ala 20 25 30

Glu Arg Asn Lys His Val Asn Val Phe Leu Lys His Pro His Ser Ala 35 40 45

Lys Ile Pro Phe Ile Ile Gly Ile Ala Gly Ser Val Ala Val Gly Lys
50 55 60

Ser Thr Thr Ala Arg Ile Leu Gln Lys Leu Leu Ser Arg Leu Pro Asp 65 70 75 80

Arg Pro Lys Val Ser Leu Ile Thr Thr Asp Gly Phe Leu Phe Pro Thr 85 90 95

Ala Glu Leu Lys Lys Lys Asn Met Met Ser Arg Lys Gly Phe Pro Glu 100 105 110

Ser Tyr Asp Val Lys Ala Leu Leu Glu Phe Leu Asn Asp Leu Lys Ser 115 120 125

Gly Lys Asp Ser Val Lys Ala Pro Val Tyr Ser His Leu Thr Tyr Asp 130 135 140

Arg Glu Glu Gly Val Phe Glu Val Val Glu Gln Ala Asp Ile Val Ile 145 150 155 160

Ile Glu Gly Ile Asn Val Leu Gln Ser Pro Thr Leu Glu Asp Asp Arg 165 170 175

Glu Asn Pro Arg Ile Phe Val Ser Asp Phe Phe Asp Phe Ser Ile Tyr 180 185 190

Val Asp Ala Glu Glu Ser Arg Ile Phe Thr Trp Tyr Leu Glu Arg Phe 195 200 205

Arg Leu Leu Arg Glu Thr Ala Phe Gln Asn Pro Asp Ser Tyr Phe His 210 215 220

Lys Phe Lys Asp Leu Ser Asp Gln Glu Ala Asp Glu Met Ala Ala Ser 225 230 235 240

Ile Trp Glu Ser Val Asn Arg Pro Asn Leu Tyr Glu Asn Ile Leu Pro
245 250 255

Thr Lys Phe Arg Ser Asp Leu Ile Leu Arg Lys Gly Asp Gly His Lys

- 61 -

265

270

260

Val Glu Glu Val Leu Val Arg Arg Val 275 '<210> 62 <211> 1092 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(1089) <400> 62 atg act aaa caa aca att cgc gtt gaa ttg aca tca aca aaa aaa ccg 48 Met Thr Lys Gln Thr Ile Arg Val Glu Leu Thr Ser Thr Lys Lys Pro aaa cca gac cca aat cag ctt tcg ttc gga aga gtg ttt aca gac cac 96 Lys Pro Asp Pro Asn Gln Leu Ser Phe Gly Arg Val Phe Thr Asp His atg ttt gta atg gac tat gcc gca gat aaa ggt tgg tac gat cca aga 144 Met Phe Val Met Asp Tyr Ala Ala Asp Lys Gly Trp Tyr Asp Pro Arg atc att cct tat caa ccc tta tca atg gat cca act gca atg gtc tat 192 Ile Ile Pro Tyr Gln Pro Leu Ser Met Asp Pro Thr Ala Met Val Tyr cac tac ggc caa acc gtg ttt gaa ggg tta aag gct tac gtg tca gag 240 His Tyr Gly Gln Thr Val Phe Glu Gly Leu Lys Ala Tyr Val Ser Glu 70 gat gac cat gtt ctg ctt ttc aga ccg gaa aaa aat atg gaa cgc ctg 288 Asp Asp His Val Leu Leu Phe Arg Pro Glu Lys Asn Met Glu Arg Leu 90 aat caa tca aac gac cgc ctc tgc atc ccg caa att gat gaa gaa cag 336 Asn Gln Ser Asn Asp Arg Leu Cys Ile Pro Gln Ile Asp Glu Glu Gln 105 gtt ctt gaa ggc tta aag cag ctt gtc gca att gat aaa gac tgg att 384 Val Leu Glu Gly Leu Lys Gln Leu Val Ala Ile Asp Lys Asp Trp Ile 120 cca aat gcg gag ggc acg tcc ctt tac atc cgt ccg ttc atc atc gca 432 Pro Asn Ala Glu Gly Thr Ser Leu Tyr Ile Arg Pro Phe Ile Ile Ala 135 acc gag cct ttc ctt ggt gtt gcg gca tct cat acg tat aag ctc ttg 480 Thr Glu Pro Phe Leu Gly Val Ala Ala Ser His Thr Tyr Lys Leu Leu 150 155 atc att ctt tct ccg gtc ggc tct tat tac aaa gaa ggc att aag ccg 528

Ile Ile Leu Ser Pro Val Gly Ser Tyr Tyr Lys Glu Gly Ile Lys Pro

010177282 I >

- 62 -

				165					170					175		•
gtc Val	aaa Lys	atc Ile	gct Ala 180	gtt Val	gaa Glu	agt Ser	gaa Glu	ttt Phe 185	gtc Val	cgt Arg	gcg Ala	gta Val	aaa Lys 190	ggc Gly	gga Gly	576 '
aca Thr	gga Gly	aat Asn 195	gcc Ala	aaa Lys	acc Thr	gca Ala	gga Gly 200	aac Asn	tat Tyr	gct Ala	tca Ser	agc Ser 205	tta Leu	aaa Lys	gcg Ala	624
cag Gln	cag Gln 210	gta Val	gcc Ala	gaa Glu	gag Glu	aaa Lys 215	gga Gly	ttt Phe	tct Ser	caa Gln	gta Val 220	ctc Leu	tgg Trp	ctg Leu	gac Asp	672
ggc Gly 225	att Ile	gag Glu	aag Lys	aaa Lys	tac Tyr 230	atc Ile	gaa Glu	gaa Glu	gtc Val	gga Gly 235	.agc 'Ser	atg Met	aac Asn	atc Ile	ttc Phe 240	720
ttc Phe	aaa Lys	atc Ile	aac Asn	ggt Gly 245	gaa Glu	atc Ile	gta Val	aca Thr	ccg Pro 250	atg Met	ctg Leu	aac Asn	ggg Gly	agc Ser 255	atc Ile	768
Leu	Glu	Gly	11e 260	acg Thr	Arg	Asn	Ser	Val 265	Ile	Ala	Leu	Leu	Lys 270	His	Trp	816
Gly	Leu	Gln 275	Val	tca Ser	Glu	Arg	Lys 280	Ile	Ala	Ile	Asp	€1u 285	Val	Ile	Gln	864
Ala	His 290	Lys	Asp	@ly ggc	Ile	Leu 295	Glu	Glu	Ala	Phe	Gly 300	Thr	Gly	Thr	Ala	912
Ala 305	Val	Ile	Ser	cca Pro	Val 310	Gly	Glu	Leu	Ile	Trp 315	Gln	Asp	-Glu	Thr	Leu 320	960
Ser	Ile	Asn	Asn	325	Glu	Thr	Gly	Glu	330	Ala	Lys	Lys	Leu	Tyr 335	Asp	1008
Thr	Ile	Thr	G1y 340	/ Ile	Gln	Lys	Gly	Ala 345	Val	. Ala	Asp	Glu	tto Phe 350	Gly	tgg Trp	1056
acg Thr	acc Thr	gaa Glu 355	ı Val	gca Ala	gcg Ala	cto Leu	act Thr 360	Glu	ago Ser	: aag	j taa	1				1092

<210> 63 <211> 363 <212> PRT

<213> Bacillus subtilis

<400> 63 Met Thr Lys Gln Thr Ile Arg Val Glu Leu Thr Ser Thr Lys Lys Pro

- 63 -

1				5					10					15	
Lys	Pro	Asp	Pro 20	Asn	Gln	Leu	Ser	Phe 25	Gly	Arg	Vaļ	Phe	Thr 30	Asp	His
Met	Phe	Val 35	Met	Asp	Tyr	Ala	Ala 40	Asp	Lys	Gly	Trp	Tyr 45	Asp	Pro	Arg
Ile	Ile 50	Pro	Tyr	Gln	Pro	Leu 55	Ser	Met	Asp	Pro	Thr 60	Ala	Met,	Val	Tyr
His 65	Tyr	Gly	Gln	Thr	Val 70	Phe	Glu	Gly	Leu	Lys 75	Ala	Tyr	Val	Ser	Glu 80
Asp	Asp	His	Val	Leu 85	Leu	Phe	Arg	Pro	Glu 90	Lys	Asn	Met	Glu	Arg 95	Leu
Asn	Gln	Ser	Asn 100	Asp	Arg	Leu	Cys	Ile 105	Pro	Gln	Ile	Asp	Glu 110	Glu	Gln
Val	Leu	Glu 115	Gly	Leu	Lys	Gln	Leu 120	Val	Ala	Ile	Asp	Lys 125	Asp	Trp	Ile
' Pro	Asn 130	Ala	Glu	Gly	Thr	Ser 135	Leu	Tyr	Ile	Arg	Pro 140	Phe	Ile	Ile	Ala
Thr 145	Glu	Pro	Phe	Leu	Gly 150	Val	Ala	Ala	Ser	His 155	Thr	Tyr	Гуs	Leu	Leu 160
Ile	Ile	Leu	Ser	Pro 165	Val	Gly	Ser	Tyr	Tyr 170	Lys	Glu	Gly	Ile	Lys 175	Pro
Val	Lys	Ile	Ala 180	Val	Glu	Ser	Glu	Phe 185	Val	Arg	Ala	Val	Lys 190	Gly	Gly
Thr	Gly	Asn 195	Ala	Lys	Thr	Ala	Gly 200	Asn	Tyr	Ala	Ser	Ser 205		Lys	Ala
Gln	Gln 210	Val	Ala	Glu	Glu	Lys 215	_	Phe	Ser	Gln	Val 220	Leu	Trp	Leu	Asp
Gly 225		Glu	Lys	Lys	Tyr 230	Ile	Glu	Glu	Val	Gly 235		Met	Asn	Ile	Phe 240
Phe	Lys	Ile	Asn	Gly 245	Glu	Ile	Val	Thr	Pro 250	Met	Leu	Asn	Gly	Ser 255	
Leu	Glu	Gly	11e 260	Thr	Arg	Asn	Ser	Val 265		Ala	Leu	Leu	Lys 270		Trp
Gly	Leu	Gln 275		Ser	Glu	Arg	Lys 280		Ala	Ile	Asp	Glu 285		Ile	Gln
Ala	His 290	Lys	Asp	Gly	Ile	Leu 295		Glu	Ala	Phe	Gly 300		Gly	Thr	Ala
Ala 305		Ile	Ser	Pro	Val 310	_	Glu	Leu	Ile	Trp 315		Asp	Glu	Thr	Leu 320

Ser Ile Asn Asn Gly Glu Thr Gly Glu Ile Ala Lys Lys Leu Tyr Asp 325 330 335

Thr Ile Thr Gly Ile Gln Lys Gly Ala Val Ala Asp Glu Phe Gly Trp 340 345 350

Thr Thr Glu Val Ala Ala Leu Thr Glu Ser Lys 355 360

<210> 64

<211> 1071

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS

<222> (1)..(1068)

<400> 64

ttg aat aag ctt att gaa cga gaa aaa act gta tat tat aag gaa aag 48
Met Asn Lys Leu Ile Glu Arg Glu Lys Thr Val Tyr Tyr Lys Glu Lys
1 5 10 15

ccc gac ccg tct tcc ttg ggg ttt gga caa tat ttt aca gat tat atg 96 Pro Asp Pro Ser Ser Leu Gly Phe Gly Gln Tyr Phe Thr Asp Tyr Met 20 25 30

ttt gtg atg gac tac gaa gag ggg att gga tgg cat cat ccg aga att. 140
Phe Val Met Asp Tyr Glu Glu Gly Ile Gly Trp His His Pro Arg Ile
35 40 45

gcg ccg tac gca ccg ctt acg ctt gat ccg tct tca tct gtt ttt cat 192
Ala Pro Tyr Ala Pro Leu Thr Leu Asp Pro Ser Ser Ser Val Phe His
50 55 60

tac ggc cag gct gtt ttt gaa gga tta aaa gca tac aga aca gac gac 24 Tyr Gly Gln Ala Val Phe Glu Gly Leu Lys Ala Tyr Arg Thr Asp Asp 65 70 75 80

ggc agg gtg ctg ctg ttc cgt ccg gat caa aat atc aaa cgg ctg aac 288 Gly Arg Val Leu Leu Phe Arg Pro Asp Gln Asn Ile Lys Arg Leu Asn 85 90 95

aga tcg tgt gag cgc atg agc atg ccc cct tta gac gaa gag ctg gtg 336 Arg Ser Cys Glu Arg Met Ser Met Pro Pro Leu Asp Glu Glu Leu Val 100 105 110

ctt gag gca ttg acg caa tta gtt gag ctg gag aaa gat tgg gtt cca 384 Leu Glu Ala Leu Thr Gln Leu Val Glu Leu Glu Lys Asp Trp Val Pro 115 120 125

aag gaa aaa gga acg tca ctg tat att cgt cct ttt gtc att gcc aca 432 Lys Glu Lys Gly Thr Ser Leu Tyr Ile Arg Pro Phe Val Ile Ala Thr 130 135 140

qua ccq agt ctc ggt gtg aug gca tcc agg agc tat aca ttt atg atc 480

	31u 145	Pro	Ser	Leu	Gly	Val 150	Lys	Ala	Ser	Arg	Ser 155	Tyr	Thr	Phe	Met	11e 160	
,	gtg /al	ctt Leu	tcg Ser	cct Pro	gtc Val 165	ggc Gly	tcc Ser	tat Tyr	tat Tyr	ggc Gly 170	gac Asp	gat Asp	cag Gln	ctg Leu	aag Lys 175	ccg Pro	528
į	gtt /al	aga Arg	atc Ile	tat Tyr 180	gtc Val	gaa Glu	gat Asp	gag Glu	tat Tyr 185	gtg Val	agg Arg	gcg Ala	gtc Val	aac Asn 190'	Gly	gga Gly	576
Ş	jtc /al	ggg Gly	ttt Phe 195	gca Ala	aaa Lys	acg Thr	gct Ala	gga Gly 200	aac Asn	tat Tyr	gcc Ala	gcc Ala	agt Ser 205	ctt Leu	cag Gln	gca Ala	624
Ċ	ag Sln	cgg Arg 210	aaa Lys	gcg Ala	aat Asn	gaa Glu	ctg Leu 215	ggc Gly	tat Tyr	gac Asp	cag Gln	gta Val 220	ctg Leu	tgg Trp	ctg Leu	gac Asp	672
P	occ Ala 225	atc Ile	gaa Glu	aag Lys	aaa Lys	tat Tyr 230	gtg Val	gaa Glu	gaa Glu	gta Val	ggg Gly 235	agc Ser	atg Met	aac Asn	atc Ile	ttt Phe 240	720
' t	tc he	gtc Val	ata Ile	aac Asn	ggg Gly 245	gaa Glu	gct Ala	gtc Val	aca Thr	cct Pro 250	gct Ala	tta Leu	agc Ser	gga Gly	agc Ser 255	att Ile	768
ţ	ta Leu	agc Ser	ggg Gly	gtt Val 260	aca Thr	cgt Arg	gcg Ala	tct Ser	gcg Ala 265	att Ile	gaa Glu	ttg Leu	att Ile	cga Arg 270	agc Ser	tgg Trp	816
Ç	gc Hy	att Ile	ccg Pro 275	gtt Val	cgt Arg	gaa Glu	gag Glu	aga Arg 280	ata Ile	tcg Ser	att Ile	gat Asp	gag Glu 285	gtg Val	tat Tyr	gcg Ala	864
P	jcc Ala	tct Ser 290	gca Ala	cgc Arg	gga Gly	gaa Glu	ttg Leu 295	aca Thr	gag Glu	gtc Val	ttt Phe	ggc Gly 300	aca Thr	ggc Gly	acg Thr	gca Ala	912
F	gca Ala 805	gtc Val	gtt Val	acg Thr	cct Pro	gtc Val 310	ggt Gly	gaa Glu	ctc Leu	aac Asn	atc Ile 315	cat His	gga Gly	aaa Lys	acg Thr	gtg Val 320	960
]	le	gta Val	ggc Gly	gac Asp	ggg Gly 325	caa Gln	atc Ile	ggg Gly	gac Asp	ctc Leu 330	tcg Ser	aaa Lys	aag Lys	ctg Leu	tat Tyr 335	gaa Glu	1008
a	cg hr	ata Ile	aca Thr	gat Asp 340	att Ile	cag Gln	ctt Leu	ggc Gly	aag Lys 345	gta Val	aaa Lys	ggc Gly	ccg Pro	ttt Phe 350	aac Asn	tgg Trp	1056
			gaa Glu 355		tga												1071

<210> 65 <211> 356

012177282 1 5

WICHAULT WIN

<212> PRT <213> Bacillus subtilis

<400> 65
Met Asn Lys Leu Ile Glu Arg Glu Lys Thr Val Tyr Tyr Lys Glu Lys
1 5 10 15

Pro Asp Pro Ser Ser Leu Gly Phe Gly Gln Tyr Phe Thr Asp Tyr Met 20 25 30

Phe Val Met Asp Tyr Glu Glu Gly Ile Gly Trp His His Pro Arg Ile
35 40 45

Ala Pro Tyr Ala Pro Leu Thr Leu Asp Pro Ser Ser Ser Val Phe His 50 55 60

Tyr Gly Gln Ala Val Phe Glu Gly Leu Lys Ala Tyr Arg Thr Asp Asp 65 70 75 80

Gly Arg Val Leu Leu Phe Arg Pro Asp Gln Asn Iie Lys Arg Leu Asn 85 90 95

Arg Ser Cys Glu Arg Met Ser Met Pro Pro Leu Asp Glu Glu Leu Val 100 105 110

Leu Glu Ala Leu Thr Gln Leu Val Glu Leu Glu Lys Asp Trp Val Pro 115 120 125

Lys Glu Lys Gly Thr Ser Leu Tyr Ile Arg Pro Phe Val Ile Ala Thr 130 135 140

Glu Pro Ser Leu Gly Val Lys Ala Ser Arg Ser Tyr Thr Phe Met Ile 145 150 155 160

Val Leu Ser Pro Val Gly Ser Tyr Tyr Gly Asp Asp Gln Leu Lys Pro 165 170 175

Val Arg Ile Tyr Val Glu Asp Glu Tyr Val Arg Ala Val Asn Gly Gly 180 185 190

Val Gly Phe Ala Lys Thr Ala Gly Asn Tyr Ala Ala Ser Leu Gln Ala 195 200 205

Gln Arg Lys Ala Asn Glu Leu Gly Tyr Asp Gln Val Leu Trp Leu Asp 210 215 220

Ala Ile Glu Lys Lys Tyr Val Glu Glu Val Gly Ser Met Asn Ile Phe 225 230 235 240

Phe Val Ile Asn Gly Glu Ala Val Thr Pro Ala Leu Ser Gly Ser Ile 245 250 255

Leu Ser Gly Val Thr Arg Ala Ser Ala Ile Glu Leu Ile Arg Ser Trp 260 265 270

Gly Ile Pro Val Arg Glu Glu Arg Ile Ser Ile Asp Glu Val Tyr Ala 275 280 .285

.

Ala Ser Ala Arg Gly Glu Leu Thr Glu Val Phe Gly Thr Gly Thr Ala 290 Ala Val Val Thr Pro Val Gly Glu Leu Asn Ile His Gly Lys Thr Val 'Ile Val Gly Asp Gly Gln Ile Gly Asp Leu Ser Lys Lys Leu Tyr Glu Thr Ile Thr Asp Ile Gln Leu Gly Lys Val Lys Gly Pro Phe Asn Trp Thr Val Glu Val 355 <210> 66 <211> 1428 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(1425) <400> 66 atg tta aac ggc caa aaa gaa tat cgc gtg gaa aaa gac ttc ctt ggg Met Leu Asn Gly Gln Lys Glu Tyr Arg Val Glu Lys Asp Phe Leu Gly gaa aaa caa att gaa gca gat gtt tat tac gga att cag acg ctc cgt 96 Glu Lys Gln Ile Glu Ala Asp Val Tyr Tyr Gly Ile Gln Thr Leu Arg gct tct gaa aat ttt ccg atc aca gga tac aaa atc cat gag gaa atg 144 Ala Ser Glu Asn Phe Pro Ile Thr Gly Tyr Lys Ile His Glu Glu Met 40 att aac gca ctg gcg att gtg aaa aaa gct gcg gct ctt gcc aac atg 192 Ile Asn Ala Leu Ala Ile Val Lys Lys Ala Ala Leu Ala Asn Met 55 gac gtg aaa cgg ctg tat gaa gga att ggc caa gct atc gta caa gcc 240 Asp Val Lys Arg Leu Tyr Glu Gly Ile Gly Gln Ala Ile Val Gln Ala gct gac gag att ctg gaa ggc aag tgg cac gat cag ttt atc gtc gat Ala Asp Glu Ile Leu Glu Gly Lys Trp His Asp Gln Phe Ile Val Asp ccg att cag ggc ggt gcc gga act tct atg aac atg aac gcg aat gag Pro Ile Gln Gly Gly Ala Gly Thr Ser Met Asn Met Asn Ala Asn Glu gtt atc gga aac cgg gcg ctt gaa atc atg gga cat aaa aag gga gat Val Ile Gly Asn Arg Ala Leu Glu Ile Met Gly His Lys Lys Gly Asp 120

tat Tyr	atc Ile 130	cat His	tta Leu	agt Ser	Pro	aac Asn 135	aca Thr	cat His	gtg Val	aac Asn	atg Met 140	tca 4 Ser 4	cag · Gln	tct Ser	cag Gln	432
aac Asn 145	gat Asp	gtg Val	ttc Phe	ccg Pro	act Thr 150	gct Ala	atc Ile	cat His	att Ile	tcc Ser 155	aca Thr	ttg Leu	aag Lys	ctc Leu	tta Leu 160	480
gaa Glu	aaa Lys	ctg Leu	ctg Leu	aaa Lys 165	aca Thr	atg Met	gaa Glu	gat Asp	atg Met 170	cat His	agt Ser	gtg Val	ttt Phe	aaa Lys 175	caa Gln	528 .
aaa Lys	gca Ala	cag Gln	gag Glu 180	ttt Phe	cac His	tct Ser	gtt Val	att Ile 185	aaa Lys	atg Met	ggc	cgg Arg	aca Thr 190	cac His	ctt Leu	576
caa Gln	gat Asp	gcg Ala 195	Val	ccg Pro	atc Ile	cgt Arg	ctt Leu 200	ggc Gly	cag Gln	gaa Glu	ttc Phe	gaa Glu 205	gct Ala	tac Tyr	agc Ser	624
cgt Arg	gtt Val 210	Leu	gag Glu	cgt Arg	gat Asp	atc Ile 215	aaa Lys	cga Arg	atc Ile	aag Lys	caa Gln 220	teg Ser	.cgc Arg	cag Gln	.cac His	· 6 72
ctg Leu 225	Туг	gaa Glu	gtc Val	aac Asn	atg Met 230	ggc	gca Ala	act Thr	gct Ala	gtt Val 235	ggt Gly	aca Thr	Gly	ctg Leu	aac Asn 240	720
gct Ala	gat Asp	cct Pro	gaa Glu	tat Tyr 245	Ile	aaa Lys	cag Gln	gta Val	gta Val 250	Lys	cac His	ctt Leu	gct Ala	gat Asp 255	116	768
ago Sei	gg Gl	g cti y Lei	cct Pro 260	Leu	gtc Val	ggc Gly	gct Ala	gat Asp 265	His	ctt Leu	gtt Val	gat Asp	gcg Ala 270	Thr	caa Gln	.816
aat Ası	ace n Th	a gat r Ası 27	p Ala	c tat a Tyr	aca Thr	gag Glu	gta Val 280	Ser	gct Ala	tca Ser	tta Leu	aaa Lys 285	Val	tgc Cys	atg Met	864
at Me	g aa t As 29	n Me	g tc	g aaq r Lys	g ato s Ile	gca Ala 295	Asr	gac Asp	cto Lev	g cgo Aro	tta g Lev 300	ı Met	gcg Ala	te Sei	g gga Gly	912
cc Pr 30	o Ar	c gc g Al	c gg a Gl	a ctt y Lei	gcq Ala 310	a Glu	a att	t-ct e Sei	cto Lei	g cct 2 Pro 31	o Ala	a egt a Arg	: caç Glr	Pro	g ggt 5 Gly 320	960
tc Se	a to r Se	t at	t at e Me	g ccc t Pro 32	o G l	g aaa y Lys	a gto s Val	c aat l Asi	2 CCC 1 Pro 330	o Vai	t ato	g gcg t Ala	g gaq a Glu	g cte Let 33	g atc u Ile 5	1008
aa As	c ca n Gl	a at n Il	t gc e Al 34	a Ph	c -cad e Gl	g gti n Vai	t ate	c gg e G1 34	y Ası	t gad n Asp	c aat p Asi	t aca n Thi	a ate 110 350	е Су	c ctt s Leu	1056
gc Al	t to a Se	a ga er Gl	a go .u Al	c gg a Gl	c ca y Gl	g -ct [.] n -Le	t ga u Gl	g tte	g aa u As	c gt n Va	c at	g gaq t Gl	g ∢cc u Pr	c gt o Va	g ctt l Leu	1104

		355					360					365				
gtc Val	ttt Phe 370	aat Asn	ttg Leu	ctt Leu	caa, Gln	tcc Ser 375	atc Ile	agc Ser	atc Ile	atg Met	aac Asn 380	aac Asn	ggc Gly	ttc Phe	cgt Arg	1152
tcg Ser 385	ttc Phe	act Thr	gac Asp	aac Asn	tgc Cys 390	tta Leu	aaa Lys	ggc Gly	att Ile	gaa Glu 395	gcc Ala	aac Asn	gaa Glu	aag Lys	cgt Arg 400	1200
atg Met	aag Lys	caa Gln	tac Tyr	gta Val 405	gaa Glu	aaa Lys	agc Ser	gca Ala	ggc Gly 410	gtg Val	atc Ile	aca Thr	gct Ala	gtc Val 415	aat Asn	1248
ccg Pro	cat His	ctt Leu	ggg Gly 420	tat Tyr	gaa Glu	gcg Ala	gca Ala	gct Ala 425	aga Arg	att Ile	gcc Ala	agg Arg	gaa Glu 430	gca Ala	att Ile	1296
atg Met	aca Thr	ggg Gly 435	caa Gln	tct Ser	gtc Val	cgg Arg	gat Asp 440	ctt Leu	tgt Cys	ctg Leu	cag Gln	cat His 445	gat Asp	gtg Val	ctg Leu	1344
act Thr	gaa Glu 450	gaa Glu	gaa Glu	ttg Leu	gat Asp	att Ile 455	att Ile	tta Leu	aac Asn	cca Pro	tat Tyr 460	gag Glu	atg Met	acc Thr	aaa Lys	1392
cca Pro 465	ggt Gly	atc Ile	gca Ala	ggg Gly	aaa Lys 470	gaa Glu	cta Leu	tta Leu	gaa Glu	aaa Lys 475	taa					1428
<211 <212	0> 67 L> 47 2> PB B> Ba	75	.us s	subti	llis						·					
)> 67 Leu		Gly	Gln 5	Lys	Glu	Tyr	Arg	Val 10	Glu	Lys	Asp	Phe	Leu 15	Gly	
Glu	Lys	Gln	Ile 20	Glu	Ala	Asp	Val	Tyr 25	Tyr	Gly	Ile	Gln	Thr 30	Leu	Arg	
Ala	Ser	Glu 35	Asn	Phe	Pro	Ile	Thr 40	Gly	Tyr	Lys	Ile	His 45	Glu	Glu	Met	
Ile	Asn 50	Ala	Leu	Ala	Ile	Val 55	Lys	Lys	Ala	Ala	Ala 60	Leu	Ala	Asn	Met	
Asp 65	Val	Lys	Arg	Leu	Tyr 70	Glu	Gly	Ile	Gly	Gln 75	Ala	Ile	Val	Gln	Ala 80	
Ala	Asp	Glu	Ile	Leu 85	Glu	Gly	Lys	Trp	His 90	Asp	Gln	Phe	Ile	Val 95	Asp	
Pro	Ile		Gly 100	Gly	Ala	Gly	Thr	Ser 105	Met	Asn	Met	Asn	Ala 110	Asn	Glu	

1000010- JAMO 010177080 I

Val	Ile	Gly 115	Asn	Arg	Ala	Leu	Glu 120	Ile	Met	Gly	His	Lys 125	Lys	Gly	Asp
Tyr	Ile 130	His	Leu	Ser	Pro	Asn 135	Thr	His	Val	Asn	Met 140	Ser	Gln	Ser	Gln
Asn 145	Asp	Val	Phe	Pro	Thir 150	Ala	Ile	His	Ile	Ser 155	Thr	Leu	Lys	Leu	Leu 160
Glu	Lys	Leu	Leu	Lys 165	Thr	Met	Glu	Asp	Met 170	His	Ser	Val	Phe	Lys 175	Gln
Lys	Ala	Gln	Glu 180	Phe	His	Ser	Val	Ile 185	Łys	Met	Gly	Arg	Thr 190	His	Leu
Gln	Asp	Ala 195	Val	Pro	Ile	Arg	Leu 200	Gly	Gln	Glu	₽he	Glu 205	Ala	Tyr	Ser
Arg	Val 210		Glu	Arg	Asp	Ile 215		Arg	Ile	Lys	Gln 220		Arg	Gln	His
Leu 225	-	Glu	Val	Asn	Met 230	Gly	Ala	Thr	Ala	Val 235		Thr	Gly	Leu	Asn 240
Ala	Asp	Pro	Glu	Tyr 245		Lys	Gln ,	Val	Val 250	Lys	His	Leu	Ala ,	Asp 255	Ile
Ser	Gly	Leu	260		Val	Gly	Ala	Asp 265		Leu	ı Val	Asp	Ala 270	Thr	Gln
Asn	Thr	275		Tyr	Thr	Glu	val 280		Ala	Ser	Leu	289	val	*Cys	Met
Met	290		Ser	Lys	: Ile	Ala 299		Asp	Let	ı Arç	300		: Ala	Ser	Gly
Pro 305	-	g Ala	a Gly	/ Let	310		ı Ile	Sez	: Leu	315	o Ala	Arq	g Gln	Pro	320
Sea	r Se	r Ile	e Met	325		/ Ly:	s Val	Ası	330		l Met	: Ala	a ∙Glu	335	ı Ile
Ası	n Gl	n Il	e Ala 34		e Glr	n Va	l Ile	345		n Ası	p Ası	n Th	r 11e 350		s Lev
Al	a ·Se	r Gl 35		a Gl	y Glr	n Le	u Glu 360		u Ası	n Va	l Me	t G1 36		va:	l Lev
Va	1 Ph 37		n Le	u Le	u Gli	n Se 37		e Se	r Il	e Me	t As: 38:		n Gly	y Pho	e Aro
Se 38		e Th	r As	p As	n Cy: 39		u Ly:	s Gl	y Il	e .G1 39		a As	n Gl	u Ly	400
Ме	t Ly	s Gl	n Ty	r Va 40		u Ly	s Se	r Al	a Gl 41		1 11	e Th	r Al	a Va 41	l Ası 5
Pr	o Hi	s Le	u Gl	у Ту	r Gl	u Al	a Al	a Al	a Ar	g Il	e Al	a Ar	g Gl	u Al	a Ile

- 71 -

420 425 430 Met Thr Gly Gln Ser Val. Arg Asp Leu Cys Leu Gln His Asp Val Leu 440 Thr Glu Glu Glu Leu Asp Ile Ile Leu Asn Pro Tyr Glu Met Thr Lys 455 Pro Gly Ile Ala Gly Lys Glu Leu Leu Glu Lys 470 <210> 68 <211> 768 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(765) <400> 68 atg aaa cga gaa agc aac att caa gtg ctc agc cgt ggt caa aaa gat 48 Met Lys Arg Glu Ser Asn Ile Gln Val Leu Ser Arg Gly Gln Lys Asp 10 cag cct gtg agc cag att tat caa gta tca aca atg act tct cta tta Gln Pro Val Ser Gln Ile Tyr Gln Val Ser Thr Met Thr Ser Leu Leu gac gga gta tat gac gga gat ttt gaa ctg tca gag att ccg aaa tat 144 Asp Gly Val Tyr Asp Gly Asp Phe Glu Leu Ser Glu Ile Pro Lys Tyr 40 gga gac ttc ggt atc gga acc ttt aac aag ctt gac gga gag ctg att 192 Gly Asp Phe Gly Ile Gly Thr Phe Asn Lys Leu Asp Gly Glu Leu Ile ggg ttt gac ggc gaa ttt tac cgt ctt cgc tca gac gga acc gcg aca Gly Phe Asp Gly Glu Phe Tyr Arg Leu Arg Ser Asp Gly Thr Ala Thr ccg gtc caa aat gga gac cgt tca ccg ttc tgt tca ttt acg ttc ttt 288 Pro Val Gln Asn Gly Asp Arg Ser Pro Phe Cys Ser Phe Thr Phe Phe 85 aca ccg gac atg acg cac aaa att gat gcg aaa atg aca cgc gaa gac 336 Thr Pro Asp Met Thr His Lys Ile Asp Ala Lys Met Thr Arg Glu Asp 100 105 ttt gaa aaa gag atc aac agc atg ctg cca agc aga aac tta ttt tat 384 Phe Glu Lys Glu Ile Asn Ser Met Leu Pro Ser Arg Asn Leu Phe Tyr 120 gca att cgc att gac gga ttg ttt aaa aag gtg cag aca aga aca gta 432 Ala Ile Arg Ile Asp Gly Leu Phe Lys Lys Val Gln Thr Arg Thr Val 135 140

gaa Glu 145	ctt Leu	caa Gln	gaa Glu	aaa Lys	ect Pro 150	tac Tyr	gtg Val	cca Pro	atg Met	gtt Val 155	gaa Glu	gcg Ala	gtc Val	aaa Lys	aca Thr 1 6 0	480
cag Gln	ccg Pro	att Ile	ttc Phe	aac Asn 165	ttc Phe	gac Asp	aac Asn	gtg Val	aga Arg 170	gga Gly	acg Thr	att Ile	gta Val	ggt Gly 175	ttc Phe	528
ttg Leu	aca Thr	cca Pro	gct Ala 180	tat Tyr	gca Ala	aac Asn	gga Gly	atc Ile 185	gcc Ala	gtt Val	tct Ser	-ggc Gly	tat Tyr 190	cac His	ctg Leu	576
cac His	ttc Phe	att Ile 195	gac Asp	gaa Glu	gga Gly	ege Arg	aat Asn 200	tca Ser	ggc Gly	gga Gly	cac His	gtt Val 205	ttt Phe	gac Asp	tat Tyr	·624
gtg Val	ctt Leu 210	gag Glu	gat Asp	tgc Cys	acg Thr	gtt Val 215	acg Thr	att Ile	tct Ser	caa Gln	aaa Lys 220	atg Met	aac Asn	atg Met	aat Asn	:67:2
ctc , Leu 225	Arg	ct.t Leu	ccg Pro	aac Asn	aca Thr 230	gcg Ala	gat Asp	ttc Phe	ttt Phe	aat Asn 235	gcg Ala	aat Asn	ctg Leu	gat Asp	aac Asn 240	720
cct Pro	gat Asp	ttt Phe	gcg Ala	aaa Lys 245	Asp	atc	gaa Glu	Thr	act Thr 250	Glu	gga Gly	agc Ser	cct Pro	gaa Glu 255	taa	768
<21 <21	0> 6 1> 2 2> P 3> B	55 RT	lus	subt	ilis											
<21 <21 <21	1> 2 2> P 3> B 0> 6 Lys	55 RT acil			Asn	Ile	Gln	Val	Leu 10		Arg	-Gly	Gln	Lys 15		
<21 <21 <21 <40 Met	1> 2 2> P 3> B 00> 6 Lys	55 RT acil 9 Arg	Glu	Ser 5	Asn				10 Ser	1		Gly Thr		15 Leu		,
<21 <21 <21 <40 Met 1	1> 2 2> P 3> B 00> 6 Lys	55 RT acil 9 Arg	Glu Ser 20	Ser 5 Gln	Asn	Туг	Gln	Val 25 Glu	10 Ser	Thr	Met	Thr	Ser 30	15 Leu		
<21 <21 <21 <40 Met 1 G1r	1> 2 2> P 3> B 00> 6 Lys n Pro	55 RT acil 9 Arg Val Val	Glu Ser 20	Ser 5 Gln	Asn Ile	Tyr Asp	Gln Phe 40	Val 25 Glu	Ser Leu	Thr	Met •Glu	Thr Ile 45	Ser 30 Pro	Leu Lys	Leu	
<21 <21 <40 Met 1 Glr Asr	1> 2 2> P 3> B 00> 6 Lys 1 Pro 50 4 Asp 50 7 Phe	55 RT acil 9 Arg Val Val	Glu Ser 20 Tyr G	Ser 5 Glm Asp	Asn Ile Gly	Tyr Asp Thr 55	Gln Phe 40	Val 25 Glu Asn	Ser Leu Lys	Thr Ser	Met Glu Asp 60	Thr Ile 45	Ser 30 Pro	Leu Lys	Leu Tyr	
<21 <21 <40 Met Glr Glr Gly Gly 65	1> 2 2> P 3> B 00> 6 Lys 1 Pro 5 Gly 7 Asp 50 7 Phe	55 RT acil 9 Arg Val Val 35 Phe	Glu Ser 20 Tyr 6	Ser 5 Glm Asp	Asn Ile Gly Gly Phe 70 Asp	Tyr Asp Thr 55	Gln Phe 40 Phe Arg	Val 25 Glu Asn	Lev Lys	Thr Ser Ser Ser 75	Met Glu Asp 60	Thr Ile 45 Gly	Ser 30 Pro	Leu Lys Leu Ala	Leu Tyr The 80	
<21 <21 <21 <40 Met 1 Glr Asr Gly Gly 69	1> 2 2> P 3> B 00> 6 Lys n Pro O Gly Asp 50 Val	55 RT acil 9 Arg Val 7 Val 35 Phe	Glu Ser 20 Tyr 6 Gly Co Gly	Ser 5 Gln Asp Glv Glv Strh	Asn Ile OGly Coly Asp	Tyr Asp Thr 55 Tyr	Phe 40 Phe Arg	Val 25 Glu Asn Leu	Leu Lys Arc	Thr Ser J Ser 75	Met Glu Asp 60 Asp	Thr Ile 45 Gly Gly	Ser 30 Pro Glu Thr	Leu Lys Leu Ala Phe	Leu Tyr The 80	

010177040 1 -

Ala Ile Arg Ile Asp Gly Leu Phe Lys Lys Val Gln Thr Arg Thr Val +135 Glu Leu Gln Glu Lys Pro Tyr Val Pro Met Val Glu Ala Val Lys Thr 150 Gln Pro Ile Phe Asn Phe Asp Asn Val Arg Gly Thr Ile Val Gly Phe Leu Thr Pro Ala Tyr Ala Asn Gly Ile Ala Val Ser Gly Tyr His Leu 185 His Phe Ile Asp Glu Gly Arg Asn Ser Gly Gly His Val Phe Asp Tyr Val Leu Glu Asp Cys Thr Val Thr Ile Ser Gln Lys Met Asn Met Asn 210 220 Leu Arg Leu Pro Asn Thr Ala Asp Phe Phe Asn Ala Asn Leu Asp Asn 230 Pro Asp Phe Ala Lys Asp Ile Glu Thr Thr Glu Gly Ser Pro Glu 245 250 <210> 70 <211> 1254 <212> DNA <213> Escherichia coli <220> <221> CDS <222> (1)..(1251) . <400> 70 atg aca tto too ott ttt ggt gac aaa ttt acc ogc cac too ggc att 48 Met Thr Phe Ser Leu Phe Gly Asp Lys Phe Thr Arg His Ser Gly Ile 10 acg ctg ttg atg gaa gat ctg aac gac ggt tta cgc acg cct ggc gcg 96 Thr Leu Leu Met Glu Asp Leu Asn Asp Gly Leu Arg Thr Pro Gly Ala att atg ctc ggc ggc ggt aat ccg gcg cag atc ccg gaa atg cag gac Ile Met Leu Gly Gly Gly Asn Pro Ala Gln Ile Pro Glu Met Gln Asp 40 tac ttc cag acg cta ctg acc gac atg ctg gaa agt ggc aaa gcg act 192 Tyr Phe Gln Thr Leu Leu Thr Asp Met Leu Glu Ser Gly Lys Ala Thr 55 gat gca ctg tgt aac tac gac ggt cca cag ggg aaa acg gag cta ctc 240 Asp Ala Leu Cys Asn Tyr Asp Gly Pro Gln Gly Lys Thr Glu Leu Leu aca ctg ctt gcc gga atg ctg cgc gag aag ttg ggt tgg gat atc gaa 288 Thr Leu Leu Ala Gly Met Leu Arg Glu Lys Leu Gly Trp Asp Ile Glu

				85					90					95		
cca Pro	cag Gln	aat Asn	att Ile 100	gca Ala	cta Leu	aca Thr	Asn ·	ggc Gly 105	agc Ser	cag Gln	agc Ser	gcg (Ala (ttt Phe 110	ttc Phe	tac Tyr	336
tta Leu	ttt Phe	aac Asn 115	ctg Leu	ttt Phe	gcc Ala	gga Gly	cgc Arg 120	cgt Arg	gcc Ala	gat Asp	ggt Gly	cgg Arg 125	gtc Val	aaa Lys	aaa Lys	384
gtg Val	ctg Leu 130	ttc Phe	ccg Pro	ctt Leu	gca Ala	ccg Pro 135	gaa Glu	tac Tyr	att Ile	ggc Gly	tat Tyr 140	gct Ala	gac Asp	gcc Ala	gga Gly	432
ctg Leu 145	gaa Glu	gaa Glu	gat Asp	ctg Leu	ttt Phe 150	gtc Val	tct Ser	gcg Ala	cgt Arg	ccg Pro 155	aat Asn	att Ile	gaa Glu	ctg Leu	ctg Leu 160	48.0
ccg Pro	gaa Glu	ggc Gly	cag Gln	ttt Phe 165	aaa Lys	tac Tyr	cac His	gtc Val	gat Asp 170	ttt Phe	gag Glu	cat His	ctg Leu	cat His 175	att Ile	528
ggc Gly	gaa Glu	gaa Glu	acc Thr 180	Gly	atg Met	att Ile	tgc Cys	gtc Val 185	tcc Ser	cgg Arg	-ccg Pro	acg Thr	aat Asn 190	cca Pro	aca Thr	576
ggc Gly	aat Asn	gto Val	Ile	act Thr	gac Asp	gaa Glu	gag Glu 200	ttg Leu	ctg Leu	aag Lys	ctt Leu	igac Asp 205	.gcg Ala	ctg Leu	elà dec	624
aat Asr	cas n Glr 210	Hi	gg¢ Gly	att / Ile	-ccg Pro	ctg Leu 215	Val	att Ile	gat Asp	aac Asn	gct Als 220	Tyr	:G1 y	gtc Val	ccg	672
tto Phe 22	e Pro	g gg	t ato	ato e Ile	tto Phe 230	Ser	gaa Glu	gcg Ala	.cgc	Pro 235	rer.	tgg Trp	aat Asn	ecg Pro	aat Asn 240	720
ate	c gto e Vai	g ct	g tg u Cy	c ato s Met 245	t Ser	ctt Leu	tcc Ser	aag Lys	cto Lev 250	ı Gly	t cta y Lew	a cct 1 Pro	ggc Gly	Ser 259	cgc Arg	7:68
tg Cy	c gg s Gl	c at y Il	t at e Il 26	e Il	c gco e Ala	a aat a Asr	gaa n Glu	a aaa Lys 269	i Ile	e Ile	c acc	c gco r Ala	270	ร เมง	aat Asn	816
at Me	g aa t As	c gg n G1 27	y Il	t ate	c age	c cto	g gca u Ala 280	a Pro	Gl;	gg:	t at	t ggt e Gly 285	y Pro	g gco	g atg a Met	864
at Me	g tg t Cy 29	s Gl	a at .u Me	g at t Il	t aa e Ly	g cg s Are 29	g As	c ga n Ası	t cto	g-ct- u Le	g cg u Ar 30	g Le	g tc u Se	t ga r Gl	a aca u Thr	912
٧a	c at	c aa e Ly	aa co /s Pi	g tt o Ph	t ta e Ty 31	r Ty	c ca r Gl	g cg n Ar	t gt g Va	t ca 1 Gl 31	n GI	a ac u Th	t at r·Il	c gc e Al	c atc a Ile 320	960

att Ile	cgc Arg	cgc Arg	tat Tyr	tta Leu 325	ccg Pro	gaa Glu	aat Asn	cgc Arg	tgc Cys 330	ctg Leu	att Ile	cat His	aaa Lys	ccg Pro 335	gaa Glu	, 1008
gga Gly	gcc Ala	att Ile	ttc Phe 340	ctc Leu	tgg Trp	cta Leu	tgg Trp	ttt Phe 345	aag Lys	gat Asp	ttg Leu	ccc Pro	att Ile 350	acg Thr	acc Thr	1056
aag Lys	cag Gln	ctc Leu 355	tat Tyr	cag Gln	cgc Arg	ctg Leu	aáa Lys 360	gca Ala	cgc Arg	ggc Gly	gtg Val	ctg Leu 365	atg Met	gtg Val	ccg Pro	1104
ggg Gly	cac His 370	aac Asn	ttc Phe	ttc Phe	cca Pro	ggg Gly 375	ctg Leu	gat Asp	aaa Lys	ccg Pro	tgg Trp 380	ccg Pro	cat His	acg Thr	cat His	1152
caa Gln 385	tgt Cys	atg Met	cgc Arg	atg Met	aac Asn 390	tac Tyr	gta Val	cca Pro	gag Glu	ccg Pro 395	gag Glu	aaa Lys	att Ile	gag Glu	gcg Ala 400	1200
ggg Gly	gtg Val	aag Lys	att Ile	ctg Leu 405	gcg Ala	gaa Glu	gag Glu	ata Ile	gaa Glu 410	aga Arg	gcc Ala	tgg Trp	gct Ala	gaa Glu 415	agt Ser	1248
cac His	taa															1254
<210> 71 <211> 417 <212> PRT <213> Escherichia coli																
)/ E:	scnei	cichi	La CC	11											
	?> 71	L				Glv	Δsn	Luc	Phe	ጥኮ~	0.50	u: a	Co.w.	C1	71-	
	?> 71	L				Gly	Asp	Lys	Phe 10	Thr	Arg	His	Ser	Gly 15	Ile	
Met 1 Thr)> 7] Thr Leu	Phe Leu	Ser Met 20	Leu 5 Glu	Phe Asp	Leu	Asn	Asp 25	10 Gly	Leu	Arg	Thr	Pro 30	15 Gly	Ala	
Met 1 Thr)> 7] Thr Leu	Phe Leu	Ser Met 20	Leu 5 Glu Gly	Phe Asp Gly	Leu Asn	Asn Pro	Asp 25 Ala	10 Gly Gln	Leu Ile		Thr Glu	Pro 30 Met	15 Gly	Ala	
Met l Thr	P> 71 Thr Leu Met	Phe Leu Leu 35	Ser Met 20	Leu 5 Glu Gly	Phe Asp Gly	Leu Asn	Asn Pro 40	Asp 25 Ala	Gly	Leu Ile	Arg Pro	Thr Glu 45	Pro 30 Met	15 Gly Gln	Ala	
Met 1 Thr Ile Tyr Asp 65	D> 71 Thr Leu Met Phe 50	Phe Leu 35 Gln Leu	Ser Met 20 Gly Thr	Leu 5 Glu Gly Leu Asn	Phe Asp Gly Leu Tyr 70	Leu Asn Thr 55 Asp	Asn Pro 40 Asp Gly	Asp 25 Ala Met	Gly Gln Leu Gln	Leu Ile Glu Gly 75	Arg Pro Ser 60 Lys	Thr Glu 45 Gly Thr	Pro 30 Met Lys Glu	Gly Gln Ala Leu	Ala Asp Thr Leu 80	
Met 1 Thr Ile Tyr Asp 65	D> 71 Thr Leu Met Phe 50	Phe Leu 35 Gln Leu	Ser Met 20 Gly Thr	Leu 5 Glu Gly Leu Asn	Phe Asp Gly Leu Tyr 70	Leu Asn Thr 55 Asp	Asn Pro 40 Asp Gly	Asp 25 Ala Met	Gly Gln Leu Gln	Leu Ile Glu Gly 75	Arg Pro Ser 60	Thr Glu 45 Gly Thr	Pro 30 Met Lys Glu	Gly Gln Ala Leu	Ala Asp Thr Leu 80	

105

Leu Phe Asn Leu Phe Ala Gly Arg Arg Ala Asp Gly Arg Val Lys Lys 115 120 125

Val	Leu 130	Phe	Pro	Leu		Pro 1	Glu	Tyr	Ile	Gly	Tyr :	Ala	Asp	Ala	Gly
Leu 145	Glu	Glu	Asp	Leu	Phe 150	Val	Ser	Ala	Arg	Pro 155	Asn	Ile	Glu	Leu	Leu 160
Pro	Glu	Gly	Gln	Phe 165	Lys	Tyr	His	Val	Asp 170	Phe	Glu	His	Leu	His ,175	Ile
Gly	Glu	Glu	Thr 180	Gly	Met	Ile	Cys	Val 185	Ser	Arg	Pro	Thr	Asn 190	Pro	Thr
Gly	Asn	Val 195	Ile	Thr	Asp	Glu	Glu 200	Leu	Leu	Lys	Leu	Asp 205	Ala	Leu	Gly
Asn	Gln 210		Gly	Ile	Pro	Leu 215	Val	Ile	Asp	Asn	Ala 220	Tyr	Gly	Val	₽ro
Phe 225	Pro	Gly	Ile	Ile	Phe 230	Ser	Glu	Ala	Arg	Pro 235	Leu	Trp	Asn	Pro	Asn 240
				Met 245					250					255	
+			260			,	٠,	265					270	1	Asn
		275	•				280					285	•		Met
	290)				295					300)			Thr
305	5				310					315	•				320
		-		325	5				330	0				33	
			341	0				345	5				35	U	r Thr
		35	5				360)				36	5		l Pro
	37	0				37	5				386	0			r His
38	5				390)				39	5				u Ala 400
Gl	y Va	l Ly	s Il	e Le 40		a Gl	u Gl	u Il	e Gl 41	u Ar O	g Al	a Tr	p Al	a Gl 41	u Ser 5

His

<210> 72 <211> 8803 <212> DNA <213> Artificial Sequence

(213) Withinity Seddelice

<220>

<400> 72

<223> Description of Artificial Sequence: Recombinant pAN294 plasmid

tgcqccqcta caqqqcqcqt ccattcqcca ttcaggctgc gcaactgttg ggaagggcga 60 teggtgeggg cetetteget attaegeeag etggegaaag ggggatgtge tgeaaggega 120 ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagtgaa 180 ttgtaatacg actcactata gggcgaattg ggcccgacgt cgcatgctgg atgaaaagcc 240 gatgaccgct tttcaggtct gtcagcagct ttttcctgct gtatatgaaa aggaattgtt 300 tttaacgatg tcagaaacgg caggtcacct tgatgtgttg gaggctgaag aagccatcac 360 gtcatattgg gaaggaaata ccgtatactt taaaacaatg aagaggtgaa atgggtgaaa 420 catatagcgg gaaaaaggat ttggataacc ggcgcttcag gagggcttgg agaaagaatc 480 gcatacttat gcgcggctga aggagcccat gtcctgctgt cggctagacg cgaggatcgt 540 ttgatagaaa tcaaaaggaa aataaccgag gaatggagcg gacagtgtga gatttttcct 600 ctggatgtcg gccgcctaga ggatatcgcc cgggtccgcg atcagatcgg ctcgattgat 660 gtactgatta acaatgcagg cttcggtata tttgaaacgg ttttagactc tacattggat 720 gacatgaaag cgatgtttga tgtgaatgtc ttcggcctga tcgcctgtac aaaagcggtg 780 cttccgcaaa tgcttgagca aaaaaaggga catatcatca atatcgcctc tcaagcgggg 840 aaaatcgcca caccgaagtc tagcctgtat tccgcgacca aacatgccgt gttaggttac 900 tcaaacqctt tqcqqatqqa qctttcqgga accggcattt atgtgacaac agtcaacccg 960 ggcccgattc agacggactt tttttccatt gctgataaag gcggggacta cgccaaaaat 1020 gtcggccgct ggatgcttga tcctgatgac gtggcagctc aaattacagc tgcaattttt 1080 acgaaaaagc gggagatcaa tcttccgcgt ttaatgaatg ccggcactaa gctgtatcag 1140 ctgtttccag ctcttgtaga aaagctggca ggacgcgcgc tcatgaaaaa ataatgatag 1200 aactgcctgt ggtggagtgg cttgtttctc acggggcagt ttttgatagt ggaagggaga 1260 qattqttgaa tgtcagttca ttcagaagtc cttcatgctc tgcttaaaga tccgtttatt 1320 caqaaactga ttgatgcaga gcctgtattc tgggcaaatt caggcaagaa agaggggcca 1380 ttaccccgtg cagatgagtg ggcaaccgag atagcggaag cggaaaaaag aatgcagcgg 1440 tttgcacctt acattgccga ggtgtttcct gagacgaaag gcgctaaagg aatcatcgag 1500 tctccgcttt ttgaggtgca gcatatgaag ggaaagctgg aagcggcata tcagcagcca 1560 tttcccggaa gatggctttt aaagtgcgac catgagcttc cgatttcagg atcgattaaa 1620 gcgaggggcg ggatttatga agtgttaaag tatgctgaaa atctcgcgct tcaagaagga 1680 atgetteagg aaacegatga ttacegeate ttacaggaag ageggtttac egggtttte 1740 tecegetatt egattgetgt eggttegaea ggaaatetag gtttaageat eggeateate 1800 ggcgcggcac tcgggtttcg cgtgacagtg catatgtccg ccgatgctaa gcagtggaaa 1860 aaggatotoo toogocaaaa gggagtoaot gttatggagt acgaaacaga ttacagtgaa 1920 gcggtgaacg aagggagacg gcaggcggaa caagatccat tctgttattt tattgatgat 1980 gaacattoto gtoagotgtt ottaggatat gotgttgotg caagoogatt aaaaacacag 2040 cttgactgta tgaatataaa gccaagtett gagacgeeet tgtttgtgta tetgeegtge 2100 ggagtcggcg gaggaccggg cggtgtagca tttgggctga agcttttata cggagatgat 2160 gttcatgtgt ttttcgcaga accaactcat tcaccttgta tgctgttagg gctttattca 2220 ggacticacg agaagatete egiceaggat aleggeeigg alaateagae ggeigeigae 2280 🕙 ggacttgccg tagggaggcc gtcaggattt gtcggcaagc tgattgaacc gcttctgagc 2340 ggctgttata cggtagagga caatacgctt tatactttgc ttcatatgct ggctgtatct 2400 gaagataaat atttagagcc ctctyctctt gctggcatgt tcgggccggt tcagcttttt 2460 togacagaag agggaaggog ctatgotoag aaatataaga tggaacatgo ogtacatgto 2520 gtctggggaa cgggaggaag catggttcca aaagatgaaa tggctgcgta taaccgaatc 2580 ggtgctgatt tgctaaaaaa acgaaatgga aaataagcag acagtgaaaa ggttttecgt 2640 tacaatottt gtaagggttt taacotacag agagtcaggt gtaaacagtg aaaaataaag 2700 aacttaacct acatacttta tatacacagc acaatcggga gtcttggtct ggttttgggg 2760 ggcatttgtc gattgctgta tctgaagaag aggcaaaagc tgtggaagga ttgaatgatt 2820 atctatctgt tgaagaagtg gagacgatct atattccgct tgttcgcttg cttcatttac 2880 atgtcaagtc tgcggctgaa cgcaataagc atgtcaatgt ttttttgaag cacccacatt 2940 cagocaaaat toogtttatt atoggoattg coggoagtgt cgcagtogga aaaagcacga 3000 cggcgcggat cttgcagaag ctgctttcgc gtttgcctga ccgtccaaaa gtgagcctta 3060 tcacgacaga tggtttttta tttcctactg ccgagctgaa aaagaaaaat atgatgtcaa 3120 gaaaaggatt tootgaaago tatgatgtaa aggogotgot ogaatttttg aatgaottaa 3180

2	aatcaggaaa	ggacagcgta	aaggccccgg	tgtattccca	tctaacctat	gaccgcgagg	3240
ć	aaggtgtgtt	cgaggttgta	gaacaggcgg	atattgtgat	tattgaaggc	attaatgttc	3300
1	ttcagtcgcc	caccttggag	gatgaccggg	aaaacccgcg	tatttttgtt	tccgatttct	3360
1	ttgatttttc	gatttatgtg	gatgcggagg	aaagccggat	tttcacttgg	tatttagagc	3420
4	gttttcgcct	gcttcgggaa	acagcttttc	aaaatcctga	ttcatatttt	cataaattta	3480
•	aagacttgtc	cgatcaggag	gctgacgaga	tggcagcctc	gatttgggag	agtgtcaacc	3540
•	ggccgaattt	atatgaaaat	attttgccaa	ctaaattcag	gtcagatctc	attttgcgta	3600
	agggagacgg	gcataaggtc	gaggaagtgt	tggtaaggag	ggtatgaaat	gtgctgcagc	3,660
	tcgagcaata	gttaccctta	ttatcaagat	aagaaagaaa	aggattttc	gctacgctca	3720
	aatcctttaa	aaaaacacaa	aagaccacat	tttttaatgt	ggtctttatt	cttcaactaa	3780
	agcacccatt	agttcaacaa	acgaaaattg	gataaagtgg	gatatttta	aaatatatat	3840
	ttatgttaca	gtaatattga	cttttaaaaa	aggattgatt	ctaatgaaga	aagcagacaa	3900
	gtaagcctcc	taaattcact	ttagataaaa	atttaggagg	catatcaaat	gaactttaat	3960
	aaaattgatt	tagacaattg	gaagagaaaa	gagatattta	atcattattt	gaaccaacaa	4020
	acgactttta	gtataaccac	agaaattgat	attagtgttt	tataccgaaa	cataaaacaa	4080
	gaaggatata	aattttaccc	tgcatttatt	ttcttagtga	caagggtgat	aaactcaaat	4140
	acagctttta	gaactggtta	caatagcgac	ggagagttag	gttattggga	taaqttagag	42.00
	ccactttata	caatttttga	tggtgtatct	aaaacattct	ctggtatttg	gactcctgta	4260
	aagaatgact	tcaaagagtt	ttatgattta	tacctttctg	atgtagagaa	atataatggt	4320
	tcggggaaat	tgtttcccaa	aacacctata	cctgaaaatg	ctttttctct	ttctattatt	4380
	ccatggactt	catttactgg	gtttaactta	aatatcaata	ataatagtaa	ttaccttcta	4440
	cccattatta	cagcaggaaa	attcattaat	aaaggtaatt	caatatatt	accgctatct	4500
	ttacaggtac	atcattctgt	: ttgtgatggt	: tatcatgcag	gattgtttat	: gaactctatt	4560
	caggaattgt	cagataggco	taatgactgo	cttttataat	atgagataat	gccgactgta	4620
	ctttttacag	g tcggttttct	aatgtcacta	acctgccccg	, ttagttgaag	g aaggtttta	4680
	tattacagct	gtcgactcgt	gatcttcgga	caggctgttc	agetttte	caatgcgatc	4740
	cagctgcgct	tttcggtttt	tcgcatactt	gaagcctgta	acageegea	a agacgacago	4800
	ggcaaatata	a ataaatacaa	acagetgaaa	a catcacatca	cctatattc	a tgttcttcac	4860
							4020

cgggaacggt gttttgtggt ctccattttc atttgccgat aggcgaacgc taaaaatggc 4980 aggeegagea gggtaatgee geteaggaea gaaaaaatat aaateggeeg geeagegeea 5040 aacaggtota tacatatoco cocgacocaa gggoogatga ogtttoogag otgtggaaaa 5100 ccgattgccc cgaaataagt gccttttaat cctggttttg caatctggtc tacatacaaa 5160 tocatcatag agaataaaag cacttogoog attgtaaatg tgatgacaat catcacaatt 5220 gatggaacac cgtgtgatac ggtgaaaatg gccatgctga tgctaaccat cacattaccg 5280 agcatcagag aacaaagcgg cgaaaaccgt tttgcaaaat ggacaatggg aaattgcgtc 5340 gccaacacaa cgattgcgtt taatgtcagc atcagcccat acagcttcgt tccattgccg 5400 atcaaggggt tctgcgccat atactgaggg aatgtggaac tgaattgtga gtagccgaag 5460 gtgcatageg taatgeegae caaageaatg gtaaaaagat aateettig egtgaeeata 5520 aacgetteee geacgeteat atttegggae tgggetggtg etgataagga tggatgtttt 5580 ttaaattgga gggcaagcac aattccgtat agtccgtaaa tgactgcagg caccaaaaag 5640 ggcgtagtcg attgcgatga gccgaaatat aggccaagca caggtccgaa gacaacgccg 5700 atattaatag cogcatagog taaattaaaa actagoagto togiittito tioigicata 5760 🕟 tcagacaaca aggeetttga agegggetea aacagtgatt tgcaaagace gtttaatgeg 5820 tttactacaa aaaacaccca gagattagat gctgccgcaa agcctgcaaa taccagcatc 5880 cateegaaaa tegatacaag cateatgttt tttetgeega atttatetga gatatateeg 5940 ccgtaaaagc ttgcgaggat gccgactgat gagctcgcgg cgatgaccag ccctgcatag 6000 gaagetgatg egeettggae ggetgteaaa taaategeta aaaaaggaat geteategat 6060 gttgccattc tgccgaaaat ggttccgatt ataattgtac gcgttggatg catagcttga 6120 gtattctata gtgtcaccta aatagcttgg cgtaatcatg gtcatagctg tttcctgtgt 6180 gaaattgtta teegeteaca atteeacaca acataegage eggaageata aagtgtaaag 6240 cctggggtgc ctaatgagtg agctaactca cattaattgc gttgcgctca ctgcccgctt 6300 tccagtcggg aaacctgtcg tgccagctgc attaatgaat cggccaacgc gcggggagag 6360 geggtttgeg tattgggege tetteegett cetegeteae tgactegetg egeteggteg 6420 ttcggctgcg gcgagcggta tcagctcact caaaggcggt aatacggtta tccacagaat 6480 caggggataa cgcaggaaag aacatgtgag caaaaggcca gcaaaaggcc aggaaccgta 6540 aaaaggccgc gttgctggcg tttttcgata ggctccgccc ccctgacgag catcacaaaa 6600 atcgacgctc aagtcagagg tggcgaaacc cgacaggact ataaagatac caggcgtttc 6660

cccctggaag ctccctcgtg cgctctcctg ttccgaccct gccgcttacc ggatacctgt 6720 cogcettet ecetteggga agegtggege ttteteatag eteaegetgt aggtatetea 6780 gttcggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaacccccc gttcagcccg 6840 accgctgcgc cttatccggt aactatcgtc ttgagtccaa cccggtaaga cacgacttat 6900 cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta ggcggtgcta 6960 cagagttctt gaagtggtgg cctaactacg gctacactag aaggacagta tttggtatct 7020 gcgctctgct gaagccagtt accttcggaa aaagagttgg tagctcttga tccggcaaac 7080 aaaccaccgc tggtagcggt ggtttttttg tttgcaagca gcagattacg cgcagaaaaa 7140 aaggatetea agaagateet ttgatetttt etaeggggte tgaegeteag tggaaegaaa 7200 actcacgtta agggattttg gtcatgagat tatcaaaaag gatcttcacc tagatccttt 7260 taaattaaaa atgaagtttt aaatcaatct aaagtatata tgagtaaact tggtctgaca 7320 gttaccaatg cttaatcagt gaggcaccta tctcagcgat ctgtctattt cgttcatcca 7380 tagttgcctg actccccgtc gtgtagataa ctacgatacg ggagggctta ccatctggcc 7440 ccagtgctgc aatgataccg cgagacccac gctcaccggc tccagattta tcagcaataa 7500 accagecage eggaagggee gagegeagaa geggteeege aactttatee geeteeatee 7560 agtotattaa tigitgoogg gaagotagag taagtagtio gooagitaat agtitgogoa 7620 acgttgttgg cattgctaca ggcatcgtgg tgtcacgctc gtcgtttggt atggcttcat 7680 tcagctccgg ttcccaacga tcaaggcgag ttacatgatc ccccatgttg tgcaaaaaa 7740 eggttagete etteggteet eegategttg teagaagtaa gttggeegea gtgttateae 7800 tcatggttat ggcagcactg cataattctc ttactgtcat gccatccgta agatgctttt 7860 ctgtgactgg tgagtactca accaagtcat tctgagaata ccgcgcccgg cgaccgagtt 7920 gctcttgccc ggcgtcaata cgggataata gtgtatgaca tagcagaact ttaaaagtgc 7980 tcatcattgg aaaacgttct tcggggcgaa aactctcaag gatcttaccg ctgttgagat 8040 ccagttcgat gtaacccact cgtgcaccca actgatcttc agcatctttt actttcacca 8100 gcgtttctgg gtgagcaaaa acaggaaggc aaaatgccgc aaaaaaggga ataagggcga 8160 cacggaaatg ttgaatactc atactettee tttttcaata ttattgaage atttatcagg 8220 gttattgtct catgagcgga tacatatttg aatgtattta gaaaaataaa caaatagggg 8280 ttccgcgcac atttccccga aaagtgccac ctgtatgcgg tgtgaaatac cgcacagatg 8340 cgtaaggaga aaataccgca tcaggcgaaa ttgtaaacgt taatattttg ttaaaattcg 8400

ogttaaatat ttgttaaatc ageteattt ttaaceaata ggeegaaate ggeaaaatee 8460 ettataaate aaaagaatag acegagatag ggttgagtgt tgtteeagtt tggaacaaga 8520 gteeactatt aaagaacgtg gaeteeaacg teaaagggeg aaaaacegte tateagggeg 8580 atggeecact acgtgaacea teaeceaaat caagttttt geggtegagg tgeegtaaag 8640 etetaaateg gaaceetaaa gggageecee gatttagage ttgaegggga aageeggega 8700 acgtggegag aaaggaaggg aagaaagega aaggageggg egetagggeg etggeaagtg 8760 tageggteae getgeegta aceaecacae eegeegget taa 8803

<210> 73

<211> 8320

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
 pAN296 plasmid

<400> 73 tgcgccgcta cagggcgcgt ccattcgcca ttcaggctgc gcaactgttg ggaagggcga 60 teggtgeggg cetetteget attacgeeag etggegaaag ggggatgtge tgeaaggega 120 ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagtgaa 180 ttgtaatacg actcactata gggcgaattg ggcccgacgt cgcatgctgg atgaaaagcc 240 gatgaccgct tttcaggtct gtcagcagct ttttcctgct gtatatgaaa aggaattgtt 300 tttaacgatg tcagaaacgg caggtcacct tgatgtgttg gaggctgaag aagccatcac 360 gtcatattgg gaaggaaata ccgtatactt taaaacaatg aagaggtgaa atgggtgaaa 420 catatagcgg gaaaaaggat ttggataacc ggcgcttcag gagggcttgg agaaagaatc 480 gcatacttat gcgcggctga aggagcccat gtcctgctgt cggctagacg cgaggatcgt 540 ttgatagaaa tcaaaaggaa aataaccgag gaatggagcg gacagtgtga gatttttcct 600 ctggatgtcg gccgcctaga ggatatcgcc cgggtccgcg atcagatcgg ctcgattgat 660 gtactgatta acaatgcagg cttcggtata tttgaaacgg ttttagactc tacattggat 720 gacatgaaag cgatgtttga tgtgaatgtc ttcggcctga tcgcctgtac aaaagcggtg 780 cttccgcaaa tgcttgagca aaaaaaggga catatcatca atatcgcctc tcaagcgggg 840 aaaatcgcca caccgaagtc tagcctgtat tccgcgacca aacatgccgt gttaggttac 900 tcaaacgctt tgcggatgga gctttcggga accggcattt atgtgacaac agtcaacccg 960 ggcccgattc agacggactt tttttccatt gctgataaag gcggggacta cgccaaaaat 1020 gtcggccgct ggatgcttga tcctgatgac gtggcagctc aaattacagc tgcaattttt 1080 acgaaaaagc gggagatcaa tcttccgcgt ttaatgaatg ccggcactaa gctgtatcag 1140 ctgtttccag ctcttgtaga aaagctggca ggacgcgcgc tcatgaaaaa ataatgatag 1200 aactgcctgt ggtggagtgg cttgtttctc acggggcagt ttttgatagt ggaagggaga 1260 gattgttgaa tgtcagttca ttcagaagtc cttcatgctc tgcttaaaga'tccgtttatt 1320 cagaaactga ttgatgcaga gcctgtattc tgggcaaatt caggcaagaa agaggggcca 1380 ttaccccgtg cagatgagtg ggcaaccgag atagcggaag cggaaaaaag aatgcagcgg 1440 tttgcacctt acattgccga ggtgtttcct gagacgaaag gcgctaaagg aatcatcgag 1500 tctccgcttt ttgaggtgca gcatatgaag ggaaagctgg aagcggcata tcagcagcca 1560 tttcccggaa gatggctttt aaagtgcgac catgagcttc cgatttcagg atcgattaaa 1620 gcgaggggcg ggatttatga agtgttaaag tatgctgaaa atctcgcgct tcaagaagga 1680 atgcttcagg aaaccgatga ttaccgcatc ttacaggaag agcggtttac cgggttttc 1740 tecegetatt egattgetgt eggttegaea ggaaatetag gtttaageat eggeateate 1800 qqcqcqqcac tcqqqtttcq cqtqacaqtq catatqtccq ccqatqctaa gcaqtqgaaa 1860 aaggatetee teegeeaaaa gggagteaet gttatggagt acgaaacaga ttacagtgaa 1920 gcqqtgaacg aagggagacg gcaggcggaa caagatccat tctgttattt tattgatgat 1980 gaacattete gteagetgtt ettaggatat getgttgetg caageegatt aaaaacacag 2040 cttgactgta tgaatataaa gccaagtctt gagacgccct tgtttgtgta tctgccgtgc 2100 ggagtcggcg gaggaccggg cggtgtagca tttgggctga agcttttata cggagatgat 2160 gttcatgtgt ttttcgcaga accaactcat tcaccttgta tgctgttagg gctttattca 2220 ggacttcacg agaagatctc cgtccaggat atcggcctgg ataatcagac ggctgctgac 2280 qqacttgeeg tagggaggee gteaggattt gteggeaage tgattgaace gettetgage 2340 qqctqttata cggtagagga caatacgctt tatactttgc ttcatatgct ggctgtatct 2400 qaaqataaat atttagagcc ctctgctctt gctggcatgt tcgggccggt tcagcttttt 2460 tcgacagaag agggaaggcg ctatgctcag aaatataaga tggaacatgc cgtacatgtc 2520 gtctggggaa cgggaggaag catggttcca aaagatgaaa tggctgcgta taaccgaatc 2580 ggtgctgatt tgctaaaaaa acgaaatgga aaataagcag acagtgaaaa ggttttccgt 2640 tacaatettt gtaagggttt taacetacag agagteaggt gtaaacagtg aaaaataaag 2700 aacttaacct acatacttta tatacacagc acaatcggga gtcttctgca gctcgagcaa 2760

tagttaccct tattatcaag ataagaaaga aaaggatttt tcgctacgct caaatccttt 2820 aaaaaaacac aaaagaccac atttttaat gtggtcttta ttcttcaact aaagcaccca 2880 ttagttcaac aaacgaaaat tggataaagt gggatatttt taaaatatat atttatgtta 2940 cagtaatatt gacttttaaa aaaggattga ttctaatgaa gaaagcagac aagtaagcct 3000 cctaaattca ctttagataa aaatttagga ggcatatcaa atgaacttta ataaaattga 3060 tttagacaat tggaagagaa aagagatatt taatcattat ttgaaccaac aaacgacttt 3120 tagtataacc acagaaattg atattagtgt tttataccga aacataaaac aagaaggata 3180 taaattttac cctgcattta ttttcttagt gacaagggtg ataaactcaa atacagcttt 3240 tagaactggt tacaatagcg acggagagtt aggttattgg gataagttag agccacttta 3300 tacaattttt gatggtgtat ctaaaacatt ctctggtatt tggactcctg taaagaatga 3360 cttcaaagag ttttatgatt tatacctttc tgatgtagag aaatataatg gttcggggaa 3420 attgtttccc aaaacaccta tacctgaaaa tgctttttct ctttctatta ttccatggac 3480 ttcatttact gggtttaact taaatatcaa taataatagt aattaccttc tacccattat 3540 tacaqcaqqa aaattcatta ataaaggtaa ttcaatatat ttaccgctat ctttacaggt 3600 acatcattct gtttgtgatg gttatcatgc aggattgttt atgaactcta ttcaggaatt 3660 gtcagatagg cctaatgact ggcttttata atatgagata atgccgactg tacttttac 3720 agtoggtttt otaatgtoac taacotgooc ogttagttga agaaggtttt tatattacag 3780 ctgtcgacta aggtcgagga agtgttggta aggagggtat gaaatgtgca tcatattgaa 3840 ctgtatgtct ctgatttgga ggcgtctagg cggttttggg gctggttctt aaaagaactt 3900 ggttataaag agtatcaaaa atggagctca ggcatcagct ggaagaaaga tcgtttttac 3960 ctagtgattg tgcaggcgaa agagccattt ctagagccgg aataccatag atgccgagtc 4020 ggtctgaacc atctcgcatt tcatgctgaa tccaagcttc aagtcgatca gatgactgaa 4080 aaattgacgg caaaaggcta tcgtgtgttg taccgagaca ggcatccttt tgccggagga 4140 gacgggcatt atgcagtctt ttgtgaggat ccagaccgga ttaaggtaga gctcgttgcc 4200 ccaagetgtt aategtgate tteggacagg etgtteaget tttteteaat gegateeage 4260 tgcgcttttc ggtttttcgc atacttgaag cctgtaacag ccgcaaagac gacagcggca 4320 aatataataa atacaaacag ctgaaacatc acatcaccta tattcatgtt cttcacctca 4380 tgtttgcggg agagattcat tctcttccgt tttttattta aagcggcttt tccagacggg 4440 aacqqtqttt tqtqqtctcc attttcattt qccqataqqc gaacqctaaa aatqqcaqqc 4500

0121772A2 | >

cgagcagggt aatgccgctc aggacagaaa aaatataaat cggccggcca gcgccaaaca 4560 ggtctataca tatccccccg acccaagggc cgatgacgtt tccgagctgt ggaaaaccga 4620 ttgccccgaa ataagtgcct tttaatcctg gttttgcaat ctggtctaca tacaaatcca 4680 tcatagagaa taaaagcact tcgccgattg taaatgtgat gacaatcatc acaattgatg 4740 gaacaccgtg tgatacggtg aaaatggcca tgctgatgct aaccatcaca ttaccgagca 4800 tcagagaaca aagcggcgaa aaccgttttg caaaatggac aatgggaaat tgcgtcgcca 4860 acacaacgat tgcgtttaat gtcagcatca gcccatacag cttcgttcca ttgccgatca 4920 aggggttctg cgccatatac tgagggaatg tggaactgaa ttgtgagtag ccgaaggtgc 4980 atagogtaat googaccaaa goaatggtaa aaagataato ottttgogtg accataaacg 5040 cttcccgcac gctcatattt cgggactggg ctggtgctga taaggatgga tgttttttaa 5100 attggagggc aagcacaatt ccgtatagtc cgtaaatgac tgcaggcacc aaaaagggcg 5160 tagtcgattg cgatgagccg aaatataggc caagcacagg tccgaagaca acgccgatat 5220 taatageege atagegtaaa ttaaaaaeta geagtetegt tttttettet gteatateag 5280 acaacaaggc ctttgaagcg ggctcaaaca gtgatttgca aagaccgttt aatgcgttta 5340 ctacaaaaaa cacccagaga ttagatqctg ccgcaaaqcc tqcaaatacc aqcatccatc 5400 cgaaaatcga tacaagcatc atgttttttc tgccgaattt atctgagata tatccgccgt 5460 aaaagettge gaggatgeeg actgatgage tegeggegat gaccageeet geataggaag 5520 ctgatgcgcc ttggacggct gtcaaataaa tcgctaaaaa aggaatgctc atcgatgttg 5580 ccattctgcc gaaaatggtt ccgattataa ttgtaacgcg ttggatgcat agcttgagta 5640 ttctatagtg tcacctaaat agcttggcgt aatcatggtc atagctgttt cctgtgtgaa 5700 attgttatcc gctcacaatt ccacacaaca tacgagccgg aagcataaag tgtaaagcct 5760 ggggtgccta atgagtgagc taactcacat taattgcgtt gcgctcactg cccgctttcc 5820 agtegggaaa eetgtegtge eagetgeatt aatgaategg eeaaegegeg gggagaggeg 5880 gtttgcgtat tgggcgctct tccgcttcct cgctcactga ctcgctgcgc tcggtcgttc 5940 ggctgcggcg agcggtatca gctcactcaa aggcggtaat acggttatcc acagaatcag 6000 qqqataacgc aggaaagaac atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa 6060 aggccgcgtt gctggcgttt ttcgataggc tccgccccc tgacgagcat cacaaaaatc 6120 gacgeteaag teagaggtgg egaaaceega eaggaetata aagataceag gegttteeee 6180 ctggaagete cetegtgege tetectgtte egaceetgee gettaeegga taeetgteeg 6240

cettletece ttegggaage gtggegettt eteatagete aegetgtagg tateteagtt 6300 cggtgtaggt cgttcgctcc aagctgggct gtgtgcacga accccccgtt cagcccgacc 6360 gctgcgcctt atccggtaac tatcgtcttg agtccaaccc ggtaagacac gacttatcgc 6420 cactggcagc agccactggt aacaggatta gcagagcgag gtatgtaggc ggtgctacag 6480 agttettgaa gtggtggeet aactaegget acactagaag gacagtattt ggtatetgeg 6540 ctctgctgaa gccagttacc ttcggaaaaa gagttggtag ctcttgatcc ggcaaacaaa 6600 ccaccgctgg tagcggtggt ttttttgttt gcaagcagca gattacgcgc agaaaaaaag 6660 gateteaaga agateetttg atettteeta eggggtetga egeteagtgg aacgaaact 6720 cacgttaagg gattttggtc atgagattat caaaaaggat cttcacctag atccttttaa 6780 attaaaaatg aagttttaaa tcaatctaaa gtatatatga gtaaacttgg tctgacagtt 6840 accaatgett aatcagtgag geacetatet cagegatetg tetatttegt teatecatag 6900 ttgcctgact ccccgtcgtg tagataacta cgatacggga gggcttacca tctggcccca 6960 gtgctgcaat gataccgcga gacccacgct caccggctcc agatttatca gcaataaacc 7020 agceageegg aagggeeyag egeagaagtg gteetgeaac titateegee teeateeagt 7080 ctattaattg ttgccgggaa gctagagtaa gtagttcgcc agttaatagt ttgcgcaacg 7140 ttgttggcat tgctacaggc atcgtggtgt cacgctcgtc gtttggtatg gcttcattca 7200 getoeggtte ccaacgatea aggegagtta catgateece catgttgtge aaaaaagegg 7260 ttagctcctt cggtcctccg atcgttgtca gaagtaagtt ggccgcagtg ttatcactca 7320 tggttatggc agcactgcat aattetetta etgteatgcc ateegtaaga tgettteetg 7380 tgactggtga gtactcaacc aagtcattct gagaataccg cgcccggcga ccgagttgct 7440 cttgcccggc gtcaatacgg gataatagtg tatgacatag cagaacttta aaagtgctca 7,500 tcattggaaa acgttcttcg gggcgaaaac tctcaaggat cttaccgctg ttgagatcca 7560 gttcgatgta acceactcgt gcacceaact gatettcage atettttact ttcaccageg 7620 tttctgggtg agcaaaaaca ggaaggcaaa atgccgcaaa aaagggaata agggcgacac 7680 ggaaatgttg aatactcata ctcttccttt ttcaatatta ttgaagcatt tatcagggtt 7740 attgtctcat gagcggatac atatttgaat gtatttagaa aaataaacaa ataggggttc 7800 cgcgcacatt tccccgaaaa gtgccacctg tatgcggtgt gaaataccgc acagatgcgt 7860 aaggagaaaa taccgcatca ggcgaaattg taaacgttaa tattttgtta aaattcgcgt 7920 taaatatttg ttaaatcagc tcattttta accaataggc cgaaatcggc aaaatccctt 7980 ataaatcaaa agaatagacc gagatagggt tgagtgttgt tccagtttgg aacaagagtc 8040 cactattaaa gaacgtggac tccaacgtca aagggcgaaa aaccgtctat cagggcgatg 8100 gcccactacg tgaaccatca cccaaatcaa gtttttgcg gtcgaggtgc cgtaaagctc 8160 taaatcggaa ccctaaaggg agcccccgat ttagagcttg acggggaaag ccggcgaacg 8220 tggcgagaaa ggaagggaag aaagcgaaag gagcggcgc tagggcgctg gcaagtgtag 8280 cggtcacgct gcgcgtaacc accacacccg ccgcgcttaa 8320

<210> 74

<211> 250

<212> PRT

<213> Clostridium acetobutylicum

<400> 74

Asn Lys Arg Ala Ala Phe Met Leu Leu Leu Phe Leu Arg Ser Val Leu

1 10 , 15

Lys Val Ile Leu Val Leu Asp Val Gly Asn Thr Asn Ile Val Leu Gly 20 25 30

Ile Tyr Asn Asp Thr Lys Leu Thr Ala Glu Trp Arg Leu Ser Thr Asp 35 40 45

Val Leu Arg Ser Ala Asp Glu Tyr Gly Ile Gln Val Met Asn Leu Phe 50 55 60

Gln Gln Asp Lys Leu Asp Pro Thr Leu Val Glu Gly Val Ile Ile Ser 65 70 75 80

Ser Val Val Pro Asn Ile Met Tyr Ser Leu Glu His Met Ile Arg Lys 85 90 95

Tyr Phe Lys Ile Asn Pro Leu Val Val Gly Pro Gly Ile Lys Thr Gly 100 105 110

Ile Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile 115 120 125

Val Asn Ala Val Ala Ala His Glu Ile Tyr Lys Arg Ser Leu Ile Ile 130 135 140

Ile Asp Phe Gly Thr Ala Thr Thr Phe Cys Ala Val Arg Glu Asn Gly 145 150 155 160

Asp Tyr Leu Gly Gly Ala Ile Cys Pro Gly Ile Lys Val Ser Ser Glu 165 170 175

Ala Leu Phe Glu Lys Ala Ala Lys Leu Pro Arg Val Glu Leu Ile Lys 180 185 190

Pro Ala Tyr Ala Ile Cys Lys Asn Thr Ile Ser Ser Ile Gln Ser Gly 195 200 205 Ile Val Tyr Arg Tyr Leu Arg Gln Val Lys Tyr Leu Phe Glu Lys Leu 210 215 220

Lys Glu Asn Leu Pro Asp Gly Arg Arg Thr Arg Thr Ser Leu Val Leu 225 230 235 240

Ala Thr Gly Gly Leu Ala Lys Leu Ile Asn 245 250

<210>.75

<211> 258

<212> PRT

<213> Rhodobacter capsulatus

<400> 75

Met Leu Leu Cys Ile Asp Cys Gly Asn Thr Asn Thr Val Phe Ser Val 1 5 10 15

Trp Asp Gly Thr Asp Phe Ala Ala Thr Trp Arg Ile Ala Thr Asp His
20 25 30

Arg Arg Thr Ala Asp Glu Tyr Phe Val Trp Leu Asn Thr Leu Met Gln 35 40 45

Leu Lys Gly Leu Gln Gly Arg Ile Ser Glu Ala Ile Ile Ser Ser Thr 50 55 60

Ala Pro Arg Val Val Phe Asn Leu Arg Val Leu Cys Asn Arg Tyr Phe 65 70 75 8C

Asp Cys Arg Pro Tyr Val Val Gly Lys Pro Gly Cys Glu Leu Pro Val 85 90 95

Ala Pro Arg Val Asp Pro Gly Thr Thr Val Gly Pro Asp Arg Leu Val
100 105 110

Asn Thr Val Ala Gly Tyr Asp Arg His Gly Gly Asp Leu Ile Val Val 115 120 125

Asp Phe Gly Thr Ala Thr Thr Phe Asp Val Val Ala Pro Asp Gly Ala 130 135 140

Tyr Ile Gly Gly Val Ile Ala Pro Gly Val Asn Leu Ser Leu Glu Ala 145 150 155 160

Leu His Met Ala Ala Ala Leu Pro His Val Asp Val Thr Lys Pro 165 170 175

Gln Gly Val Ile Gly Thr Asn Thr Val Ala Cys Ile Gln Ser Gly Val 180 185 190

Tyr Trp Gly Tyr Ile Gly Leu Val Glu Gly Ile Val Arg Gln Ile Arg 195 200 205

Met Glu Arg Asp Arg Pro Met Lys Val Ile Ala Thr Gly Gly Leu Ala 210 215 220 Ser Leu Phe Asp Leu Gly Phe Asp Leu Phe Asp Lys Val Glu Asp Asp 225 230 235 240

Leu Thr Met His Gly Leu Arg Leu Ile Phe Asp Tyr Asn Lys Gly Leu 245 250 255

Gly Ala

<210> 76

<211> 10801

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
pAN240 plasmid

<400> 76 gaattttgcg gccgcttcga aagctgtaat ataaaaacct tcttcaacta acggggcagg 60 ttagtgacat tagaaaaccg actgtaaaaa gtacagtcgg cattatctca tattataaaa 120 gccagtcatt aggcctatct gacaattcct gaatagagtt cataaacaat cctgcatgat 180 aaccatcaca aacagaatga tgtacctgta aagatagcgg taaatatatt gaattacctt 240 tattaatgaa ttttcctgct gtaataatgg gtagaaygta attactatta ttattgatat 300 ttaagttaaa cccagtaaat gaagtccatg gaataataga aagagaaaaa gcattttcag 360 gtataggtgt tttgggaaac aatttccccg aaccattata tttctctaca tcagaaaggt 420 ataaatcata aaactctttg aagtcattct ttacaggagt ccaaatacca gagaatgttt 480 tagatacacc atcaaaaatt gtataaagtg gctctaactt atcccaataa cctaactctc 540 cgtcgctatt gtaaccagtt ctaaaagctg tatttgagtt tatcaccctt gtcactaaga 600 aaataaatgc agggtaaaat ttatatcctt cttgttttat gtttcggtat aaaacactaa 660 tatcaatttc tgtggttata ctaaaagtcg tttgttggtt caaataatga ttaaatatct 720 cttttctctt ccaattgtct aaatcaattt tattaaagtt catttgatat gcctcctaaa 780 tttttatcta aagtgaattt aggaggctta cttgtctgct ttcttcatta gaatcaatcc 840 ttttttaaaa gtcaatatta ctgtaacata aatatatatt ttaaaaatat cccactttat 900 ccaattttcg tttgttgaac taatgggtgc tttagttgaa gaataaagac cacattaaaa 960 aatgtggtct tttgtgtttt tttaaaggat ttgagcgtag cgaaaaatcc ttttcttct 1020 tatcttgata ataagggtaa ctattgaatt cggtaccaag agtttgtaga aacgcaaaaa 1080 ggccatccgt caggatggcc ttctgcttaa tttgatgcct ggcagtttat ggcgggcgtc 1140

ctgcccgcca ccctccgggc cgttgcttcg caacgttcaa atccgctccc ggcggatttg 1200 toctactcag gagagogtto accgacaaac aacagataaa acgaaaggoo cagtotttog 1260 actgageett tegttttatt tgatgeetgg cagtteeeta etetegeatg gggagaeece 1320 acactaccat cggcgctacg gcgtttcact tctgagttcg gcatggggtc aggtgggacc 1380 ' accgcgctac tgccgccagg caaattctgt tttatcagac cgcttctgcg ttctgattta 1440 atctgtatca ggctgaaaat cttctctcat ccgccaaaac aggatcctac ggaaatggag 1500 cggcaaaacc gttttactct caaaatctta aaagaaaacc cccgataaag ggggcttttc 1560 ttctacaaaa ttgtacgggc tggttcgttc cccagcattt gttcaatttt gttttgatca 1620 ttcagaacag ccactttcgg ctcatggctt gccgcttctt gatcagacat cattttgtag 1680 gaaataataa tgaccttatc teetteetge acaaggegtg eggetgeace gtttaageat 1740 atgacgccgc ttccccgttt accaggaata atatacgttt caagacgtgc tccattatta 1800 ttattcacaa tttgtacttt ttcattagga agcattccca cagcatcaat gagatcttca 1860 tcaattgtaa tgcttcccac atagttcagg tttgcttccg taacagttgc cctgtgaagt 1920 ttgccgctca tcattgttcg atacatatta tattctctcc atttctcgaa tatcaataat 1990 gatattatet attaaacgeg ettttgaaaa ageaactgea acagegagaa teatettee 2040 agcaatttca ttcacagget egagtteegg ataggaataa agetetacat agtetatggt 2100 teegetagte gttteaatga tatettttge agettttate accgetteag gatetettte 2160 accggcttgg acaagttccg cacttgtttg aagggcccga tacagcttag gcgcttcttt 2220 tettteetea getgttaagt atacattgeg agagettttg getaageegt etteetetet 2280 gaccytatcy acaggaacca attcaatatc catgaagaag tegetgatta acceatcaac 2340 aacagctacc tgctgcgcat cttttaaacc gaaataggca cgagtcggct tgactagatt 2400 gaaaagette gteagtaega tegegaeeee gteaaaatgt eettetettg agegeeegea 2460 taacacgtct gtgcgtcttt ctacatgaat cgtgacattc ttttcaccgg gatacatatc 2520 atgagcatet ggegtaaaaa gaatategae teeggegttt tetgeaagag etgeateeeg 2580 ctcaatatcg cgcggatatg cttcaaaatc ttcattaggg ccgaattgtg caggattcac 2640 aaaaatactc ataataacgg cgtcgttttc ttgtcttgct ttgtctgcta aggttaaatg 2700 eccetcatge agaaacecca tegteggaac aaateegatt gaettgeeet etgaatggta 2760 ttgttttatg gcttctttca gctgtgaaat atcagtaatc tgtctcatct tattttcccc 2820 cgtacaagcc gtcaagcact gtctggttca tttgaaagga atgcttttgt tcagggaaag 2880

cacgatgtct tacatectga acatateege tgattgetgt ttegatggtt teateaatge 2940 gcgtatattg ctttacaaat ttaggtgttc tctcaacacc gtggccgata atatcatgat 3000 aaacgagaac tigiccgicc gcittcacac cagccccgat iccaatgacc ggiatgcita 3060 gcgtctcggc aattttggct gtgagttctg ccggcacaca ttccagcaca agcatcatag 3120 ctcctgcttc ttcgcatttt atactgtctt ctattaattt tttggcgctt tgttcgtctt 3180 tgccctgtac tttatagccg cccagtacgc cgactgactg cggtgtcaaa cctaagtgac 3240 tgactactgg aatgcctcca agcgtcaatg cgcgaatgga ttcaaacacg ccttctccgc 3300 ceteaagett cagtgegtea geteegettt cetgaacgat ageegetgea tttttcageg 3360 tatetteett agacaggtga taagacataa aeggeatate tgteacaata aaggtatteg 3420 gcgcacccct tttaacggct tttgtatgat ggatcatgtc cgcaactgtc acaccgacag 3480 ttgaatcaag gccgaggacg accattccaa gtgaatcacc gactaaaatc atgtcaactc 3540 ccgcttgttc agcaagttta gctgccggat aatcataagc ggtcagcatg acaatcggtt 3600 gagogtoctg ctgttgttaa gattattata coacaccttg tagataaagt caacaacttt 3720 ttgcaaaatt tttcaggaat tttagcagag gttgttctgg atgtagaaca aaacatcttt 3780 ccgctcttgt gctgttagga tatctttctt ggaagctagg taggcctcga gttatggcag 3840 ttggttaaaa ggaaacaaaa agaccgtttt cacacaaaac ggtcttttc gatttctttt 3900 tacagtcaca gccacttttg caaaaaccgg acagcttcat gccttataac tgctgtttcg 3960 gtcgacgatg atctgccgtt ttcttctgca agccaaaaaa ccttccgtta caacgagaag 4020 gattetteae tttetaaagt teggegagtt teatecetet gteecagtee ttttttggat 4080 caaggcagac tgctgcaatg tctatctatt ttaataatag gtgcagttcg caggcgatac 4140 tgcccaatgg aagtatacca aaatcaacgg gcttgtacca acacattagc ccaattcgat 4200 atcggcagaa tagattttt taatgccttc gttcgtttct aaaagcagaa cgccttcatc 4260 atctatacct aacgccttac cgtaaaaggt tccgtttaac gttctggctc tcatattagt 4320 gccaataccg agcgcatage tttcccataa aagcttaatc ggcgtaaatc cgtgcgtcat 4380 ataatcccgg taccgtttct caaagcatag taaaatatgc tggatgacgc cggcccgatc 4440 aattttttcc ccagcagctt ggctgaggct tgtcgcgatg tccttcaatt catctggaaa 4500 atcattaggc tgctggttaa cgttaatgcc gatcccaatg atcactgaac gtacgcggtc 4560 ttcttcagcc tgcatttccg ttaggatacc gactgttttt tttccgttaa tcaaaatatc 4620

atttggccat	ttaatateeg	tttggatgcc	ractacter	tetatteeet	gcacaacagc	4000
tactgcagca	agcagagtca	gctgcggtgt	tttttggagc	ggaatgtcag	gccgcaaaat	4740
caggctcatc	caaacaccgt	ttocttcttg	agaatgccat	accctagaca	tteggcccct	4800
tccggctgtt	tgtttgtcag	ccaccacaag	ggtgccttcc	ggtgcgttat	ţattcgcgag	48'60'
ctcatgagcc	gttttttgcg	tgcttgaaag	aacgtcatgg	taaataagat	gctggcccat	4920
cacttccgtt	tttaatccaa	aacgaatttc	gctttcactg	agttttccgg	gttttttgat	4980
gagccgatat	ccttttcttc	taacggcttc	tacttcataa	ccctctttcc	gaagetette	5040
aatatgcttc	cacacagcag	ttcttgaaca	gccgagagca	tcactgattt	tttggccgga	5100
aataaattca	ttgccggcct	gagaaaataa	ttcaataagg	tetttetta	atgttgaccg	5160
catgtcttca	gccactcctc	tatgtgtttc	ttttgattgg	agagettece	tgtcacaaca	52-20
gcctgctcga	tccactgtaa	ttcttctgac	acccattttc	cggccggccg	gtttcgaagc	5280
gcaagcaagt	ccttacccgt	gatatcaaga	teettaagge	ttttgatcgg	caggttttga	5340
taagcgtact	gaatgtcctt	cagtttcttt	tcatccagtt	tttcgttttg	ccgaagctgc	5400
gatattttgg	ccgctgagag	cagtgctttt	ttoccageto	tgtacattgt	cattgcgtca	54·60
aggctctggc	caaacgtatc	ggcaatgtga	atggcttcct	tgatcacttt	toccgggagc	5520
ttccaggctt	tcaggaaaag	gggcgcgtct	ttcaaaacta	tgccaaggtt	aattaaaaga	5580
gcagcccaaa	gctcctcacg	ggatgttaaa	gagaagaato	gaaactcact	cgttgaaatc	5640
aggttttctc	gtttatgata	aaaaccagga	agctcttcat	acaatctcgt	ttgaatgagt	5700
gtttgaagcg	cctggcgaga	agctcttccc	: tgcagcaatt	tctcaaactc	tatagttttt	5760
cgttcgactq	g aaacatggga	gaggagtgat	ttttctttc	g caatggcttc	ttctgtttcc	5820
ggtgaaagc	g taaagccaag	ctggctcata	aagcgtacg	g ctctcagcat	. acgaagcgca	5880
tectettgaa	a atctatcctc	aggettteca	acggttcga	tcactttctg	, atcaatatct	5940
ttcttgccg	caaaataato	aagcacctto	ccgtccgct	g tcatggccat	cgcattgatc	6000
gttaaatct	tgcgttttag	atcctcttct	aatgatgag	a taaattgcad	ttctgacggt	6060
cttctgaaa	t caacataato	: agattcagtd	cggaatgtc	g tgacttcata	ggtttcatco	6120
tcccagagc	a caataatggt	cccgtgctct	ttgcctaca	t caacagtcc	g ctgaaacago	6180
cgttctact	t gatcaggtgo	cgcatctgto	gcgatatcg	a catctccgat	cgttcgtttc	6240
atatagctg	t cacgaactgo	geceegae	a aaataagcc	t gatggcccg	c ttcgattaaq	g 6300
atgcggagc	a cgggaagtgo	tttgataaa	a acttttcc	a tgtgatcaci	ccggttctg	6360

taaatcggca taaatctgtt catactggct gacaattttt ttagaagaaa attcattttc 6420 aagcatetet attgeegeet ttgtaaaaeg attgettage tgtteatett etaaaatget 6480 categegegg getgttgegg cegtaacate acegacatee aceaaaaate egeteacatt 6540 gttttttata acctcaggga taccgccaat gtttgttcca atacaaggca ctccgcaagc 6600 catcgcttca agcaggacaa ggccaaagct ttcttttca gatagcagca gcttcaaatc 6660 gctaatagaa taaagatett caacaeggte ttgattteca agcattaaga ettggtette 6720 caagccatat tttctgataa gctcgcaggc tgtcgatttc tccggaccgt ctccgactaa 6780 aagcagette getttegttt tgecagegat attgeggaac acaeggatga cateetgeac 6840 gcgtgcagcc actggtaaca ggattagcag agcgaggtat gtaggcggtg ctacagagtt 6900 cttgaagtgg tggcctaact acggctacac tagaaggaca gtatttggta tctgcgctct 6960 getgaageea gttacetteg gaaaaagagt tggtagetet tgateeggea aacaaaceae 7020 cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc 7080 tcaagaagat cctttgatct tttctacggg gtctgacgct cagtggaacg aaaactcacg 7140 ttaagggatt ttggtcatga gattatcaaa aaggatcttc acctagatcc ttttaaatta 7200 aaaatgaagt tttaaatcaa tctaaagtat atatgagtaa acttggtctg acagttacca 7260 atgettaate agtgaggeae etateteage gatetgteta tttegtteat ceatagttge 7320 ctgactcccc gtcgtgtaga taactacgat acgggagggc ttaccatctg gccccagtgc 7380 tgcaatgata ccgcgagacc cacgctcacc ggctccagat ttatcagcaa taaaccagcc 7440 agccggaagg gccgagcgca gaagtggtcc tgcaacttta tccgcctcca tccagtctat 7500 taattgttgc cgggaagcta gagtaagtag ttcgccagtt aatagtttgc gcaacgttgt 7560 tgecattget acaggeateg tggtgteaeg etegtegttt ggtatggett catteagete 7620 cggttcccaa cgatcaaggc gagttacatg atcccccatg ttgtgcaaaa aagcggttag 7680 cteetteggt ceteegateg ttgteagaag taagttggee geagtgttat cacteatggt 7740 tatggcagca ctgcataatt ctcttactgt catgccatcc gtaagatgct tttctgtgac 7800 tggtgagtac tcaaccaagt cattctgaga atagtgtatg cggcgaccga gttgctcttg 7860 eceggegtea ataegggata ataeegegee acatageaga aetttaaaag tgeteateat 7920 tggaaaacgt tettegggge gaaaaetete aaggatetta eegetgttga gateeagtte 7980 gatgtaaccc actcgtgcac ccaactgatc ttcagcatct tttactttca ccagcgtttc 8040 tgggtgagca aaaacaġgaa ggcaaaatgc cgcaaaaaag ggaataaggg cgacacggaa 8100

atgttgaata ctcatactct teetttttea atattattga ageatttate agggttattg 8160 totcatgago ggatacatat ttgaatgtat ttagaaaaat aaacaaatag gggttoogog 8220 cacatttece egaaaagtge cacetgaegt etaagaaace attattatea tgacattaae 8280 ctataaaaat aggcgtatca cgaggccctt tcgtctcgca tgcggatcag tgagggtttg 8340 caactgcggg tcaaggatct ggatttcgat cacggcacga tcatcgtgcg ggagggcaag 8400 ggctccaagg atcgggcctt gatgttaccc gagagcttgg cacccagcct gcgcgagcag 8460 gggaattgat ccggtggatg accttttgaa tgacctttaa tagattatat tactaattaa 8520 ttggggaccc tagaggtccc ctttttatt ttaaaaattt tttcacaaaa cggtttacaa 8580 gcataacggg ttttgctgcc cgcaaacggg ctgttctggt gttgctagtt tgttatcaga 8640 atogoagato eggetteagg titgeegget gaaagegeta titetteeag aattgeeatg 8700 attttttccc cacgggaggc gtcactggct cccgtgttgt cggcagcttt gattcgataa 8760 gcagcatege etgtttcagg etgtetatgt gtgactgttg agetgtaaca agttgtetca 8820 ggtgttcaat ttcatgttct agttgctttg ttttactggt ttcacctgtt ctattaggtg 8880 ttacatgctg ttcatctgtt acattgtcga tctgttcatg gtgaacagct ttaaatgcac 8940 canaaactcg taaaagctct gatgtatcta tcttttttac accgttttca tctgtgcata 9000 tggacagitt tecetitgat atctaaeggi gaacagitgi tetaetitig titgitagie 9060 ttgatgette actgatagat acaagageea taagaacete agateettee gtatttagee 9120 agtatgttct ctagtgtggt tcgttgtttt tgcgtgagcc atgagaacga accattgaga 9180 tcatgcttac tttgcatgtc actcaaaaat tttgcctcaa aactggtgag ctgaattttt 9240 qcagttaaag catcgtgtag tgtttttctt agtccgttac gtaggtagga atctgatgta 9300 atggttgttg gtattttgtc accattcatt tttatctggt tgttctcaag ttcggttacg 9360 agatocattt gtotatotag ttoaacttgg aaaatcaacg tatoagtogg goggootogo 9420 ttatcaacca ccaatttcat attgctgtaa gtgtttaaat ctttacttat tggtttcaaa 9480 acccattggt taagcctttt aaactcatgg tagttatttt caagcattaa catgaactta 9540 aattcatcaa ggctaatctc tatatttgcc ttgtgagttt tcttttgtgt tagttctttt 9600 aataaccact cataaatcct catagagtat ttgttttcaa aagacttaac atgttccaga 9660 ttatatttta tgaatttttt taactggaaa agataaggca atatctcttc actaaaaact 9720 aattctaatt tttcgcttga gaacttggca tagtttgtcc actggaaaat ctcaaagcct 9780 ttaaccaaag gattootgat ttocacagtt otogtoatca gotototggt tgotttagot 9840

aatacaccat aagcattttc cctactgatg ttcatcatct gagcgtattg gttataagtg 9900 aacgataccg tccgttcttt ccttgtaggg ttttcaatcg tggggttgag tagtgccaca 9960 cagcataaaa ttagcttggt ttcatgctcc gttaagtcat agcgactaat cgctagttca 10020 tttgctttga aaacaactaa ttcagacata catctcaatt ggtctaggtg attttaatca 10080 ctataccaat tgagatgggc tagtcaatga taattactag tccttttcct ttgagttgtg 10140 ggtatctgta aattctgcta gacctttgct ggaaaacttg taaattctgc tagaccctct 10200 gtaaattccg ctagaccttt gtgtgttttt tttgtttata ttcaagtggt tataatttat 10260 agaataaaga aagaataaaa aaagataaaa agaatagatc ccagccctgt gtataactca 10320 ctactttagt cagttccgca gtattacaaa aggatgtcgc aaacgctgtt tgctcctcta 10380 caaaacagac cttaaaaccc taaaggctta agtagcaccc tcgcaagctc gggcaaatcg 10440 ctgaatattc cttttgtctc cgaccatcag gcacctgagt cgctgtcttt ttcgtgacat 10500 tcagttcgct gcgctcacgg ctctggcagt gaatgggggt aaatggcact acaggcgcct 10560 tttatggatt catgcaagga aactacccat aatacaagaa aagcccgtca cgggcttctc 10620 agggcgtttt atggcgggtc tgctatgtgg tgctatctga cttttcgctg ttcagcagtt 10680 cetgecetet gattitecag tetgaceaet teggattate eegtgacagg teatteagae 10740 tggctaatgc acccagtaag gcagcggtat catcaacagg cttacccgtc ttactgtcaa 10800 C 10801

<210> 77

<211> 8654

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN236 plasmid

<400> 77

ctcgaggcct acctagcttc caagaaagat atcctaacag cacaagagcy gaaagatgtt 60 ttgttctaca tccagaacaa cctctgctaa aattcctgaa aaattttgca aaaagttgtt 120 gactttatct acaaggtgtg gtataataat cttaacaaca gcaggacgct ctagaggagg 180 agacatcatg aaaattggaa ttatcggcgg aggctccgtt ggtcttttat gcgcctatta 240 tttgtcactt tatcacgacg tgactgttgt gacgaggcgg caagaacagg ctgcggccat 300 tcagtctgaa ggaatccggc tttataaagg cggggaggaa ttcagggctg attgcagtgc 360

ggacacgagt atcaattogg actttgacct gcttgtcgtg acagtgaagc agcatcagct 420 tcaatctgtt ttttcgtcgc ttgaacgaat cgggaagacg aatatattat ttttgcaaaa 480 cggcatgggg catatccacg acctaaaaga ctggcacgtt ggccattcca tttatgttgg 540 aatcgttgag cacggagctg taagaaaatc ggatacagct gttgatcata caggcctagg 600 tgcgataaaa tggagcgcgt tcgacgatgc tgaaccagac cggctgaaca tcttgtttca 660 gcataaccat teggatttte egatttatta tgagaeggat tggtaeegte tgetgaeggg 720 caagetgatt gtaaatgegt gtattaatee tttaaetgeg ttattgeaag tgaaaaatgg 780 agaactgctg acaacgccag cttatctggc ttttatgaag ctggtatttc aggaggcatg 840 ccgcatttta aaacttgaaa atgaagaaaa ggcttgggag cgggttcagg ccgtttgtgg 900 gcaaacgaaa gagaatcgtt catcaatgct ggttgacgtc attggaggcc ggcagacgga 960 agctgacgcc attatcggat acttattgaa ggaagcaagt cttcaaggtc ttgatgccgt 1020 ccacctagag tttttatatg gcagcatcaa agcattggag cgaaatacaa acaaagtctt 1080 ttgagetttt teggtaacat getataetea ttteggatea etaaetattt attggaggat 1140 cctgttttgg cggatgagag aagattttca gcctgataca gattaastca gaacgcagaa 1200 geggtetgat aaaacagaat ttgeetggeg geagtagege ggtggteeca «cetyacecca 1260 tgccgaactc agaagtgaaa cgccgtagcg ccgatggtag tgtggggtct ccccatgcga 1320 gagtagggaa ctgccaggca tcaaataaaa cgaaaggctc agtcgaaaga ctgggccttt 1380 cgttttatct gttgtttgtc ggtgaacgct ctcctgagta ggacaaatcc gccgggagcg 1440 gatttgaacg ttgcgaagca acggcccgga gggtggcggg caggacgccc gccataaact 1500 gccaggcatc aaattaagca gaaggccatc ctgacggatg gcctttttgc gtttctacaa 1560 actottggta cocagaaaaa goggcaaaag oggotgttaa aaaagogaaa togaagaago 1620 tgtctgccgc taagacggaa tatcaaaagc gttctgctgt tgtgtcatct ttaaaagtca 1680 cageegatga ateecageaa gatgteetaa aataettgaa caeecagaaa gataaaggaa 1740 atgcagacca aattcattet tattatgtgg tgaaegggat tgetgtteat geetcaaaag 1800 aggttatgga aaaagtggtg cagtttcccg aagtggaaaa ggtgcttcct aatgagaaac 1860 ggcagctttt taagtcatcc tececattta atatgaaaaa agcacagaaa getattaaag 1920 caactgacgg tgtggaatgg aatgtagacc aaatcgatgc cccaaaagct tgggcacttg 1980 gatatgatgg aactggcacg gttgttgcgt ccattgatac cggggtggaa tggaatcatc 2040 cggcattaaa agagaaatat cgcggatata atccggaaaa tcctaatgag cctgaaaatg 2100

aaatgaactg gtatgatgcc gtagcaggcg aggcaagccc ttatgatgat ttggctcatg 2160 gaacccacgt gacaggcacg atggtgggct ctgaacctga tggaacaaat caaatcggtg 2220 tagcacctgg cgcaaaatgg attgctgtta aagcgttctc tgaagatggc ggcactgatg 2280 ctgacatttt ggaagctggt gaatgggttt tagcaccaaa ggacgcggaa ggaaatcccc 2340 acceggaaat ggctcctgat gttgtcaata actcatgggg agggggctct ggacttgatg 2400 aatggtacag agacatggtc aatgcctggc gttcggccga tattttccct gagttttcag 2460 cggggaatac ggatctcttt attcccggcg ggcctggttc tatcgcaaat ccggcaaact 2520 atccagaatc gtttgcaact ggagcgactg agaattccaa ttccccatgg agagaaaaga 2580 aaatcgctaa tgttgattac tttgaacttc tgcatattct tgaatttaaa aaggctgaaa 2640 gagtaaaaga ttgtgctgaa atattagagt ataaacaaaa tcgtgaaaca ggcgaaagaa 2700 agttgtatcg agtgtggttt tgtaaatcca ggctttgtcc aatgtgcaac tggaggagag 2760 caatgaaaca tggcattcag tcacaaaagg ttgttgctga agttattaaa caaaagccaa 2820 cagttegttg gttgtttete acattaacag ttaaaaatgt ttatgatgge gaagaattaa 2880 ataagagttt gtcagatatg gctcaaggat ttcgccgaat gatgcaatat aaaaaaatta 2940 ataaaaatct tgttggtttt atgcgtgcaa cggaagtgac aataaataat aaagataatt 3000 cttataatca gcacatgcat gtattggtat gtgtggaacc aacttatttt aagaatacag 3060 aaaactacgt gaatcaaaaa caatggattc aattttggaa aaaggcaatg aaattagact 3120 atgatccaaa tgtaaaagtt caaatgattc gaccgaaaaa taaatataaa tcggatatac 3180 aatcggcaat tgacgaaact gcaaaatatc ctgtaaagga tacggatttt atgaccgatg 3240 atgaagaaaa gaatttgaaa cgtttgtctg atttggagga aggtttacac cgtaaaaggt 3300 taatctccta tggtggtttg ttaaaagaaa tacataaaaa attaaacctt gatgacacag 3360 aagaaggcga tttgattcat acagatgatg acgaaaaagc cgatgaagat ggattttcta 3420 ttattgcaat gtggaattgg gaacggaaaa attattttat taaagagtag ttcaacaaac 3480 gggccatatt gttgtataag tgatgaaata ctgaatttaa aacttagttt atatgtggta 3540 caattaaaag agggaagcgt atcattaacc ctataaacta cgtctgccct cattattgga 3660 gggtgaaatg tgaatacatc ctattcacaa tcgaatttac gacacaacca aattttaatt 3720 tggctttgca ttttatcttt ttttagcgta ttaaatgaaa tggttttgaa cgtctcatta 3780 cctgatattg caaatgattt taataaacca cctgcgagta caaactgggt gaacacagcc 3840

tttatgttaa ccttttccat tggaacagct gtatatggaa agctatctga tcaattaggc 3900 atcaaaaggt tactcctatt tggaattata ataaattgtt tcgggtcggt aattgggttt 3960 gttggccatt ctttcttttc cttacttatt atggctcgtt ttattcaagg ggctggtgca 4020 gctgcatttc cagcactcgt aatggttgta gttgcgcgct atattccaaa ggaaaatagg 4080 ggtaaagcat ttggtcttat tggatcgata gtagccatgg gagaaggagt cggtccagcg 4140 attggtggaa tgatageeca ttatatteat tggteetate ttetaeteat teetatgata 4200 acaattatca ctgttccgtt tcttatgaaa ttattaaaga aagaagtaag gataaaaggt 4260 cattttgata tcaaaggaat tatactaatg tctgtaggca ttgtatttt tatgttgttt 4320 acaacatcat atagcatttc ttttcttatc gttagcgtgc tgtcattcct gatatttgta 4380 aaacatatca ggaaagtaac agateetttt gttgateeeg gattagggaa aaatataeet 4440 tttatgattg gagttetttg tgggggaatt atatttggaa cagtagcagg gtttgtetet 4500 atggttcctt atatgatgaa agatgttcac cagctaagta ctgccgaaat cggaagtgta 4560 attattttcc ctggaacaat gagtgtcatt attttcggct acattggtgg gatacttgtt 4620 gatagaagag gtcctttata cgtgttaaac atcggagtta catttcttc tgttagcttt 4680 traactgctt cotttotttt agaaacaaca toatggttca tgacaattat aatogtattt 4740 gttttaggtg ggctttcgtt caccaaaaca gttatatcaa caattgtttc aagtagcttg 4800 aaacagcagg aagctggtgc tggaatgagt ttgcttaact ttaccagctt tttatcagag 4860 ggaacaggta ttgcaattgt aggtggttta ttatccatac ccttacttga tcaaaggttg 4920 ttacctatgg aagttgatca gtcaacttat ctgtatagta atttgttatt actttttca 4980 ggaatcattg tcattagttg gctggttacc ttgaatgtat ataaacattc tcaaagggat 5040 ttctaaatcg ttaagggatc aactttggga gagagttcaa aattgatcct ttttttataa 5100 cagttcgaag cggccgcaat tcttgaagac gaaagggcct cgtgatacgc ctattttat 5160 aggttaatgt catgataata atggtttett agaegteagg tggcaetttt eggggaaatg 5220 tgcgcggaac ccctatttgt ttatttttct aaatacattc aaatatgtat ccgctcatga 5280 gacaataacc ctgataaatg cttcaataat attgaaaaag gaagagtatg agtattcaac 5340 attteegtgt egecettatt ecettttttg eggeattttg eetteetgtt titgeteace 5400 cagaaacgct ggtgaaagta aaagatgctg aagatcagtt gggtgcacga gtgggttaca 5460 tegaactgga teteaacage ggtaagatee ttgagagttt tegeceegaa gaacgtttte 5520 caatgatgag cacttttaaa gttctgctat gtggcgcggt attatcccgt attgacgccg 5580

ggcaagagca acteggtege egcatacaet atteteagaa tgaettggtt gagtaeteae 5640 cagtcacaga aaagcatett aeggatggea tgacagtaag agaattatge agtgetgeea 5700 taaccatgag tgataacact gcggccaact tacttctgac aacgatcgga ggaccgaagg 5760 agctaaccgc ttttttgcac aacatggggg atcatgtaac tcgccttgat cgttgggaac 5820 eggagetgaa tgaagecata ecaaaegaeg agegtgaeae eaegatgeet geageaatgg 5880 caacaacgtt gcgcaaacta ttaactggcg aactacttac tctagcttcc cggcaacaat 5940 taatagactg gatggaggcg gataaagttg caggaccact tetgegeteg geeetteegg 6000 ctggctggtt tattgctgat aaatctggag ccggtgagcg tgggtctcgc ggtatcattg 6060 cagcactggg gccagatggt aagccctccc gtatcgtagt tatctacacg acggggagtc 6120 aggcaactat ggatgaacga aatagacaga tcgctgagat aggtgcctca ctgattaagc 6180 attggtaact gtcagaccaa gtttactcat atatacttta gattgattta aaacttcatt 6240 tttaatttaa aaggatctag gtgaagatcc tttttgataa tctcatgacc aaaatccctt 6300 aacgtgagtt ttcgttccac tgagcgtcag accccgtaga aaagatcaaa ggatcttctt 6360 gagateettt ttttetgege gtaatetget gettgeaaac aaaaaaacca eegetaceag 6420 eggtggtttg tttgeeggat caagagetae caactetttt teegaaggta aetggettea 6480 gcagagcgca gataccaaat actgtccttc tagtgtagcc gtagttaggc caccacttca 6540 agaactetgt ageacegeet acataceteg etetgetaat eetyttacea gtggetgetg 6600 ccagtggcga taagtcgtgt cttaccgggt tggactcaag acgatagtta ccggataagg 6660 cgcagcggtc gggctgaacg gggggttcgt gcacacagcc cagcttggag cgaacgacct 6720 acaccgaact gagataccta cagcgtgagc tatgagaaag cgccacgctt cccgaaggga 6780 gaaaggegga caggtateeg gtaageggea gggteggaae aggagagege acgagggage 6840 ttccaggggg aaacgcctgg tatctttata gtcctgtcgg gtttcgccac ctctgacttg 6900 agcgtcgatt tttgtgatgc tcgtcagggg ggcggagcct atggaaaaac gccagcaacg 6960 eggeettttt aeggtteetg geettttget ggeettttge teacatgtte ttteetgegt 7020 tateceetga ttetgtggat aacegtatta eegeetttga gtgagetgat aeegetegee 7080 gcagccgaac gaccgagcgc agcgagtcag tgagcgagga agcggaagag cgcctgatgc 7140 ggtattttct ccttacgcat ctgtgcggta tttcacaccg catatggtgc actctcagta 7200 caatctgctc tgatgccgca tagttaagcc agtatacact ccgctatcgc tacgtgactg 7260 ggtcatggct gcgccccgac acccgccaac acccgctgac gcgccctgac gggcttgtct 7320

geteceggea tecgettaca gacaagetgt gacegtetee gggagetgea tgtgteagag 7380 gttttcaccg tcatcaccga aacgcgcgag gcagctgcgg taaagctcat cagcgtggtc 7440 gtgaagcgat tcacagatgt ctgcctgttc atccgcgtcc agctcgttga gtttctccag 7500 aagcgttaat gtctggcttc tgataaagcg ggccatgtta agggcggttt tttcctgttt 7560 ggtcacttga tgcctccgtg taagggggaa tttctgttca tgggggtaat gataccgatg 7620 aaacgagaga ggatgctcac gatacgggtt actgatgatg aacatgcccg gttactggaa 7680 cgttgtgagg gtaaacaact ggcggtatgg atgcggcggg accagagaaa aatcactcag 7740 ggtcaatgcc agcgcttcgt taatacagat gtaggtgttc cacagggtag ccagcagcat 7800 cctgcgatgc agatccggaa cataatggtg cagggcgctg acttccgcgt ttccagactt 7860 tacgaaacac ggaaaccgaa gaccattcat gttgttgctc aggtcgcaga cgttttgcag 7920 cagcagtege tteaegtteg etegegtate ggtgatteat tetgetaace agtaaggeaa 7980 ccccgccagc ctagccgggt cctcaacgac aggagcacga tcatgcgcac ccgtggccag 8040 gacccaacgo tgcccgagat gcgccgcgtg cggctgctgg agatggcgga cgcgatggat 8100 atgitetgee aagggitiggi itgegeatte acagtietee geaagaatig attggeteea 3160 attettggag tggtgaatcc gttagcgagg tgccgccggc ttccattcag gtcgaggtgg 8220 cccggctcca tycaccgcga cgcaacgcgg ggaggcagac aaggtatagg gcggcgccta 8280 caatccatgo caaccogtto catgtgotog cogaggoggo ataaatogoo gtgacgatca 8340 gcggtccagt gatcgaagtt aggctggtaa gagccgcgag cgatccttga agctgtccct 8400 gatggtcgtc atctacctgc ctggacagca tggcctgcaa cgcgggcatc ccgatgccgc 8460 cggaagcgag aagaatcata atggggaagg ccatccagcc tcgcgtcggc ggccgcttcg 8520 tcgaccgaaa cagcagttat aaggcatgaa gctgtccggt ttttgcaaaa gtggctgtga 8580 ctgtaaaaag aaatcgaaaa agaccgtttt gtgtgaaaac ggtctttttg tttcctttta 8640 8654 accaactgcc ataa

<210> 78

<211> 8093

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
pAN423 plasmid

<400> 78 ggcggccgct tcgtcgaccg aaacagcagt tataaggcat gaagctgtcc ggtttttgca 60

aaagtggctg tgactgtaaa aagaaatcga aaaagaccgt tttgtgtgaa aacggtcttt 120 ttgtttcctt ttaaccaact gccataactc gaggcctacc tagcttccaa gaaagatatc 180 ctaacagcac aagageggaa agatgttttg ttetacatee agaacaacet etgetaaaat 240 tcctgaaaaa ttttgcaaaa agttgttgac tttatctaca aggtgtggta taataatctt 300 aacaacagca ggacgctcta gaaaaggagg aatttaaatg tatcgtacga tqatgaqcqq 360 caaacttcac agggcaactg ttacggaagc aaacctgaac tatgtgggaa gcattacaat 420 tgatgaagat ctcattgatg ctgtgggaat gcttcctaat gaaaaagtac aaattgtgaa 480 taataataat ggagcacgtc ttgaaacgta tattattcct ggtaaacggg gaagcggcgt 540 catatgetta aacggtgcag cegeacgeet tgtgcaggaa ggagataagg teattattat 600 ttcctacaaa atgatgtctg atcaagaagc ggcaagccat gagccgaaag tggctgttct 660 gaatgatcaa aacaaaattg aacaaatgct ggggaacgaa ccagcccgta caattttgta 720 aaggateetg tittggegga tgagagaaga titteageet gatacagatt aaateagaac 780 gcagaagcgg tctgataaaa cagaatttgc ctggcggcag tagcgcggtg gtcccacctg 840 accocatgee gaacteagaa gtgaaaegee gtagegeega tggtagtgtg gggteteece 900 atgcgagagt agggaactgc caggcatcaa ataaaacgaa aggctcagtc gaaagactgg 960 geetttegtt ttatetgttg tttgteggtg aacgetetee tgagtaggae aaateegeeg 1020 ggagcggatt tgaacgttgc gaagcaacgg cccggagggt ggcgggcagg acgcccqcca 1080 taaactgcca ggcatcaaat taagcagaag gccatcctga cggatggcct ttttgcgttt 1140 ctacaaactc ttggtaccca gaaaaagcgg caaaagcggc tgttaaaaaa gcgaaatcqa 1200 agaagetgte tgeegetaag aeggaatate aaaagegtte tgetgttgtg teatetttaa 1260 aagtcacage egatgaatee cageaagatg teetaaaata ettgaacace cagaaagata 1320 aaggaaatgc agaccaaatt cattcttatt atgtggtgaa cgggattgct gttcatqcct 1380 caaaagaggt tatggaaaaa gtggtgcagt ttcccgaagt ggaaaaggtg cttcctaatg 1440 agaaacggca gctttttaag tcatcctccc catttaatat gaaaaaagca cagaaagcta 1500 ttaaagcaac tgacggtgtg gaatggaatg tagaccaaat cgatgcccca aaagcttggg 1560 cacttggata tgatggaact ggcacggttg ttgcgtccat tgataccggg gtggaatgga 1620 atcatccggc attaaaagag aaatatcgcg gatataatcc ggaaaatcct aatgagcctg 1680 aaaatgaaat gaactggtat gatgccgtag caggcgaggc aagcccttat gatgatttgg 1740 ctcatggaac ccacgtgaca ggcacgatgg tgggctctga acctgatgga acaaatcaaa 1800 toggtgtage acctggcgca aaatggattg ctgttaaage gttetetgaa gatggcggca 1860 ctgatgctga cattttggaa gctggtgaat gggttttagc accaaaggac gcggaaggaa 1920 atococacco ggaaatggot cotgatgttg toaataacto atggggaggg ggototggac 1980 ttgatgaatg gtacagagac atggtcaatg cctggcgttc ggccgatatt ttccctgagt 2040 tttcagcggg gaatacggat ctctttattc ccggcgggcc tggttctatc gcaaatccgg 2100 caaactatcc agaatcgttt gcaactggag cgactgagaa ttccaattcc ccatggagag 2160 aaaagaaaat cgctaatgtt gattactttg aacttctgca tattcttgaa tttaaaaagg 2220 ctgaaagagt aaaagattgt gctgaaatat tagagtataa acaaaatcgt gaaacaggcg 2280 aaagaaagtt gtatcgagtg tggttttgta aatccaggct ttgtccaatg tgcaactgga 2340 ggagagcaat gaaacatggc attcagtcac aaaaggttgt tgctgaagtt attaaacaaa 2400 agccaacagt togttggttg tttctcacat taacagttaa aaatgtttat gatggcgaag 2460 aattaaataa gagttigica gataiggete aaggattieg eegaatgaig caatataaaa 2520 aaattaataa aaatcttgtt ggttttatgc gtgcaacgga agtgacaata aataataaag 2580 ataattetta taateageae atgeatgtat tggtatgtgt ggaaceaaet tattttaaga 2640 atacagaaaa ctacgtgaat caaaaacaat ggattcaatt ttggaaaaag gcaatgaaat 2700 tagactatga tocaaatgta aaagttoaaa tgattogaco gaaaaataaa tataaatogg 2760 atatacaatc ggcaattgac gaaactgcaa aatatcctgt aaaggatacg gattttatga 2820 ccgatgatga agaaaagaat ttgaaacgtt tgtctgattt ggaggaaggt ttacaccgta 2880 aaaggttaat ctcctatggt ggtttgttaa aagaaataca taaaaaatta aaccttgatg 2940 acacagaaga aggcgatttg attcatacag atgatgacga aaaagccgat gaagatggat 3000 tttctattat tgcaatgtgg aattgggaac ggaaaaatta ttttattaaa gagtagttca 3060 acaaacgggc catattgttg tataagtgat gaaatactga atttaaaact tagtttatat 3120 gtggtaaaat gttttaatca agtttaggag gaattaatta tgaagtgtaa tgaatgtaac 3180 agggttcaat taaaagaggg aagcgtatca ttaaccctat aaactacgtc tgccctcatt 3240 attggagggt gaaatgtgaa tacatcctat tcacaatcga atttacgaca caaccaaatt 3300 ttaatttggc tttgcatttt atctttttt agcgtattaa atgaaatggt tttgaacgtc 3360 tcattacctg atattgcaaa tgattttaat aaaccacctg cgagtacaaa ctgggtgaac 3420 acageettta tgttaacett tteeattgga acagetgtat atggaaaget atetgateaa 3480 ttaggcatca aaaggttact cctatttgga attataataa attgtttcgg gtcggtaatt 3540

gggtttgttg gccattcttt cttttcctta cttattatgg ctcgttttat tcaaggggct 3600 ggtgcagctg catttccagc actcgtaatg gttgtagttg cgcgctatat tccaaaggaa 3660 aataggggta aagcatttgg tcttattgga tcgatagtag ccatgggaga aggagtcggt 3720 ccagcgattg gtggaatgat agcccattat attcattggt cctatcttct actcattcct 3780 ' atgataacaa ttatcactgt tccgtttctt atgaaattat taaagaaaga'agtaaggata 3840 aaaggtcatt ttgatatcaa aggaattata ctaatgtctg taggcattgt atttttatg 3900 ttgtttacaa catcatatag catttcttt cttatcgtta gcgtgctgtc attcctgata 3960 tttgtaaaac atatcaggaa agtaacagat ccttttgttg atcccggatt agggaaaaat 4020 atacctttta tgattggagt tctttgtggg ggaattatat ttggaacagt agcagggttt 4080 gtctctatgg ttccttatat gatgaaagat gttcaccagc taagtactgc cgaaatcgga 4140 agtgtaatta ttttccctgg aacaatgagt gtcattattt tcggctacat tggtgggata 4200 cttgttgata gaagaggtcc tttatacgtg ttaaacatcg gagttacatt tctttctgtt 4260 agetttttaa etgetteett tettttagaa acaacateat ggtteatgae aattataate 4320 gtatttgttt taggtgggct ttcgttcacc aaaacagtta tatcaacaat tgtttcaagt 4380 agettgaaac agcaggaage tggtgetgga atgagtttge ttaaetttae cagettttta 4440 tcagagggaa caggtattgc aattgtaggt ggtttattat ccataccctt acttgatcaa 4500 aggitightac ctatggaagt tgatcagtca acttatctgt atagtaattt gitattacit 4560 ttttcaggaa tcattgtcat tagttggctg gttaccttga atgtatataa acattctcaa 4620 agggatttct aaatcgttaa gggatcaact ttgggagaga gttcaaaatt gatccttttt 4680 ttataacagt tcgaagcggc cgcaattctt gaagacgaaa gggcctcgtg atacgcctat 4740 ttttataggt taatgtcatg ataataatgg tttcttagac gtcaggtggc acttttcggg 4800 gaaatgtgcg cggaacccct atttgtttat ttttctaaat acattcaaat atgtatccgc 4860 tcatgagaca ataaccctga taaatgcttc aataatattg aaaaaggaag agtatgagta 4920 ttcaacattt ccgtgtcgcc cttattccct tttttgcggc attttgcctt cctgtttttg 4980 ctcacccaga aacgctggtg aaagtaaaag atgctgaaga tcagttgggt gcacgagtgg 5040 gttacatcga actggatctc aacagcggta agatccttga gagttttcgc cccgaagaac 5100 gttttccaat gatgagcact tttaaagttc tgctatgtgg cgcggtatta tcccgtattg 5160 acgccgggca agagcaactc ggtcgccgca tacactattc tcagaatgac ttggttgagt 5220 actcaccagt cacagaaaag catcttacgg atggcatgac agtaagagaa ttatgcagtg 5280

ctgccataac catgagtgat aacactgcgg ccaacttact tctgacaacg atcggaggac 5340 cqaaqqagct aaccgctttt ttgcacaaca tgggggatca tgtaactcgc cttgatcgtt 5400 gggaaccgga gctgaatgaa gccataccaa acgacgagcg tgacaccacg atgcctgcag 5460 caatqqcaac aacqttqcqc aaactattaa ctqqcqaact acttactcta gcttcccqqc 5520 aacaattaat agactggatg gaggcggata aagttgcagg accacttctg cgctcggccc 5580 ttccggctgg ctggtttatt gctgataaat ctggagccgg tgagcgtggg tctcgcggta 5640 tcattgcagc actggggcca gatggtaagc octoocgtat ogtagttate tacacgacqg 5700 qqaqtcaqqc aactatgqat gaacqaaata gacagatcqc tgayataggt gcctcactga 5760 ttaagcattg gtaactgtca gaccaagttt actcatatat actttagatt gatttaaaac 5820 ttcattttta atttaaaagg atctaggtga agatcctttt tgataatctc atgaccaaaa 5880 tocottaacg tgagttttcg ttocactgag cgtcagaccc cgtagaaaag atcaaaggat 5940 cttcttgaga tcctttttt ctgcgcgtaa tctgctgctt gcaaacaaaa aaaccaccgc 6000 taccagoggt ggtttgtttg coggatcaag agctaccaac tottttccg aaggtaactg 6060 getteageag agegeagata ceaaataetg teettetagt gtageegtag ttaggeeace 6120 acticaagaa ctctgtagca ccgcctacat acctcgctct gctaatcctg ttaccagigg 6180 ctgctgccag tggcgataag tcgtgtctta ccgggttyga ctcaagacga tagitaccgg 6240 ataaggegea geggteggge tgaaeggggg gttegtgeae acageecage ttggagegaa 6300 cqacctacac cqaactgaqa tacctacagc gtgagctatg agaaagcgcc acgcttcccg 6360 aagggagaaa ggcggacagg tatccggtaa gcggcagggt cggaacagga gagcgcacga 6420 qqqaqcttcc aqqqqqaaac gcctqqtatc tttataqtcc tqtcqqqttt cqccacctct 6480 qacttgaqcg tcgatttttg tgatgctcgt caggggggcg gagcctatgg aaaaacgcca 6540 qcaacgcqqc ctttttacqq ttcctqqcct tttqctqqcc ttttqctcac atqttctttc 6600 ctgcgttatc ccctgattct gtggataacc gtattaccgc ctttgagtga gctgataccg 6660 ctcgccgcag ccgaacgacc gagcgcagcg agtcagtgag cgaggaagcg gaagagcgcc 6720 tgatgcggta titteteett aegeateigt geggtatite acaeegeata tggigeacie 6780 tragtaraat regreetgat grogratage taagroagta taractrogr tatrgetarg 6840 tgactgggtc atggctgcgc cccgacaccc gccaacaccc gctgacgcgc cctgacgggc 6900 ttqtctqctc ccqqcatccq cttacaqaca agctqtqacc qtctccqqqa qctqcatqtq 6960 tcagaggttt tcaccqtcat caccqaaacq cqcqaqqcaq ctqcqqtaaa qctcatcaqc 7020

gtggtcgtga agcgattcac agatgtctgc ctgttcatcc gcgtccagct cgttgagttt 7080 ctccagaagc gttaatgtct ggcttctgat aaagcgggcc atgttaaggg cggtttttc 7140 ctgtttggtc acttgatgcc tccgtgtaag ggggaatttc tgttcatggg ggtaatgata 7200 ccgatgaaac gagagaggat gctcacgata cgggttactg atgatgaaca tgcccggtta 7260 ctggaacgtt gtgagggtaa acaactggcg gtatggatgc ggcgggacca gagaaaaatc 7320 actcagggtc aatgccagcg cttcgttaat acagatgtag gtgttccaca gggtagccag 7380 cagcatectg egatgeagat ceggaacata atggtgeagg gegetgaett eegegtttee 7440 agactttacg aaacacggaa accgaagacc attcatgttg ttgctcaggt cgcagacgtt 7500 ttgcagcagc agtcgcttca cgttcgctcg cgtatcggtg attcattctg ctaaccagta 7560 aggcaacccc gccagcctag ccgggtcctc aacgacagga gcacgatcat gcgcacccgt 7620 ggccaggacc caacgctgcc cgagatgcgc cgcgtgcggc tgctggagat ggcggacgcg 7680 atggatatgt tctgccaagg gttggtttgc gcattcacag ttctccgcaa gaattgattg 7740 gctccaattc ttggagtggt gaatccgtta gcgaggtgcc gccggcttcc attcaggtcg 7800 aggtggcccg gctccatgca ccgcgacgca acgcggggag gcagacaagg tatagggcgg 7860 egectacaat ecatgecaac cegttecatg tgetegeega ggeggeataa ategeegtga 7920 cgatcagcgg tccagtgatc gaagttaggc tggtaagagc cgcgagcgat ccttgaagct 7980 gtccctgatg gtcgtcatct acctgcctgg acagcatggc ctgcaacgcg ggcatcccga 8040 tgccgccgga agcgagaaga atcataatgg ggaaggccat ccagcctcgc gtc 8093

<210> 79

<211> 8098

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN429 plasmid

<400> 79

ggcggccgct tcgtcgaccg aaacagcagt tataaggcat gaagctgtcc ggttttgca 60
aaagtggctg tgactgtaaa aagaaatcga aaaagaccgt tttgtgtgaa aacggtcttt 120
ttgtttcctt ttaaccaact gccataactc gaggcctacc tagcttccaa gaaagatatc 180
ctaacagcac aagagcggaa agatgttttg ttctacatcc agaacaacct ctgctaaaat 240
tcctgaaaaa ttttgcaaaa agttgttgac tttatctaca aggtgtggta taataatctt 300

acaacagca	ggacgctcta	gattagaaag	gaggtttaat	taatgtatcg	Cacgatgatg	3.00
gcggcaaac	ttcacagggc	aactgttacg	gaagcaaacc	tgaactatgt	gggaagcatt	420
acaattgatg	aagatctcat	tgatgctgtg	ggaatgcttc	ctaatgaaaa	agtacaaatt	480
gtgaataata	ataatggagc	acgtcttgaa	acgtatatta	t-t-cctggtaa	acggggaagc	540
ggcgtcatat	gcttaaacgg	tgcagccgca	cgccttgtgc	aggaaggaga	taaggtcatt	6 00 .
attatttcct	acaaaatgat	gtctgatcaa	gaagcggcaa	gccatgagcc	gaaagtggct	660
gttctgaatg	atcaaaacaa	aattgaacaa	atgctgggga	acgaaccagc	ccgtacaatt	720
ttgtaaagga	tcctgttttg	gcggatgaga	gaagattttc	agcctgatac	agattaaatc	780
agaacgcaga	agcggtctga	taaaacagaa	tttgcctggc	ggcagtagcg	cggtggtccc	840
acctgacccc	atgccgaact	cagaagtgaa	acgccgtago	gccgatggta	gtgtggggtc	900
tccccatgcg	agagtaggga	actgccaggc	atcaaataaa	acgaaaggct	cagtcgaaag	960
actgggcctt	tcgttttatc	: tgttgtttgt	cggtgaacg	c tctcctgagt	aggacaaatc	1020
cyccyggago	ggatttgaac	gttgcgaago	: aacggcccg	g agggtggcgg	gcaggacgco	1080
cgccataaac	tgccaggcat	caaattaago	agaaggcca	t cctgacggat	ggccttttg	1140
cgtttctaca	a aactcttggt	acccagaaaa	a agcggcaaa	a gcggctgtta	a .aaaaagcgaa	1200
atcgaagaa	g ctgtctgcc	g ctaagacgga	a atatcaaaa	g cgttctgctq	g tegtgtcate	1260
tttaaaagt	c acageegate	g aatcccagca	a agatgtcct	a aaatactty	a acacccagaa	1320
agataaagg	a aatgcagac	c aaattcatte	c ttattatgt	g gtgaacggg	a ttgctgttca	1380
tgcctcaaa	a gaggttatg	g aaaaagtgg	t gcagtttcc	c gaagtggaa	a aggtgcttco	1440
					a aagcacagaa	
					g ccccaaaag	
					a ccggggtgg	
					a atcctaatg	
					c cttatgatg	
					g atggaacaa	
		•			t ctgaagatg	
cggcactga	at gctgacatt	t tggaagctg	gg tgaatggg	tt ttagcacca	a agga cg cgg	a 1920
aggaaatco	ce cacceggaa	a tggctcctg	ga tgttgtca	at aactcatgo	gg gagggggct	c 1980
tagactta	at gaatggtag	a gagacatgo	t caatgoot	gg cgttcggc	g atattttcc	c 2046

tgagttttca gcggggaata cggatctctt tattcccggc gggcctggtt ctatcgcaaa 2100 teeggeaaac tateeagaat egittgeaac tggagegaet gagaatteea atteeceatg 2160 gagagaaaag aaaatcgcta atgttgatta ctttgaactt ctgcatattc ttgaatttaa 2220 aaaggctgaa agagtaaaag attgtgctga aatattagag tataaacaaa atcgtgaaac 2280 aggcgaaaga aagttgtatc gagtgtggtt ttgtaaatcc aggctttgtc caatgtgcaa 2340 ctggaggaga gcaatgaaac atggcattca gtcacaaaag gttgttgctg aagttattaa 2400 acaaaagcca acagttcgtt ggttgtttct cacattaaca gttaaaaatg tttatgatgg 2460 cgaagaatta aataagagtt tgtcagatat ggctcaagga tttcgccgaa tgatgcaata 2520 taaaaaaatt aataaaaatc ttgttggttt tatgcgtgca acggaagtga caataaataa 2580 taaagataat tottataato agoacatgoa tgtattggta tgtgtggaac caacttattt 2640 taagaataca gaaaactacg tgaatcaaaa acaatggatt caattttgga aaaaggcaat 2700 gaaattagac tatgatccaa atgtaaaagt tcaaatgatt cgaccgaaaa ataaatataa 2760 atcggatata caatcggcaa ttgacgaaac tgcaaaatat cctgtaaagg atacggattt 2820 tatgaccgat gatgaagaaa agaatttgaa acgtttgtct gatttggagg aaggtttaca 2880 ccgtaaaagg ttaatctcct atggtggttt gttaaaagaa atacataaaa aattaaacct 2940 tgatgacaca gaagaaggcg atttgattca tacagatgat gacgaaaaag ccgatgaaga 3000 tggattttct attattgcaa tgtggaattg ggaacggaaa aattatttta ttaaagagta 3060 gttcaacaaa cgggccatat tgttgtataa gtgatgaaat actgaattta aaacttagtt 3120 tatatgtggt aaaatgtttt aatcaagttt aggaggaatt aattatgaag tgtaatgaat 3180 gtaacagggt tcaattaaaa gagggaagcg tatcattaac cctataaact acgtctgccc 3240 tcattattgg agggtgaaat gtgaatacat cctattcaca atcgaattta cgacacaacc 3300 aaattttaat ttggctttgc attttatctt tttttagcgt attaaatgaa atggttttga 3360 acgtctcatt acctgatatt gcaaatgatt ttaataaacc acctgcgagt acaaactggg 3420 tgaacacagc ctttatgtta accttttcca ttggaacagc tgtatatgga aagctatctg 3480 atcaattagg catcaaaagg ttactcctat ttggaattat aataaattgt ttcgggtcgg 3540 taattgggtt tgttggccat tctttctttt ccttacttat tatggctcgt tttattcaag 3600 gggctggtgc agctgcattt ccagcactcg taatggttgt agttgcgcgc tatattccaa 3660 aggaaaatag gggtaaagca tttggtctta ttggatcgat agtagccatg ggagaaggag 3720 tcggtccagc gattggtgga atgatagccc attatattca ttggtcctat cttctactca 3780

DISCOURT - WAY 012177282 1

ttoctatgat aacaattato actgttoogt ttottatgaa attattaaag aaagaagtaa 3840 ggataaaagg tcattttgat atcaaaggaa ttatactaat gtctgtaggc attgtatttt 3900 ttatgttgtt tacaacatca tatagcattt cttttcttat cgttagcgtg ctgtcattcc 3960 tgatatttgt aaaacatatc aggaaagtaa cagatccttt tgttgatccc ggattaggga 4020 aaaatatacc ttttatgatt ggagttcttt gtgggggaat tatatttgga acagtagcag 4080 ggtttgtctc tatggttcct tatatgatga aagatgttca ccagctaagt actgccgaaa 4140 teggaagtgt aattatttte eetggaacaa tgagtgteat tattttegge tacattggtg 4200 ggatacttgt tgatagaaga ggtcctttat acgtgttaaa catcggagtt acatttcttt 4260 ctgttagctt tttaactgct tcctttcttt tagaaacaac atcatggttc atgacaatta 4320 taatcgtatt tgttttaggt gggctttcgt tcaccaaaac agttatatca acaattgttt 4380 caagtagett gaaacageag gaagetggtg etggaatgag tttgettaae tttaccaget 4440 ttttatcaga gggaacaggt attgcaattg taggtggttt attatccata eccttacttg 4500 atcaaaggtt gttacctatg gaagttgatc agtcaactta tetgtatagt aatttgttat 4560 tacttttttc aggaatcatt gtcattagtt ggctggttac cttgaatgta tataaacatt 4620 ctcaaaggga tttctaaatc gttaagggat caactttggg agagagttca aaattgatcc 4680 tttttttata acagttogaa goggoogoaa ttottgaaga ogaaagggoo togtgataog 4740 cctattttta taggttaatg tcatgataat aatggtttct tagacgtcag gtggcacttt 4800 teggggaaat gtgegeggaa eccetatteg tetatette taaatacatt caaatatgta 4860 tecgeteatg agacaataac eetgataaat getteaataa tattgaaaaa ggaagagtat 4920 gagtattcaa cattteegtg tegecettat teeetttttt geggeatttt geetteetgt 4980 ttttgctcac ccagaaacgc tggtgaaagt aaaagatgct gaagatcagt tgggtgcacg 5040 agtgggttac atcgaactgg atctcaacag cggtaagatc cttgagagtt ttcgccccga 5100 agaacgtttt ccaatgatga gcacttttaa agttctgcta tgtggcgcgg tattatcccg 5160 tattgacgcc gggcaagagc aactcggtcg ccgcatacac tattctcaga atgacttggt 5220 tgagtactca ccagtcacag aaaagcatct tacggatggc atgacagtaa gagaattatg 5280 cagtgctgcc ataaccatga gtgataacac tgcggccaac ttacttctga caacgatcgg 5340 aggaccgaag gagctaaccg cttttttgca caacatgggg gatcatgtaa ctcgccttga 5400 tegttgggaa eeggagetga atgaageeat accaaacgae gagegtgaca eeaegatgee 5460 tgcagcaatg gcaacaacgt tgcgcaaact attaactggc gaactactta ctctagcttc 5520

ccggcaacaa ttaatagact ggatggaggc ggataaagtt gcaggaccac ttctgcgctc 5580 ggcccttccg gctggctggt ttattgctga taaatctgga gccggtgagc gtgggtctcg 5640 cggtatcatt gcagcactgg ggccagatgg taagccctcc cgtatcgtag ttatctacac 5700 gacggggagt caggcaacta tggatgaacg aaatagacag atcgctgaga taggtgcctc 5160 actgattaag cattggtaac tgtcagacca agtttactca tatatacttt agattgattt 5820 aaaacttcat ttttaattta aaaggatcta ggtgaagatc ctttttgata atctcatgac 5880 caaaatccct taacgtgagt tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa 5940 aggatettet tgagateett tttttetgeg egtaatetge tgettgeaaa caaaaaaace 6000 accgctacca gcggtggttt gtttgccgga tcaagagcta ccaactcttt ttccgaaggt 6060 aactggcttc agcagagcgc agataccaaa tactgtcctt ctagtgtagc cgtagttagg 6120 ccaccactte aagaactetg tageacegee tacatacete getetgetaa teetgttace 6180 agtggctgct gccagtggcg ataagtcgtg tcttaccggg ttggactcaa gacgatagtt 6240 accggataag gegeageggt egggetgaae ggggggtteg tgeacacage ceagettgga 6300 gcgaacgacc tacaccgaac tgagatacct acagcgtgag ctatgagaaa gcgccacgct 6360 tecegaaggg agaaaggegg acaggtatee ggtaagegge agggteggaa caggagageg 6420 cacgagggag cttccagggg gaaacgcctg gtatctttat agtcctgtcg ggtttcgcca 6480 cctctyactt gagcgtcgat ttttgtgatg ctcgtcaggg gggcggagcc tatggaaaaa 6540 cgccagcaac gcggcctttt tacggttcct ggccttttgc tggccttttg ctcacatgtt 6600 ctttcctgcg ttatcccctg attctgtgga taaccgtatt accgcctttg agtgagctga 6660 taccgctcgc cgcagccgaa cgaccgagcg cagcgagtca gtgagcgagg aagcggaaga 6720 gcgcctgatg cggtattttc tccttacgca tctgtgcggt atttcacacc gcatatggtg 6780 cacteteagt acaatetget etgatgeege atagttaage cagtatacae teegetateg 6840 ctacgtgact gggtcatggc tgcgccccga cacccgccaa cacccgctga cgcgccctga 6900 cgggcttgtc tgctcccggc atccgcttac agacaagctg tgaccgtctc cgggagctgc 6960 atgtgtcaga ggttttcacc gtcatcaccg aaacgcgcga ggcagctgcg gtaaagctca 7020 tcagcgtggt cgtgaagcga ttcacagatg tctgcctgtt catccgcgtc cagctcgttg 7080 agtttctcca gaagcgttaa tgtctggctt ctgataaagc gggccatgtt aagggcggtt 7140 ttttcctgtt tggtcacttg atgcctccgt gtaaggggga atttctgttc atgggggtaa 7200 tgataccgat gaaacgagag aggatgctca cgatacgggt tactgatgat gaacatgccc 7260

agattactgga acgttgtgag ggtaaacaac tggcggtatg gatgcggcgg gaccagagaa 7320

aaatcactca gggtcaatgc cagcgcttcg ttaatacaga tgtaggtgtt ccacagggta 7380

gccagcagca tcctgcgatg cagatccgga acataatggt gcagggcgct gacttccgcg 7440

tttccagact ttacgaaaca cggaaaccga agaccattca tgttgttgct caggtcgcag 7500

acgttttgca gcagcagtcg cttcacgttc gctcgcgtat cggtgattca ttctgctaac 7560

cagtaaggca accccggcag cctagccggg tcctcaacga caggagcacg atcatgcgca 7620

cccgtggcca ggacccaacg ctgcccgaga tgcgccgct gcggctgctg gagatggcgg 7680

acgcgatgga tatgttctgc caagggttgg tttgcgcatt cacagttctc cgcaagaatt 7740

gattggctcc aattcttgga gtggtgaatc cgttagcgag gtgccgccgg cttccattca 7800

ggtcgaggtg gcccggctcc atgcaccgc acgcaacgcg gggaggcaga caaggtatag 7860

ggcggcgcct acaatccatg ccaacccgtt ccatgtgctc gccgaggcgg cataaatcgc 7920

cgtgacgatc agcggtccag tgatcgaagt taggctggta agagccgca acgcaccttg 7980

aagctgtccc tgatggtcgt catctacctg cctggacagc atggcctgca acgcgggcat 8040

cccgatgccg ccggaagcga gaagaatcat aatggggaag gccatccagc ctcgcgtc 8098

<400> 80
ccatcgaatg gecagatgat taattectaa tttttgttga cactetatea ttgatagagt 60

tattttacca ctccctatea gtgatagaga aaagtgaaat gaatagteeg acaaaaatet 120
agattagaaa ggaggattta aatatgagae agattactga tatttcacag ctgaaagaag 180

ccataaaaaca ataccattea gagggcaagt caateggatt tgttccgacg atggggttte 240

tgcatgaggg gcatttaace ttagcagaca aagcaagaca agaaaacgae gccgttatta 300

tgagtatttt tgtgaateet gcacaatteg gccctaatga agattttgaa gcatateege 360

gcgatattga gcgggatgea gctcttgcag aaaacgccgg agtcgatatt cttttacge 420

cagatgctca tgatatgtat cccggtgaaa agaatgtcae gattcatgta gaaagacgca 480

cagacgtgtt atgcgggcge tcaagagaag gacattttga cggggtcgcg atcgtactga 540

cgaagctttt caatctagte aagccgacte gtgcctattt cggtttaaaa gatgcgcage 600

<210> 80

<211> 4450

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN443 plasmid

aggtagetgt tgttgatggg ttaateageg aettetteat ggatattgaa ttggtteetg 660 tcgatacggt cagagagaa gacggcttag ccaaaagctc tcgcaatgta tacttaacag 720 ctgaggaaag aaaagaagcg cctaagctgt atcgggccct tcaaacaagt gcggaacttg 780 tccaagccgg tgaaagagat cctgaagcgg tgataaaagc tgcaaaagat atcattgaaa 840 cgactagcgg aaccatagac tatgtagagc tttattccta tccggaactc gagcctgtga 900 atgaaattgc tggaaagatg attctcgctg ttgcagttgc tttttcaaaa gcgcgtttaa 960 tagataatat cattattgat attcgtagaa aggaggtgaa ttaatatgta tcgtacgatg 1020 atgageggea aaetteaeag ggeaaetgtt aeggaageaa aeetgaaeta tgtgggaage 1080 attacaattg atgaagatct cattgatgct gtgggaatgc ttcctaatga aaaagtacaa 1140 attgtgaata ataataatgg agcacgtctt gaaacgtata ttattcctgg taaacgggga 1200 agcggcgtca tatgcttaaa cggtgcagcc gcacgccttg tgcaggaagg agataaggtc 1260 attattattt cctacaaaat gatgtctgat caagaagcgg caagccatga gccgaaagtg 1320 gctgttctga atgatcaaaa caaaattgaa caaatgctgg ggaacgaacc agcccgtaca 1380 attttgtaaa ggatcccccg gggatccctc gaggtcgacc tgcaggggga ccatggtctc 1440 agegettgga gecaccegea gttegaaaaa taataagett gaeetgtgaa gtgaaaaatg 1500 gegeacattg tgegacattt tttttgtetg eegtttaeeg etaetgegte aeggatetee 1560 acgcgccctg tagcggcgca ttaagcgcgg cgggtgtggt ggttacgcgc agcgtgaccg 1620 cgttcgccgg ctttccccgt caagctctaa atcgggggct ccctttaggg ttccgattta 1740 gtgctttacg gcacctcgac cccaaaaaac ttgattaggg tgatggttca cgtagtgggc 1800 categeeetg atagaeggtt tttegeeett tgaegttgga gteeaegtte tttaatagtg 1860 gactettgtt ccaaactgga acaacactca accetatete ggtetattet tttgatttat 1920 aagggatttt gccgatttcg gcctattggt taaaaaatga gctgatttaa caaaaattta 1980 acgcgaattt taacaaaata ttaacgctta caatttcagg tggcactttt cggggaaatg 2040 tgcgcggaac ccctatttgt ttattttct aaatacattc aaatatgtat ccgctcatga 2100 gacaataacc ctgataaatg cttcaataat attgaaaaag gaagagtatg agtattcaac 2160 atttccgtgt cgcccttatt cccttttttg cggcattttg ccttcctgtt tttgctcacc 2220 cagaaacgct ggtgaaagta aaagatgctg aagatcagtt gggtgcacga gtgggttaca 2280 tcgaactgga tctcaacagc ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc 2340

caatgatgag cacttttaaa gttctgctat gtggcgcggt attatcccgt attgacgccg 2400 qqcaaqaqca actcqqtcqc cqcatacact attctcaqaa tqacttqqtt qaqtactcac 2460 cagtcacaga aaagcatctt acggatggca tgacagtaag agaattatgc agtgctgcca 2520 taaccatgag tgataacact geggecaact tacttetgac aacgategga ggacegaagg 2580 agctaaccgc ttttttgcac aacatggggg atcatgtaac tcgccttgat cgttgggaac 2640 cggagctgaa tgaagccata ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg 2700 caacaacgtt gcgcaaacta ttaactggcg aactacttac tctagcttcc cggcaacaat 2760 tgatagactg gatggaggcg gataaagttg caggaccact tetgegeteg gecetteegg 2820 ctggctggtt tattgctgat aaatctggag ccggtgagcg tggctctcgc ggtatcattg 2880 cagcactggg gccagatggt aagccctccc gtatcgtagt tatctacacg acggggagtc 2940 aggcaactat ggatgaacga aatagacaga togotgagat aggtgootca otgattaago 3000 attggtagga attaatgatg totogtttag ataaaagtaa agtgattaac agcgcattag 3060 agetgettaa tgaggtegga ategaaggtt taacaaceeg taaactegee cagaagetag 3120 gtgtagagca gcctacattg tattggcatg taaaaaaataa gcgggctttg ctcgacgcct 3180 tagccattga gatgttagat aggcaccata ctcacttttg ccctttagaa ggggaaagct 3240 ggcaagattt tttacgtaat aacgctaaaa gttttagatg tgctttacta agtcatcgcg 3300 atggagcaaa agtacattta ggtacacggc ctacagaaaa acagtatgaa actctcgaaa 3360 atcaattago otttttatgo caacaaggtt tttcactaga gaatgoatta tatgoactoa 3420 gcgcagtggg gcattttact ttaggttgcg tattggaaga tcaagagcat caagtcgcta 3480 aagaagaaag ggaaacacct actactgata gtatgccgcc attattacga caagctatcg 3540 aattatttga tcaccaaggt gcagagccag cettettatt eggeettgaa ttgatcatat 3600 gcggattaga aaaacaactt aaatgtgaaa gtgggtctta aaagcagcat aacctttttc 3660 cgtgatggta acttcactag tttaaaagga tctaggtgaa gatccttttt gataatctca 3720 tgaccaaaat cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga 3780 tcaaaggate ttettgagat eettttttte tgegegtaat etgetgettg caaacaaaaa 3840 aaccaccgct accagcggtg gtttgtttgc cggatcaaga gctaccaact ctttttccga 3900 aggtaactgg cttcagcaga gcgcagatac caaatactgt ccttctagtg tagccgtagt 3960 taggocacca etteaagaae tetgtageae egectacata ectegetetg etaateetgt 4020 taccagtggc tgctgccagt ggcgataagt cgtgtcttac cgggttggac tcaagacgat 4080 agttaccgga taaggcgaag cggtcgggct gaacgggggg ttcgtgcaca cagcccagct 4140 tggagcgaac gacctacacc gaactgagat acctacagcg tgagctatga gaaagcgcca 4200 cgcttcccga agggagaaag gcggacaggt atccggtaag cggcagggtc ggaacaggag 4260 agcgaccgag ggagcttcca gggggaaacg cctggtatct ttatagtcct gtcgggttc 4320 gccacctctg acttgagcgt cgatttttgt gatgctcgtc aggggggggg agcctatgga 4380 aaaacgccag caacgcggcc ttttacggt tcctggcctt ttgctggcct tttgctcaca 4440 tgacccgaca

<210> 81

<211> 10212

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
pAN251 plasmid

<400> 81 gcggccgcta aaaagagctt gaggatttgc ggagtgaaaa tcagacattg cggaatcagc 60 tagagatgac agaagaggat tacaaggcac tgatcgatat catggatcgg gccagaaaaa 120 tggttgtttc gaaggaagac ggaagaatga aaaaagcggc tcaagaaacg taaagaaacg 180 cctgaaatga accggcccta tagtaagaat aggccggttg ttttgatttc tatgcagact 240 ctcccggtgt catttcgcga tccatatcag gatgccagat gagcgggtct tcccctttgt 300 cccgcgccat atcatactta acagttttaa agttcatttt gttccaaaat tccgctgatt 360 tcattctcgg atttgtccgg atcggcattt tgaatgattt tgcaaattca accaaggctc 420 tecegtatee cetgttetgg tageceggaa gaaceteaag ettecacage tecaaataat 480 cctggcggtt gtcaaaatag ggattcgatt tgccgttaac ttgatacaga ctcattcgtg 540 ctacaagttt atcgccaaaa taaatcccgt aaaaaggcga ggtgctgtca ttttcaataa 600 tattatcctg aagttcttca agcattgaaa gctcctgaat gccgtattct ttgaatttct 660 tgaattette cagegtttta tagttgataa geagaegtte taeetttgte aaacaaatet 720 ccccctttgt tgtttctaca tatattgtaa acgctttatt taaaaaatcc aaatatttaa 780 actttaattt taagcacatg ggatctttga gaagtaattt cttcttactt ctgctatgat 840 aatacgtaaa tgcgtcgacc gaaacagcag ttataaggca tgaagctgtc cggtttttgc 900

aaaagtggct gtgactgtaa aaagaaatcg aaaaagaccg ttttgtgtga aaacggtctt 960

tttgtttcct tttaaccaac tgccataact cgaggcctac ctagcttcca agaaagatat 1020 cctaacagca caagagcgga aagatgtttt gttctacatc cagaacaacc tctgctaaaa 1080 ttcctgaaaa attttgcaaa aagttgttga ctttatctac aaggtgtggt ataataatct 1140 taacaacagc aggacgctct agaggaggag acatcatgaa aattggaatt ateggcggag 1200 geteegttgg tettttatge geetattatt tgteaettta teaegaegtg aetgttgtga 1260 cgaggcggca agaacaggct gcggccattc agtctgaagg aatccggctt tataaaggcg 1320 gggaggaatt cagggctgat tgcagtgcgg acacgagtat caattcggac tttgacctgc 1380 ttgtcgtgac agtgaagcag catcagcttc aatctgtttt ttcgtcgctt gaacgaatcg 1440 ggaagacgaa tatattattt ttgcaaaacg gcatggggca tatccacgac ctaaaagact 1500 ggcacgttgg ccattccatt tatgttggaa tcgttgagca cggagctgta agaaaatcgg 1560 atacagetgt tgateataca ggeetaggtg egataaaatg gagegegtte gaegatgetg 1620 aaccagaccg getgaacate ttgtttcage ataaccatte ggatttteeg atttattatg 1680 agacggattg gtaccgtctg ctgacgggca agctgattgt aaatgcgtgt attaatcctt 1740 taactgcgtt attgcaagtg aaaaatggag aactgctgac aacgccagct tatctggctt 1800 ttatgaaget ggtattteag gaggeatgee geattttaaa aettgaaaat gaagaaaagg 1860 cttgggagcg ggttcaggcc ytttgtgggc aaacgaaaga gaatcgttca tcaatgctgg 1920 ttgacgtcat tggaggccgg cagacggaag ctgacgccat tatcggatac ttattgaagg 1980 aagcaagtct tcaaggtctt gatgeegtee aectagagtt tttatatgge agcateaaag 2040 cattggagcg aaatacaaac aaagtetttt gagettttte ggtaacatge tatactcatt 2100 teggateact aactatttat tggagaaagg aagttetaga agatgeaget aactgaactt 2160 tocatcaaaa atcagaatgt gtttgtacag cactatatag atggcaaaga agaaatgtct 2220 tetttttttg attacagtat teateataag gaeatgtgge gegaaagaet ggaagaetta 2280 tetteceggt ttttegeaag agaggaattg geggegtaet taacetetta ecataataaa 2340 ttcggttcaa gtgcgatgca gtctgctatt gagaagctga aggacccgtc aagtgccgct 2400 gtagtcggcg gacagcaggc aggactttta acaggaccgc tttacaccat acataaaatc 2460 atttcaatca ttgttttagc aaagcaacaa gaaaaggaac tgcaagtgcc tgtcatacca 2520 atcttctggg tggctggaga agaccacgat ttggatgaga ttaattttgt tcacacatct 2580 gaagagaatg ggcctgtgaa aaaaaagctg cctcagtctt attggaagaa atcatcagca 2640 gcgagtacat cgcttgatca ggaaaagtgt gccgcgtgga tagatgatgt ttttgccgct 2700

	tttgaagaaa	cagaccatac	gaatacactt	ctcgacaatg	tgaaacgatg	tttaagggaa	2760
	tctgttacgt	ttactgactt	ctttgaactg	ctgatcgcgg	atttgttcca	agaagaggc	2820
	ttagttttat	taaattctgg	ggatcctgtt	ttggcggatg	agagaagatt	ttcagcctga	2880
1	tacagattaa	atcagaacgc	agaagcggtc	tgataaaaca	gaatttgcct	ggcggcagta	2940
	gcgcggtggt	cccacctgac	cccatgccga	actcagaagt	gaaacgccgt	agcgccgatg	3000
	gtagtgtggg	gtctccccat	gcgagagtag	ggaactgcca	ggcatcaaat	aaaacgaaag	3060
	gctcagtcga	aagactgggc	ctttcgtttt	atctgttgtt	tgtcggtgaa	cgctctcctg	3120
	agtaggacaa	atccgccggg	agcggatttg	aacgttgcga	agcaacggcc	cggagggtgg	3180
	cgggcaggac	gcccgccata	aactgccagg	catcaaatta	agcagaaggc	catcctgacg	3240
	gatggccttt	ttgcgtttct	acaaactctt	ggtacccaga	aaaagcggca	aaagcggctg	3300
	ttaaaaaagc	gaaatcgaag	aagctgtctg	ccgctaagac	ggaatatcaa	aagcgttctg	3360
	ctgttgtgtc	atctttaaaa	gtcacageeg	atgaatccca	gcaagatgtc	ctaaaatact	3420
	tgaacaccca	gaaagataaa	ggaaatgcag	accaaattca	ttcttattat	gtggtgaacg	3480
	ggattgctgt	tcatgcctca	aaagaggtta	tggaaaaagt	ggtgcagttt	cccgaagtgg	3540
	aaaaggtgct	tcctaatgag	aaacggcagc	tttttaagtc	atcctcccca	tttaatatga	3600
	aaaaagcaca	gaaagctatt	aaagcaactg	acggtgtgga	atggaatgta	gaccaaatcg	3660
	atgccccaaa	agcttgggca	cttggatatg	atggaactgg	cacggttgtt	gcgtccattg	3720
	ataccggggt	ggaatggaat	catccggcat	taaaagagaa	atatcgcgga	tataatccgg	3780
	aaaatcctaa	tgagcctgaa	aatgaaatga	actggtatga	tgccgtagca	ggcgaggcaa	3840
	gcccttatga	tgatttggct	catggaaccc	acgtgacagg	cacgatggtg	ggctctgaac	3900
	ctgatggaac	aaatcaaatc	ggtgtagcac	ctggcgcaaa	atggattgct	gttaaagcgt	3960
	tctctgaaga	tggcggcact	gatgctgaca	ttttggaagc	tggtgaatgg	gttttagcac	4020
	caaaggacgc	ggaaggaaat	ccccacccgg	aaatggctcc	tgatgttgtc	aataactcat	4080
	ggggaggggg	ctctggactt	gatgaatggt	acagagacat	ggtcaatgcc	tggcgttcgg	4140
	ccgatatttt	ccctgagttt	tcagcgggga	atacggatct	ctttattccc	ggcgggcctg	4200
	gttctatcgc	aaatccggca	aactatccag	aatcgtttgc	aactggagcg	actgagaatt	4260
	ccaattcccc	atggagagaa	aagaaaatcg	ctaatgttga	ttactttgaa	cttctgcata	4320
	ttcttgaatt	taaaaaggct	gaaagagtaa	aagattgtgc	tgaaatatta	gagtataaac	4380
	aaaatcotoa	aacaggggaa	agaaagttgt	at coact at a	attt======	.	

gtccaatgtg caactggagg agagcaatga aacatggcat tcagtcacaa aaggttgttg 4500 ctgaagttat taaacaaaag ccaacagttc gttggttgtt tctcacatta acagttaaaa 4560 atgtttatga tggcgaagaa ttaaataaga gtttgtcaga tatggctcaa ggatttcgcc 4620 gaatgatgca atataaaaaa attaataaaa atcttgttgg ttttatgcgt gcaacggaag 4680 tgacaataaa taataaagat aattottata atcagcacat gcatgtattg gtatgtgtgg 4740 aaccaactta ttttaagaat acagaaaact acgtgaatca aaaacaatgg attcaatttt 4800 ggaaaaaggc aatgaaatta gactatgatc caaatgtaaa agttcaaatg attcgaccga 4860 aaaataaata taaateggat atacaategg caattgaega aaetgeaaaa tateetgtaa 4920 aggatacgga ttttatgacc gatgatgaag aaaagaattt gaaacgtttg tctgatttgg 4980 aggaaggttt acaccgtaaa aggttaatct eetatggtgg tttgttaaaa gaaatacata 5040 aaaaattaaa ccttgatgac acagaagaag gcgatttgat tcatacagat gatgacgaaa 5100 aagccgatga agatggattt tctattattg caatgtggaa ttgggaacgg aaaaattatt 5160 ttattaaaga gtagttcaac aaacgggcca tattgttgta taagtgatga aatactgaat 5220 ttaaaactta gtitatatgt ggtaaaatgt tttaatcaag tttaggagga attaattatg 5280 aagtgtaatg aatgtaacag ggttcaatta aaagagggaa gegtatcatt aacectataa 5340 actacgtctg ccctcattat tggagggtga aatgtgaata catcctattc acaatcgaat 5400 ttacgacaca accaaatttt aatttggctt tgcattttat cttttttag cytattaaat 5460 gaaatggttt tgaacgtctc attacctgat attgcaaatg attttaataa accacctgcg 5520 agtacaaact gggtgaacac agcetttatg ttaacetttt ecattggaac agetgtatat 5580 ggaaagctat ctgatcaatt aggcatcaaa aggttactcc tatttggaat tataataaat 5640 tgtttcgggt cggtaattgg gtttgttggc cattctttct tttccttact tattatggct 5700 cgttttattc aaggggctgg tgcagctgca tttccagcac tcgtaatggt tgtagttgcg 5760 cgctatattc caaaggaaaa taggggtaaa gcatttggtc ttattggatc gatagtagcc 5820 atgggagaag gagtcggtcc agcgattggt ggaatgatag cccattatat tcattggtcc 5880 tatettetae teatteetat gataacaatt ateaetgtte egtttettat gaaattatta 5940 aagaaagaag taaggataaa aggtcatttt gatatcaaag gaattatact aatgtctgta 6000 gtgctgtcat tcctgatatt tgtaaaacat atcaggaaag taacagatcc ttttgttgat 6120 cccggattag ggaaaaatat accttttatg attggagttc tttgtggggg aattatattt 6180

ggaacagtag cagggtttgt ctctatggtt ccttatatga tgaaagatgt tcaccagcta 6240 agtactgccg aaatcggaag tgtaattatt ttccctggaa caatgagtgt cattatttc 6300 ggctacattg gtgggatact tgttgataga agaggtcctt tatacgtgtt aaacatcgga 6360 gttacatttc tttctgttag ctttttaact gcttcctttc ttttagaaac aacatcatgg 6420 ttcatgacaa ttataatcgt atttgtttta ggtgggcttt cgttcaccaa aacagttata 6480 tcaacaattg tttcaagtag cttgaaacag caggaagctg gtgctggaat gagtttgctt 6540 aactttacca gctttttatc agagggaaca ggtattgcaa ttgtaggtgg tttattatcc 6600 ataccettae ttgateaaag gttgttaeet atggaagttg ateagteaac ttatetgtat 6660 agtaatttgt tattactttt ttcaggaatc attgtcatta gttggctggt taccttgaat 6720 gtatataaac attctcaaag ggatttctaa atcgttaagg gatcaacttt gggagagagt 6780 tcaaaattga tcctttttt ataacagttc gaagcggccg caattcttga agacgaaagg 6840 gcctcgtgat acgcctattt ttataggtta atgtcatgat aataatggtt tcttagacgt 6900 caggtggcac tittcgggga aatgtgcgcg gaacccctat tigttlatit tictaaatac 6960 attcaaatat gtatccgctc atgagacaat aaccctgata aatgcttcaa taatattgaa 7020 aaaggaagag tatgagtatt caacatttee gtgtegeeet tatteeettt tttgeggeat 7080 tttgccttcc tgtttttgct cacccagaaa cgctggtgaa agtaaaagat gctgaagatc 7140 agttgggtgc acgagtgggt tacatcgaac tggatctcaa cagcggtaag atccttgaga 7200 gttttcgccc cgaagaacgt tttccaatga tgagcacttt taaagttctg ctatgtggcg 7260 cggtattatc ccgtattgac gccgggcaag agcaactcgg tcgccgcata cactattctc 7320 agaatgactt ggttgagtac tcaccagtca cagaaaagca tcttacggat ggcatgacag 7380 taagagaatt atgcagtgct gccataacca tgagtgataa cactgcggcc aacttacttc 7440 tgacaacgat cggaggaccg aaggagctaa ccgctttttt gcacaacatg ggggatcatg 7500 taactcgcct tgatcgttgg gaaccggagc tgaatgaagc cataccaaac gacgagcgtg 7560 acaccacgat gcctgcagca atggcaacaa cgttgcgcaa actattaact ggcgaactac 7620 ttactctagc ttcccggcaa caattaatag actggatgga ggcggataaa gttgcaggac 7680 cacttetgeg eteggeeett eeggetgget ggtttattge tgataaatet ggageeggtg 7740 agegtgggte tegeggtate attgeageae tggggeeaga tggtaageee teeegtateg 7800 tagttatcta cacgacgggg agtcaggcaa ctatggatga acgaaataga cagatcgctg 7860 agataggtgc ctcactgatt aagcattggt aactgtcaga ccaagtttac tcatatatac 7920

tttagattga tttaaaactt catttttaat ttaaaaggat ctaggtgaag atcctttttg 7980 ataatctcat gaccaaaatc ccttaacgtg agttttcgtt ccactgagcg tcagaccccg 8040 tagaaaagat caaaggatct tettgagate ettttttet gegegtaate tgetgettge 8100 aaacaaaaaa accaccgcta ccagcggtgg tttgtttgcc ggatcaagag ctaccaactc 8160 tttttccgaa ggtaactggc ttcagcagag cgcagatacc aaatactgtc cttctagtgt 8220 agcegtagtt aggecaecae tteaagaaet etgtageaec geetaeatae etegetetge 8280 taatcctgtt accagtggct gctgccagtg gcgataagtc gtgtcttacc gggttggact 8340 caagacgata gttaccggat aaggcgcagc ggtcgggctg aacggggggt tcgtgcacac 8400 agcccagctt ggagcgaacg acctacaccg aactgagata cctacagcgt gagctatgag 8460 aaagcgccac gcttcccgaa gggagaaagg cggacaggta tccggtaagc ggcagggtcg 8520 gaacaggaga gegeaegagg gagetteeag ggggaaaege etggtatett tatagteetg 8580 tegggttteg ceacetetga ettgagegte gatttttgtg atgetegtea ggggggegga 8640 gcctatggaa aaacgccagc aacgcggcct ttttacggtt cctggccttt tgctggcctt 8700 ttgeteacat gttettteet gegttateee etgattetgt ggataacegt attacegeet 8760 ttgagtgage tgataceget egeegeagee gaaegaeega gegeagegag teagtgageg 8820 aggaagegga agagegeetg atgeggtatt tteteettae geatetgtge ggtattteae 8880 accgcatatg gtgcactctc agtacaatct gctctgatgc cgcatagtta agccagtata 8940 cacteegeta tegetaegtg actgggteat ggetgegeee egacaeeege caacaeeege 9000 tgacgcgccc tgacgggctt gtctgctccc ggcatccgct tacagacaag ctgtgaccgt 9060 ctccgggagc tgcatgtgtc agaggttttc accgtcatca ccgaaacgcg cgaggcagct 9120 geggtaaage teateagegt ggtegtgaag egatteacag atgtetgeet gtteateege 9180 gtccagctcg ttgagtttct ccagaagcgt taatgtctgg cttctgataa agcgggccat 9240 gttaagggcg gtttttcct gtttggtcac ttgatgcctc cgtgtaaggg ggaatttctg 9300 ttcatggggg taatgatacc gatgaaacga gagaggatgc tcacgatacg ggttactgat 9360 gatgaacatg cccggttact ggaacgttgt gagggtaaac aactggcggt atggatgcgg 9420 cgggaccaga gaaaaatcac tcagggtcaa tgccagcgct tcgttaatac agatgtaggt 9480 gttccacagg gtagccagca gcatcctgcg atgcagatcc ggaacataat ggtgcagggc 9540 gctgacttcc gcgtttccag actttacgaa acacggaaac cgaagaccat tcatgttgtt 9600 getcaggtcg cagacgtttt gcagcagcag tegettcacg ttegetcgcg tateggtgat 9660

tcattctgct aaccagtaag gcaaccccgc cagcctagcc gggtcctcaa cgacaggagc 9720
acgatcatgc gcacccgtgg ccaggaccca acgctgcccg agatgcgccg cgtgcggctg 9780
ctggagatgg cggacgcgat ggatatgttc tgccaagggt tggtttgcgc attcacagtt 9840
ctccgcaaga attgattggc tccaattctt ggagtggtga atccgttagc gaggtgccgc 9900
cggcttccat tcaggtcgag gtggcccggc tccatgcacc gcgacgcaac gcggggaggc 9960
agacaaggta tagggcggc cctacaatcc atgccaaccc gttccatgtg ctcgccgagg 10020
cggcataaat cgccgtgacg atcagcggtc cagtgatcga agttaggctg gtaagagccg 10080
cgagcgatcc ttgaagctgt ccctgatggt cgtcatctac ctgcctggac agcatggcct 10140
gcaacgcggg catcccgatg ccgccggaag cgagaagaat cataatggg aaggccatcc 10200
agcctcgcgt cg

<210> 82

<211> 10426

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN267 plasmid

<400> 82
aacaaaattc tccagtcttc acatcggttt gaaaggagga agcggaagaa tgaagtaaga 60
gggatttttg actccgaagt aagtcttcaa aaaatcaaat aaggagtgtc aagaatgttt 120
gcaaaacgat tcaaaacctc tttactgccg ttattcgctg gattttatt gctgttcat 180
ttggttctgg caggaccggc ggctgcgagt gctgaaacgg cgaacaaatc gaatgagctt 240
acagcaccgt cgatcaaaag cggaaccatt cttcatgcat ggaattggtc gttcaatacg 300
ttaaaacaca atatgaagga tattcatgat gcaggatata cagccattca gacatctccg 360
attaaccaag taaaggaagg gaatcaagga gataaaagca tgtcgaactg gtactggctg 420
tatcagccga catcgtatca aattggcaac cgttacttag gtactgaaca agaatttaaa 480
gaaatgtgtg cagccgctga agaatatggc ataaaggtca ttgttgacgc ggtcatcaat 540
cataccacca gtgattatgc cgcgattcc aatgaggta agagtattcc aaactggaca 600
catggaaaca cacaaattaa aaactggtct gatcgaaata gtacataatg gatttcctta 660
cgcgaaatac gggcagacat ggcctgcccg gttattatta tttttgacac cagaccaact 720
ggtaatggta gcgaccgcg ctcaggatcg tctcggtacc aagagtttgt agaaacgcaa 780
aaaggccatc cgtcaggatg gccttctgct taatttgatg cctggcagtt tatggcgggc 840

gtectgeecg ecaeceteeg ggeegttget tegeaacgtt caaateeget eeeggeggat 900 ttgtcctact caggagagcg ttcaccgaca aacaacagat aaaacgaaag gcccagtctt 960 togactgage etttegtttt atttgatgee tggeagttee etactetege atggggagae 1020 cccacactac catcggcgct acggcgtttc acttctgagt tcggcatggg gtcaggtggg 1080 accaccgcgc tactgccgcc aggcaaattc tgttttatca gaccgcttct gcgttctgat 1140 ttaatctgta tcaggctgaa aatcttctct catccgccaa aacaggatcc atcacgaagc 1200 gtcgtatcga aaaaattaat tttgcgcaac ggagaccacc gcttccttct tcttgccttg 1260 tttcacaaac ggcatcattt cacgaagett tetteccaet acttegattt gatgttegtt 1320 ctcgcttgca ttgatagcgt tgaaacgagg acggtttact tggttttcga cgatccactc 1380 ttttgcgaat gtaccgtttt ggatatettt taataettet ttcatagatt ettttaettt 1440 agcgtccaca acgcgagggc ctgatacgaa atctccccac tgtgctgtgt cagagattga 1500 atateteatt cetgeaagte ettettegta cataaggtet acgattaatt teageteatg 1560 aagacacteg aagtatgeaa gtteaggetg ataacetget teagttaagg ttteaaatee 1620 ggotttgaca agogogotta atoogoogoa aagaactgot tyotoacoga acaaatotgt 1680 ttotgtttot totttaaatg togtttotaa tacgeooget ottgegooge egatteettt 1740 agcataagcg agggettigt etettgette tecagteaca tettgataga tigegaacaa 1800 tgcaggtacg ccagctcctt gctcatatgt tcttcttacc aagtgtcecg ggcctttagg 1860 ggccactaag aatacatcta catccgccgg aggaacaatt tgatggaaat gcacgttaaa 1920 tccatgagcg aatactaatg attttcctgc tgtcaattca tctttgattt cagcttcgta 1980 tactttttgc tgctgctcat ccggaagcag aaccatgatg atttcggctt gggctgccgc 2040 ttottttact gaaaatactt tatgtoogto ttottgggot tgagtgaaag attttottg 2100 totaacaccg acgateacgt etacteeget ttetttaagg tteagggeat gtgcgtggcc 2160 ttgcgaaccg tacccgataa ccgctactgt ttttccagcc aatacgttct ctttgatatc 2220 accyttataa tatactttta ccatttcaat ctctcccttg ttatytttta tacaatagat 2280 attgttttat tggatgacgc cttttgctgg ttcccctcgc aaaagccgtt gtacctgttc 2340 togogattto tttaatgoca taaggittta ataactcaat aagogottoa attiigitag 2400 attcacctgt cacctgaaca acgatgctgt ctctgctgac atcaacgaca gaggctctaa 2460 acggttctat gattccatta atctctgttc ttgttgaagg tgcggagaca accttgatta 2520 . aggccagete ectetggaca ategattgat ttgtgatgte tgtgaettte ageacateaa 2580 tctgtttgtt gagctgtttc gttaactgtt caacatcatt ttcaccttca acatgaacga 2640 cgaaggtgat tctggaaacg ccggctgttt ctgtgtgtcc aactgtaatg ctttcaatgt 2700 tgtaatgcct ttttgtgaat agaccggtga tccggtttaa caccccggag cggttcacca 2760 cagtcaatgt gataattctt ttcaaggttt cacccccacc atttcatgca gccctttccc 2820 ' cggagccacc atcgggaata ctttttcttc gctggcaacc cgcacgtcaa tgacaacagg 2880 ttctcttgat gttaatgcct cttccagctt ttcctttgct tccgcttctg atgaaattct 2940 gatgccttta atgccgtatg cttcggacaa tttgacgaag tcaggctgag aagcgaattt 3000 agattctgaa taacgttctt catagaaaat ttcctgccac tgtctgacca ttccgagaca 3060 agcgttattt aaaatcacta ccttgaccgg aagatttaat tcgcgaataa catcgagttc 3120 ttgaagcgtc atttggaatc cgccgtctcc gacaaccgcg acaacagtag catcttttc 3180 ggccagctqt gcgccgatcg ccgccggaag accgaatccc atcgttccaa gtccqcctqa 3240 cgtgacccat ttatctgctt tttggaacgg ataaaattgc gctgaccaca tttgatgctg 3300 gcctacatcc gttgcgacaa tggcctctcc ttttgtaaat tgatgaatat attcaatcaa 3360 tttctgaggt ttaaaacctt cttcttcatt atctacatac cagagcggat actcttcttt 3420 ccattctgcg agctgttttt tccattcgct tgaatcgctt tgtttgccgt cttgtttgat 3480 cageteetge aggacaattt tgetgtetee gactacagga atetgtgttt teatgatttt 3540 teegatttea getggateaa tategatgtg ggetatettt gegtttetgg caaagtgttt 3600 caggittect gigacacggi catcaaaacg ggcgccgata cigattaata gatcacatic 3660 atgaagggcc atattggctg tataagtacc gtgcattccc gccatcccta ggaaaagcgg 3720 atggtcagcc gggaagcctc cgagccccaa aagggtgtgt gccacaggga tttgctgctg 3780 ttcagcataa ttttttaatt cttctgacgc ttttccgtgc agtacgcccg cacccgccag 3840 gatcaccggt tttttcgcac tgctcacggc ttccacaagc ttgcggatct gcaaataatt 3900 cggctctgtt gtcggctggt atcccgggag attcatctca tgatcgtagc tgaattctcc 3960 ttcaattgtt gctacatctt tcggaatatc aatcaataca ggtccgggtc ttccagttgt 4020 tgcaatatgg aacgettett taatgatgeg eggeagatet teeggetgge gaacetggta 4080 gctgtgtttt gttactggca tcgtaatccc taaaatgtct gcttcctgaa atgcatcqct 4140 cccgattaca gaggttgcta cctgccctgt aaagacgact aacggcaatg aatcaatcat 4200 ggcatcagca aggcctgtaa caaggtttgt cgctcccggc ctgacgtggc aatgacgaca 4260 ccggtttccg gagacccttg cgtatccctc cgctgcatga attgctcctt gttcgtgacg 4320

ggaaggatat gtaccaacct gaatgtatag cttatcgtaa atcggaagca cagccccgcc 4380 cggataaccg aagatcattt ctactttctc tttttttaat gattcaatca gcattaatcg 4440 tecgeteate gtetgtgtae atteggeaga tgetgaatee acetgtaeat tagteeceat 4500 tttatctcct cctctaqaqc qtcctqctqt tqttaaqatt attataccac accttgtaga 4560 taaagtcaac aactttttgc aaaatttttc aggaatttta gcagaggttg ttctggatgt 4620 agaacaaaac atctttccgc tcttgtgctg ttaggatatc tttcttggaa gctaggtagg 4680 cctcgagtta tggcagttgg ttaaaaggaa acaaaaagac cgttttcaca caaaacggtc 4740 tttttcgatt tctttttaca gtcacagcca cttttgcaaa aaccggacag cttcatgcct 4800 tataactgct gtttcggtcg acgaagcggc cgccgtttaa acgaattcct gcagctggcg 4860 aatqqcqatt ttcqttcqtq aatacatqtt ataataacta taactaataa cgtaacgtga 4920 ctggcaagag atattttaa aacaatgaat aggtttacac ttactttagt tttatggaaa 4980 tgaaagatca tatcatatat aatctagaat aaaattaact aaaataatta ttatctagat 5040 aaaaaattta gaagccaatg aaatctataa ataaactaaa ttaagtttat ttaattaaca 5100 actatggata taaaataggt actaatcaaa atagtgagga ggatatattt gaatacatac 5160 qaacaaatta ataaaqtqaa aaaaatactt cggaaacatt taaaaaataa ccttattggt 5220 acttacatgt ttggatcagg agttgagagt ggactaaaac caaatagtga tcttgacttt 5280 ttagtcgtcg tatctgaacc attgacagat caaagtaaag aaatacttat acaaaaaatt 5340 agacctattt caaaaaaaat aggagataaa agcaacttac gatatattga attaacaatt 5400 attattcagc aagaaatggt accgtggaat catcctccca aacaagaatt tatttatgga 5460 gaatggttac aagagcttta tgaacaagga tacattcctc agaaggaatt aaattcagat 5520 ttaaccataa tgctttacca aqcaaaacqa aaaaataaaa gaatatacgg aaattatgac 5580 ttagaggaat tactacctga tattccattt tctgatgtga gaagagccat tatggattcg 5640 tcagaggaat taatagataa ttatcaggat gatgaaacca actctatatt aactttatgc 5700 cgtatgattt taactatgga cacgggtaaa atcataccaa aagatattgc gggaaatgca 5760 gtggctgaat cttctccatt agaacatagg gagagaattt tgttagcagt tcgtagttat 5820 cttqqaqaqa atattqaatq qactaatqaa aatqtaaatt taactataaa ctatttaaat 5880 aacaqattaa aaaaattata aaaaaattga aaaaatggtg gaaacacttt tttcaatttt 5940 tttgttttat tatttaatat ttgggaaata ttcattctaa ttggtaatca gattttagaa 6000 aacaataaac cettgcatag ggggategat atcegtttag getgggeggt gatagettet 6060 cgttcaggca gtacgcctct tttcttttcc agacctgagg gaggcggaaa tggtgtgagg 6120 ttcccgggga aaagccaaat aggcgatcgc gggagtgctt tatttgaaga tcaggctatc 6180 actgcggtca atagatttca caatgtgatg gctggacagc ctgaggaact ctcgaacccg 6240 aatggaaaca accagatatt tatgaatcag cgcggctcac atggcgttgt gctggcaaat 6300 gcaggttcat cctctgtctc tatcaatacg gcaacaaaat tgcctgatgg caggtatgac 6360 aataaagctg gagcgggttc atttcaagtg aacgatggta aactgacagg cacgatcaat 6420 gccaggtctg tagctgtgct ttatcctgat gatattgcaa aagcgcctca tgttttcctt 6480 gagaattaca aaacaggtgt aacacattct ttcaatgatc aactgacgat taccttgcgt 6540 gcagatgcga atacaacaaa agccgtttat caaatcaata atggaccaga cgacaggcgt 6600 ttaaggatgg agatcaattc acaatcggaa aaggagatcc aatttggcaa aacatacacc 6660 atcatgttaa aaggaacgaa cagtgatggt gtaacgagga ccgagaaata cagttttgtt 6720 aaaagagatc cagcgtcggc caaaaccatc ggctatcaaa atccgaatca ttggagccag 6780 gtaaatgctt atatctataa acatgatggg agccgagtaa ttgaattgac cggatcttgg 6840 cctggaaaac caatgactaa aaatgcagac ggaatttaca cgctgacgct gcctgcggac 6900. acggatacaa ccaacgcaaa agtgattttt aataatggca gcgcccaagt gcccggtcag 6960 aatcageetg getttgatta egtgetaaat ggtttatata atgaeteggg ettaageggt 7020 tetetteece attgagggea aggetagaeg ggaettaceg aaagaaacca teaatgatgg 7080 tttctttttt gttcataaat cagacaaaac ttttctcttg caaaagtttg tgaagtgttg 7140 cacaatataa atgtgaaata cttcacaaac aaaaagacat caaagagaaa cataccctgc 7200 aaggatgctg atattgtctg catttgcgcc ggagcaaacc aaaaacctgg tgagacacgc 7260 cttgaattag tagaaaagaa cttgaagatt ttcaaaggca tcgttagtga agtcatggcg 7320 agcggatttg acggcatttt cttagtcggg cggcacctcg ctaacggatt caccactcca 7380 agaattggag ccaatcaatt cttgcggaga actgtgaatg cgcaaaccaa cccttggcag 7440 aacatateca tegegteege cateteeage ageegeacge ggegeatete gggeagegtt 7500 gggtcctggc cacgggtgcg catgatcgtg ctcctgtcgt tgaggacccg gctaggctgg 7560 cggggttgcc ttactggtta gcagaatgaa tcaccgatac gcgagcgaac gtgaagcgac 7620 tgctgctgca aaacgtctgc gacctgagca acaacatgaa tggtcttcgg tttccgtgtt 7680 togtaaagto tggaaacgog gaagtoagog cootgoacca ttatgttoog gatotgoato 7740 gcaggatgct gctggctacc ctgtggaaca cctacatctg tattaacgaa gcgctggcat 7800

tgaccotgag tgatttttct otggtocogo ogcatocata cogocagttg tttaccotca 7860 caacgtteca gtaaceggge atgtteatea teagtaacee gtategtgag catectetet 7920 cgtttcatcg gtatcattac ccccatgaac agaaatcccc cttacacgga ggcatcagtg 7980 accaaacagg aaaaaaccgc ccttaacatg gcccgcttta tcagaagcca gacattaacg 8040' cttctggaga aactcaacga gctggacgcg gatgaacagg cagacatctg tgaatcgctt 8100 cacgaccacg ctgatgaget ttaccgcage tgeetegege gttteggtga tgaeggtgaa 8160 aacctctgac acatgcagct cccggagacg gtcacagctt gtctgtaagc ggatgccggg 8220 agcagacaag cccgtcaggg cgcgtcagcg ggtgttggcg ggtgtcgggg cgcagccatg 8280 acceagteae gtagegatag eggagtgtat actggettaa etatgeggea teagageaga 8340 ttgtactgag agtgcaccat atgcggtgtg aaataccgca cagatgcgta aggagaaaat 8400 acceptation generated gettectors teactgasts getgegeter geogttegge 8460 tgcggcgagc ggtatcagct cactcaaagg cggtaatacg gttatccaca gaatcagggg 8520 ataacgcagg aaagaacatg tgagcaaaag gccagcaaaa ggccaggaac cgtaaaaagg 8580 cegegttget ggegttttte cataggetee geeeceetga egageateae aaaaategae 8640 geteaagtea gaggtggega aaccegacag gactataaag ataccaggeg tetececetg 8700 gaageteest egtgegetet estgtteega eestgeeget taceggatae etgteegeet 8760 ttetecette gggaagegtg gegetttete atageteaeg etgtaggtat eteagttegg 8820 tgtaggtcgt tcgctccaag ctgggctgtg tgcacgaacc ccccgttcag cccgaccgct 8880 gegeettate eggtaactat egtettgagt ecaaceeggt aagacaegae ttategeeae 8940 tggcagcagc cactggtaac aggattagca gagcgaggta tgtaggcggt gctacagagt 9000 tcttgaagtg gtggcctaac tacggctaca ctagaaggac agtatttggt atctgcgctc 9060 tgctgaagcc agttaccttc ggaaaaagag ttggtagctc ttgatccggc aaacaaacca 9120 ccgctggtag cggtggtttt tttgtttgca agcagcagat tacgcgcaga aaaaaaggat 9180 ctcaagaaga teetttgate ttttctaegg ggtetgaege teagtggaae gaaaacteae 9240 gttaagggat tttggtcatg agattatcaa aaaggatctt cacctagatc cttttaaatt 9300 aaaaatgaag ttttaaatca atctaaagta tatatgagta aacttggtct gacagttacc 9360 aatgcttaat cagtgaggca cctatctcag cgatctgtct atttcgttca tccatagttg 9420 cctgactccc cgtcgtgtag ataactacga tacgggaggg cttaccatct ggccccagtg 9480 ctgcaatgat accgcgagac ccacgctcac cggctccaga tttatcagca ataaaccagc 9540

cageeggaag ggeegagege agaagtggte etgeaacttt ateegeetee ateeagteta 9600 ttaattgttg ccgggaagct agagtaagta gttcgccagt taatagtttg cgcaacgttg 9660 ttgccattgc tgcaggcatc gtggtgtcac gctcgtcgtt tggtatggct tcattcagct 9720 ccggttccca acgatcaagg cgagttacat gatcccccat gttgtgcaaa aaagcggtta 9780 gctccttcgg tcctccgatc gttgtcagaa gtaagttggc cgcagtgtta tcactcatgg 9840 ttatggcage actgcataat tetettactg teatgccate egtaagatge ttttetgtga 9900 ctggtgagta ctcaaccaag tcattctgag aatagtgtat gcggcgaccg agttgctctt 9960 gcccggcgtc aacacgggat aataccgcgc cacatagcag aactttaaaa gtgctcatca 10020 ttggaaaacg ttcttcgggg cgaaaactct caaggatctt accgctgttg agatccagtt 10080 cgatgtaacc cactcgtgca cccaactgat cttcagcatc ttttactttc accagcgttt 10140 ctgggtgagc aaaaacagga aggcaaaatg ccgcaaaaaa gggaataagg gcgacacgga 10200 aatgttgaat actcatactc ttccttttc aatattattg aagcatttat cagggttatt 10260 gtctcatgag cggatacata tttgaatgta tttagaaaaa taaacaaata ggggttccgc 10320 gcacatttcc ccgaaaagtg ccacctgacg tctaagaaac cattattatc atgacattaa 10380 cotataaaaa taggogtato acgaggooot ttogtottoa agaatt 10426

<210> 83

<211> 4191

<212> DNA

<213> Artificial Sequence

<220>

MICCOCOID. JUIC

012177282 L s

<223> Description of Artificial Sequence: Recombinant pAN263 plasmid

<400> 83

ccaatacgca aaccgcctct ccccgcgcgt tggccgattc attaatgcag cccggatccg 60
aaagaagagg atatcccggt tttacagaag gcattggatg atccaaaggt gtccatcaga 120
agacaggctg ttgtgtactt aggaatgatt gaaacacctg atgttcttcc tctattgtat 180
aaagcacttg aggacaaagc tgtatcagtc agaagaacgg ccggagactg cctgtctgat 240
atcggcgatc ctcaagccat tcctgctatg atcaagtcat taagcgactc cagcaagctt 300
gttcgctggc gtgccgccat gttcctgtac gaagtcggcg atgaaagtgc aattgaagct 360
ttgcgcgctg ccgaagatga ccccgaattt gaggtcagcc ttcaagtcaa aatggcgctt 420
gaacgtattg agcatggaga agaagcaaaa ggttctgttt ggaaacaaat gacggaaagc 480

igaaaaaaag	gcgaataaag	ataaaaaagg	tgcagatcat	gcaccttttt	tatgtgaatt	540
gtcgaccga	aacagcagtt	ataaggcatg	aagctgtccg	gtttttgcaa	aagtggctgt	600
gactgtaaaa	agaaatcgaa	aaagaccgtt	ttgtgtgaaa	acggtctttt	tgtttccttt	660
aaccaactg	ccataactcg	aggcctacct	agcttccaag	aaagatatcc	taacagcaca	720
agagcggaaa	gatgttttgt	tctacatcca	gaacaacctc	tgctaaaatt	cctgaaaaat	780
ttgcaaaaa	gttgttgact	ttatctacaa	ggtgtggtat	aataatctta	acaacagcag	840
gacgctctag	aggaggagac	accatggcag	aattacgcag	taatatgatc	acacaaggaa	900
cgatagagc	tecgcacege	agtttgcttc	gtgcagcagg	ggtaaaagaa	gaggatttcg	960
gcaagccgtt	tattgcggtg	tgtaattcat	acattgatat	cgttcccggt	catgttcact	1020
tgcaggagtt	tgggaaaatc	gtaaaagaag	caatcagaga	agcaggggc	gttccgtttg	1080
aatttaatac	cattggggta	gatgatggca	tcgcaatggg	gcatateggt	atgagatatt	1140
cgctgccaag	ccgtgaaatt	atcgcagact	ctgtggaaac	ggttgtatcc	gcacactggt	1200
ttgacggaat	ggtctgtatt	ccgaactgcg	acaaaatcac	accgggaatg	cttatggcgg	1260
caatgcgcat	-caacattccg	acgatttttg	tcagcggcgg	accgatggcg	-gcaggaagaa	1320
caagttacgg	gcgaaaaatc	tecettteet	cagtattcga	aggggtaggc	gcctaccaag	1380
cagggaaaat	caacgaaaac	gagcttcaag	aactagagca	gttcggatgc	ccaacgtgcg	1440
ggtcttgctc	aggcatgttt	acggcgaact	caatgaactg	tetgteagaa	gcacttggtc	1500
ttgctttgcc	gggtaatgga	accattctgg	caacatetee	ggaacgcaaa	gagtttgtga	1560
gaaaatcggc	tgcgcaatta	atggaaacga	ttcgcaaaga	tatcaaaccg	-cgtgatattg	1620
ttacagtaaa	agcgattgat	aacgcgtttg	cactcgatat	ggcgctcgga	ggttctacaa	1:680
ataccgttct	tcataccctt	gcccttgcaa	acgaagccgg	cgttgaatac	tctttagaac	1740
gcattaacga	agtcgctgag	cgcgtgccgc	acttggctaa	gctggcgcct	-gcatcggatg	1800
tgtttattga	agatcttcac	gaagcgggcg	gcgtttcagc	ggctctgaat	gagetttega	1860
agaaagaagg	agcgcttcat	ttagatgcgc	tgactgttac	aggaaaaact	cttggagaaa	1920
ccattgccgg	acatgaagta	aaggattatg	acgtcattca	cccgctggat	caaccattca	1980
ctgaaaaggg	aggccttgct	gttttattcg	gtaatctago	teeggaegge	gctatcatta	2040
aaacaggcgg	cgtacagaat	gggattacaa	gacacgaagg	gccggctgtc	gtattogatt	2100
ctcaggacga	ggcgcttgac	ggcattatca	accgaaaagt	aaaagaaggc	gacgttgtca	2160
tcatcagata	cgaagggcca	aaaqqcqqac	ctggcatgcc	ggaaatgcto	gcqccaacat	2220

cccaaatcgt tggaatggga ctcgggccaa aagtggcatt gattacggac ggacgttttt 2280 ccggagcctc ccgtggcctc tcaatcggcc acgtatcacc tgaggccgct gagggcgggc 2340 cgcttgcctt tgttgaaaac ggagaccata ttatcgttga tattgaaaaa cgcatcttgg 2400 atgtacaagt gccagaagaa gagtgggaaa aacgaaaagc gaactggaaa ggttttgaac 2460 cgaaagtgaa aaccggctac ctggcacgtt attctaaact tgtgacaagt gccaacaccg 2520 gcggtattat gaaaatctag acccctggcg taatagcgaa gaggcccgca ccgatcgccc 2580 ttcccaacag ttgcgcagcc tgaatggcga atgagcttgc gccgtcccgt caagtcagcg 2640 taatgetetg ceagtgttae aaccaattaa eeaattetga ttagaaaaae teategagea 2700 tcaaatgaaa ctgcaattta ttcatatcag gattatcaat accatatttt tgaaaaagcc 2760 gtttctgtaa tgaaggagaa aactcaccga ggcagttcca taggatggca agatcctggt 2820 ateggtetge gatteegaet egteeaacat caatacaace tattaattte eeetegteaa 2880 aaataaggtt atcaagtgag aaatcaccat gagtgacgac tgaatccggt gagaatggca 2940 aaaggttatg catttettte cagaettgtt caacaggeea gecattaege tegteatcaa 3000 aatcactege atcaaccaaa cegttattea ttegtgattg egeetgageg agaegaaata 3060 cgcgatcgct gttaaaagga caattacaaa caggaatcga atgcaaccgg cgcaggaaca 3120 ctgccagcgc atcaacaata ttttcacctg aatcaggata ttcttctaat acctggaatg 3180 ctgttttccc agggatcgca gtggtgagta accatgcatc atcaggagta cggataaaat 3240 gcttgatggt cggaagaggc ataaattccg tcagccagtt tagtctgacc atctcatctg 3300 taacatcatt ggcaacgcta cetttgccat gtttcagaaa caactctggc gcatcgggct 3360 · teceatacaa teaatagatt gtegeacetg attgeeegae attategega geeeatttat 3420 acceatataa atcagcatee atgttggaat ttaategegg eetegaegag caagaegttt 3480 cccgttgaat atggctcata acaccccttg tattactgtt tatgtaagca gacagtttta 3540 ttgttcatga tgatatattt ttatcttgtg caatgtaaca tcagagattt tgagacactc 3600 gacaagatga tettettgag ategttttgg tetgegegta atetettget etgaaaaega 3660 aaaaaccgcc ttgcagggcg gtttttcgaa ggttctctga gctaccaact ctttgaaccg 3720 aggtaactgg cttggaggag cgcagtcacc aaaacttgtc ctttcagttt agccttaacc 3780 ggcgcatgac ttcaagacta actcctctaa atcaattacc agtggctgct gccagtggtg 3840 cttttgcatg tctttccggg ttggactcaa gacgatagtt accggataag gcgcagcggt 3900 cggactgaac ggggggttcg tgcatacagt ccagcttgga gcgaactgcc tacccggaac 3960

tgagtgtcag gcgtggaatg agacaaacgc ggccataaca gcggaatgac accggtaaac 4020 cgaaaggcag gaacaggaga gcgcacgagg gagccgccag gggaaacgcc tggtatcttt 4080 atagteetgt egggtttege caccactgat ttgagegtea gatttegtga tgettgteag 4140 4191 qqqqqqqqq cctatggaaa aacggctttg ccgcggccct ctcacttccc t <210> 84 <211> 702 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(699) <400> 84 ttg tta ctg gtt atc gat gtg ggg aac acc aat act gta ctt ggt gta 48 Met Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val 10 tat cat gat gga aaa tta gaa tat cac tgg cyt ata gaa aca agc agg 96 Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg 20 25 cat aae aca gaa gat gag ttt ggg atg att ttg cgc tcc tta ttt gat 144 His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp 35 192 cac too ggg off atg ttt gaa day ata gat ggc att att tog toa His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser 55 gta gtg ccg cca atc atg ttt gcg tta gaa aga atg tgc aca aaa tac 240 Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr ttt cat atc gag cct caa att gtt ggt cca ggt atg aaa acc ggt tta 288 Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu 90 aat ata aaa tat gac aat ccg aaa gaa gta ggg gca gac aga atc gta 336 Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val 100 aat gct gtc gct gcg ata cac ttg tac ggc aat cca tta att gtt gtc 384 Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val gat ttc gga acc gcc aca acg tac tgc tat att gat gaa aac aaa caa 432 Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 135 tac atg ggc ggg gcg att gcc cct ggg att aca att tcg aca gag gcg 480 Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala 155 145 150

ctt Leu	tac Tyr	tcg Ser	cgt Arg	gca Ala 165	gca Ala	aag ,Lys	ctt Leu	cct Pro	cgt Arg 170	atc Ile	gaa Glu	atc Ile	acc Thr	cgg Arg 175	ccc Pro	528
gac Asp	aat Asn	att Ile	atc Ile 180	ggʻa Gly	aaa Lys	aac Asn	act Thr	gtt Val 185	agc Ser	gcg Ala	atg Met	caa Gln	tct Ser 190	gga Gly	att Ile	57,6
tta Leu	ttt Phe	ggc Gly 195	tat Tyr	gtc Val	ggc Gly	caa Gln	gtg Val 200	gaa Glu	gga Gly	atc Ile	gtt Val	aag Lys 205	cga Arg	atg Met	aaa Lys	624
tgg Trp	cag Gln 210	gca Ala	aaa Lys	cag Gln	gac Asp	cca Pro 215	agg Arg	tca Ser	ttg Leu	cga Arg	cag Gln 220	gag Gl _. u	gcc Ala	tgg Trp	cgc Arg	672
					aat Asn 230				tag			•				702
<210> 85 <211> 233 <212> PRT <213> Bacillus subtilis																
)> 85 Leu		Val	Ile 5	Asp	Val	Gly	Asr.	Thr 10	Asn	Thr	Val	Leu	Gly 15	Val	
Tyr	His	Asp	Gly 20	Lys	Leu	Glu	Tyr	His 25	Trp	Arg	Ile	Glu	Thr 30	Ser	Arg	
His	Lys	Thr 35	Glu	Asp	Glu	Phe	Gly 40	Met	Ile	Leu	Arg	Ser 45	Leu	Phe	Asp	
His	Ser 50	Gly	Leu	Met	Phe	Glu 55	Gln	Ile	Asp	Gly	Ile 60	Ile	Ile	Ser	Ser	
Val 65	Val	Pro	Pro	Ile	Met 70	Phe	Ala	Leu	Glu	Arg 75	Met	Cys	Thr	Lys	Tyr 80	
Phe	His	Ile	Glu	Pro 85	Gln	Ile	Val	Gly	Pro 90	Gly	Met	Lys	Thr	Gly 95	Leu	
Asn	Ile	Lys	Tyr 100	Asp	Asn	Pro	Lys	Glu 105	Val	Gly	Ala	Asp	Arg 110	Ile	Val	
Asn	Ala	Val 115	Ala	Ala	Ile	His	Leu 120	Tyr	Gly	Asn	Pro	Leu 125	Ile	Val	Val	
Asp	Phe 130	Gly	Thr	Ala	Thr	Thr 135	Tyr	Cys	Tyr	Ile	Asp 140	Glu	Asn	Lys	Gln	
Tyr 145	Met	Gly	Gly	Ala	Ile 150	Ala	Pro	Gly	Ile	Thr 155	Ile	Ser	Thr	Glu	Ala 160	

Leu Tyr Ser Arg Ala Ala Lys Leu Pro Arg Ile Glu Ile Thr Arg Pro 170 Asp Asn Ile Ile Gly Lys Asn Thr Val Ser Ala Met Gln Ser Gly Ile 185 180 Leu Phe Gly Tyr Val Gly Gin Val Glu Gly Ile Val Lys Arg Met Lys 200 Trp Gln Ala Lys Gln Asp Pro Arg Ser Leu Arg Gln Glu Ala Trp Arg 215 Arg Ser Leu Arg Thr Asn Gln Ile Val 230 <210> 86 <211> 1623 <212> DNA <213> Bacillus subtilis <220> ,<221> CDS <222> (1)..(1620) <400> 86 atg tat ttg gca ttc cag gtg caa aaa ttg atg cgg tat ttg acg ctt Met Tyr Leu Ala Phe Gln Val Gln Lys Leu Met Arg Tyr Leu Thr Leu 10 tac aag ata aag gac ctg aaa tta tcg ttg ccc ggc acg aac aaa acg 96 Tyr Lys Ile Lys Asp Leu Lys Leu Ser Leu Pro Gly Thr Asn Lys Thr . 25 . cag caa ttc atg gcc caa gca gtc ggc cgt tta act gga aaa ccg gga 144 Gln Gln Phe Met Ala Gln Ala Val Gly Arg Leu Thr Gly Lys Pro Gly 35 gto gtg tta gto aca toa gga cog ggt goo tot aac ttg gca aca ggo 192 Val Val Leu Val Thr Ser Gly Pro Gly Ala Ser Asn Leu Ala Thr Gly 50 240 ctg ctg aca gcg aac act gaa gga gac cct gtc gtt gcg ctt gct gga Leu Leu Thr Ala Asn Thr Glu Gly Asp Pro Val Val Ala Leu Ala Gly 65 aac gtg atc cgt gca tat cgt tta aaa cgg aca cat caa tct ttg gat Asn Val Ile Arg Ala Tyr Arg Leu Lys Arg Thr His Gln Ser Leu Asp 90 aat gcg gcg cta ttc cag ccg att aca aaa tac agt gta gaa gtt caa 336 Asn Ala Ala Leu Phe Gln Pro Ile Thr Lys Tyr Ser Val Glu Val Gln 100 105 110 gat gta aaa aat ata ccg gaa gct gtt aca aat gca ttt agg ata gcg 384 Asp Val Lys Asn Ile Pro Glu Ala Val Thr Asn Ala Phe Arg Ile Ala 120 125 115 tca gca ggg cag gct ggg gcc gct ttt gtg agc ttt ccg caa gat gtt 432

	Ser	Ala 130	Gly	Gln	Ala	Gly	Ala 135	Ala	Phe	Val	Ser	Phe 140	Pro	Gln	Asp	Val	
												gct Ala					480
												gcg Ala					528
												ggc Gly				gga Gly	576
												ttg Leu					624
												acc Thr 220					672
												ttc Phe					720
												ctg Leu					768
,	gac Asp	ccg Pro	att Ile	gaa Glu 260	tat Tyr	gat Asp	ccg Pro	aaa Lys	ttc Phe 265	tgg Trp	aat Asn	atc Ile	aat Asn	gga Gly 270	gac Asp	cgg Arg	916
	aca Thr	att Ile	atc Ile 275	cat His	tta Leu	gac Asp	gag Glu	att Ile 280	atc Ile	gct Ala	gac Asp	att Ile	gat Asp 285	cat His	gct Ala	tac Tyr	864
												tcc Ser 300					912
	atc Ile 305	gaa Glu	cac	gat Asp	gct Ala	gtg Val 310	aaa Lys	gtg Val	gaa Glu	ttt Phe	gca Ala 315	gag Glu	cgt Arg	gag Glu	cag Gln	aaa Lys 320	960
	atc Ile	ctt Leu	tct Ser	gat Asp	tta Leu 325	aaa Lys	caa Gln	tat Tyr	atg Met	cat His 330	gaa Glu	ggt Gly	gag Glu	cag Gln	gtg Val 335	cct Pro	1008
	gca Ala	gat Asp	tgg Trp	aaa Lys 340	tca Ser	gac Asp	aga Arg	gcg Ala	cac His 345	cct Pro	ctt Leu	gaa Glu	atc Ile	gtt Val 350	aaa Lys	gag Glu	1056
	ttg Leu	cgt Arg	aat Asn 355	gca Ala	gtc Val	gat Asp	gat Asp	cat His 360	gtt Val	aca Thr	gta Val	act Thr	tgc Cys 365	gat Asp	atc Ile	ggt Gly	1104

tcg cac t Ser His S	tcc att Ser Ile	tgg atg Trp Met	tca Ser 375	cgt Arg	tat Tyr	ttc Phe	Arg	agc Ser 380	tac Tyr	gag Glu	ccg Pro	tta Leu	1152
aca tta a Thr Leu I 385	atg atc Met Ile	agt aac Ser Asn 390	ggt Gly	atg Met	caa Gln	aca Thr	ctc Leu 395	ggc Gly	gtt Val	gcg Ala	ctt Leu	cct Pro 400	1200
tgg gca a	atc ggc Ile Gly	gct tca Ala Ser 405	ttg Leu	gtg Val	aaa Lys	ccg Pro 410	gga Gly	gaa Glu	aaa Lys	gtg Val	gtt Val 415	tct Ser	1248
gtc tct (Val Ser	ggt gac Gly Asp 420	Gly Gly	ttc Phe	Leu	ttc Phe 425	tca Ser	gca Ala	atg Met	gaa Glu	tta Leu 430	gag Glu	aca Thr	1296
gca gtt Ala Val	cga cta Arg Leu 435	aaa gca Lys Ala	cca Pro	att Ile 440	gta Val	cac His	att Ile	gta Val	tgg Trp 445	aac Asn	gac Asp	agc Ser	1344
aca tat Thr Tyr 450	gac atg Asp Met	gtg cat Val His	ttc Phe 455	cag Gln	caa Gln	ttg Leu	aaa Lys	aaa Lys 460	tat Tyr	aac Asn	cgt Arg	aca Thr	1392
tct gcg Ser Ala 465	gtc gat Val Asp	ttc gga Phe Gly 470	A'sn	atc Ile	gat Asp	atc Ile	gtg Val 475	aaa Lys	tat Tyr	gcg Ala	gaa Glu	agc Ser 480	1440
ttc gga Phe Gly	gca act Ala Thr	gcg ttg Ala Lev 485	g cgc n Arg	gta Val	gaa Glu	tca Ser 490	cca Pro	gac Asp	cag Gln	ctg Leu	gca Ala 495	gat Asp	1488
gtt ctg Val Leu	cgt caa Arg Glr 500	Gly Met	aac Asn	gct Ala	gaa Glu 505	ggt Gly	cct Pro	gtc Val	atc Ile	atc Ile 510	gat Asp	gtc Val	1536
ccg gtt Pro Val	gac tac Asp Tyr 515	agt gat Ser Asp	aac Asn	att Ile 520	aat Asn	tta 'Leu	gca Ala	agt Ser	gac Asp 525	aag Lys	ctt Leu	ccg Pro	1584
aaa gaa Lys Glu 530	ttc ggg Phe Gly	g gaa cto y Glu Leo	atg Met 535	Lys	acg Thr	aaa Lys	gct Ala	ctc Leu 540					1623
<210> 87 <211> 540 <212> PRT <213> Bacillus subtilis													
<400> 8 Met Tyr 1		a Phe Gl	n Val	Gln	Lys	Leu 10		Arg	Tyr	Leu	Thr 15		
Tyr Lys	Ile Ly	s Asp Le 0	u Lys	Leu	Ser 25		Pro	Gly	Thr	Asn 30		Thr	

Gln Gln Phe Met Ala Gln Ala Val Gly Arg Leu Thr Gly Lys Pro Gly Val Val Leu Val Thr Ser Gly Pro Gly Ala Ser Asn Leu Ala Thr Gly Leu Leu Thr Ala Asn Thr Glu Gly Asp Pro Val Val Ala Leu Ala Gly Asn Val Ile Arg Ala Tyr Arg Leu Lys Arg Thr His Gln Ser Leu Asp Asn Ala Ala Leu Phe Gln Pro Ile Thr Lys Tyr Ser Val Glu Val Gln 100 Asp Val Lys Asn Ile Pro Glu Ala Val Thr Asn Aia Phe Arg Ile Ala Ser Ala Gly Gln Ala Gly Ala Ala Phe Val Ser Phe Pro Gln Asp Val 135 Val Asn Glu Val Thr Asn Thr Lys Asn Val Arg Ala Val Ala Ala Pro 150 Lys Leu Gly Pro Ala Ala Asp Asp Ala Ile Ser Ala Ala Ile Ala Lys Ile Gln Thr Ala Lys Leu Pro Val Val Leu Val Gly Met Lys Gly Gly Arg Pro Glu Ala Ile Lys Ala Val Arg Lys Leu Leu Lys Lys Vai Gln 195 200 Leu Pro Phe Val Glu Thr Tyr Gln Ala Ala Gly Thr Leu Ser Arg Asp 215. Leu Glu Asp Gln Tyr Phe Gly Arg Ile Gly Leu Phe Arg Asn Gln Pro Gly Asp Leu Leu Glu Gln Ala Asp Val Val Leu Thr Ile Gly Tyr 250 Asp Pro Ile Glu Tyr Asp Pro Lys Phe Trp Asn Ile Asn Gly Asp Arg 260 265 Thr Ile Ile His Leu Asp Glu Ile Ile Ala Asp Ile Asp His Ala Tyr 280 Gln Pro Asp Leu Glu Leu Ile Gly Asp Ile Pro Ser Thr Ile Asn His Ile Glu His Asp Ala Val Lys Val Glu Phe Ala Glu Arg Glu Gln Lys 310 315 Ile Leu Ser Asp Leu Lys Gln Tyr Met His Glu Gly Glu Gln Val Pro Ala Asp Trp Lys Ser Asp Arg Ala His Pro Leu Glu Ile Val Lys Glu

WO 01/21772 PCT/US00/25993

- 134 -

			340					345					350		
Leu	Arg	Asn 355	Ala	Val	Asp	Asp	His 360	Val	Thr	Val	Thr	Cys 365	Asp	Fle	Gly
Ser	His 370	Ser	Ile	Trp	Met	Ser 375	Arg	Tyr	Phe	Arg	Ser 380	Tyr	Glu	Pro	Leu
Thr 385	Leu	Met	Ile	Ser	Asn 390	Gly	Met	Gln	Thr	Leu 395	Gly	Val	Ala	Leu	Pro 400
Trp	Ala	Ile	Gly	Ala 405		Leu	Val	Lys	Pro 410	Gly	Glu	Lys	Val	Val 415	Ser
Val	Ser	Gly	Asp 420	Gly	Gly	Phe	Leu	Phe 425	Ser	Ala	Met	Glu	Leu 430		Thr .
Ala	Val	Arg 435	Leu	Lys	Ala	Pro	Ile 440	Val	His	Ile	Val	Trp 445	Asn	Asp	Ser
Thr	Tyr 450	Asp	Met	Val	His	₽he 455	Gln	Gln	Leu	Lys	Lys 460	Tyr	Asn	Arg	Thr
Ser 465	Ala	Val	Asp	Phe	Gly 470	Asn	Ile	Asp	Ile	Val 475	Lys	Tyr	Ala	Glu	Ser 480
Phe	Gly	Ala	Thr	Ala 485	Łeu	Arg	Va).	Glu	Ser 490	Pro	Asp	Gln	Leu	Ala 495	Asp
Val	Leu	Arg	Gln 500	_	Met	Asn	Ala	-Glu 505	Gly	Pro	Val	Ile	11e 510	Asp	Val
Pro	Val	Asp 515	_	Ser	Asp	Asn	11e 520		Leu	Alə	Ser	Asp 525		Leu	Pro
Lys	Glu 530		Gly	Glu	Leu	Met 535		Thr	Lys	Ala	Leu 540				
<210> 88 <211> 23 <212> DNA <213> Artificial Sequence															
<220> <223> Description of Artificial Sequence: ribosome binding site															
<22 <22		All c	occur	renc	es o	f n	indi	cate	any	nuc	leot	ide			
	00> 8 aagg		tgar	ınnnr	ınn a	tg									

23

<210> 89 <211> 7 <212> PRT

BNSDOCID: <WO 0121772A2 1 >

```
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: PanC
      C terminus
<400> 89
Ile Arg Glu Met Glu Arg Ile
  1
<210> 90
<211> 5
<212> PRT
<213> Artificial Sequence
<223> Description of Artificial Sequence: PanC
      C terminus
<400> 90
Ile Arg Glu Arg Arg
, 1
<210> 9i
<211> 7
<212> PRT
<213> Artificial Sequence
<223> Description of Artificial Sequence: PanC
      C terminus
<400> 91
Ile Arg Arg Lys Glu Val Asn
<210> 92
<211> 6688
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Recombinant
      pAN336 plasmid
<400> 92
tgcgccgcta cagggcgcgt ccattcgcca ttcaggctgc gcaactgttg ggaagggcga 60
tcggtgcggg cctcttcgct attacgccag ctggcgaaag ggggatgtgc tgcaaggcga 120
ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagtgaa 180
ttgtaatacg actcactata gggcgaattg ggcccgacgt cgcatgcacc aggcttctca 240
ggcgctgact tagaaaacct cttgaatgaa gctgcgcttg tagcggctcg tcaaaacaag 300
```

aaaaaaatcg atgcgcgtga tattgacgaa gcgacggacc gtgtaattgc cggacccgct 360 aagaagagcc gcgttatctc caagaaagaa cgcaatatcg tggcttatca cgaaggcgga 420 cacaccetta tegeteteet titagateae geagatatee ticataaaet aaceattett 480 cctcggggcc aggctggcgg ttatgctgtt atgctgccaa gagaagaccg ttatttccaa 540 acaaagccgg agctgcttga taaaattgtc ggcctcttgg gcggacgtgt tgctgaagag 600 attatcttcg gtgaagtcag cacaggggcg cacaatgact tccagcgtgc gacgaatatt 660 gcaagacgaa tggttacaga attcggtatg tcagaaaaac tgggaccgtt gcaatttgga 720 cagteteagg geggteaggt attettagge egtgatttea acaacgaaca gaactacagt 780 gatcaaatcg cttacgaaat tgatcaggaa attcagcgca tcatcaaaga atgttatgag 840 cgtgcgaaac aaatcctgac tgaaaatcgt gacaagcttg aattgattgc ccaaacgctt 900 ctgaaagttg aaacgcttga cgctgaacaa atcaaacacc ttatcgatca tggaacatta 960 cctgagcgta atttctcaga tgatgaaaag aacgatgatg tgaaagtaaa cattctgaca 1020 aaaacagaag aaaagaaaga cgatacgaaa gagtaattcg ctttctttct aaaaaaactg 1080 coggetgacg etggcagttt ttttatgtaa atgattgget cagetgegge ttttacaate 1140 atccaattct ggtatcgatt tgtttacaaa tgagccgctg atcgtgtatg gtattgtaga 1200 atgtttgtaa aaagtaaagt agagaaacta ttcaaaagtg gtgatagagg ttgttactgg 1260 ttatcgatgt ggggaacacc ctgcagctcg agtgaaatac cgcacagatg cgtaaggaga 1320 aaataccgca tcaggcgata aacccagcga accatttgag gtgataggta agattatacc 1380 gaggtatgaa aacgagaatt ggacctttac agaattactc tatgaagcgc catatttaaa 1440 aagctaccaa gacgaagagg atgaagagga tgaggaggca gattgccttg aatatattga 1500 caatactgat aagataatat atcttttata tagaagatat cgccgtatgt aaggatttca 1560 gggggcaagg cataggcagc gcgcttatca atatatctat agaatgggca aagcataaaa 1620 acttgcatgg actaatgctt gaaacccagg acaataacct tatagcttgt aaattctatc 1680 ataattgtgg tttcaaaatc ggctccgtcg atactatgtt atacgccaac tttcaaaaca 1740 actttgaaaa agctgttttc tggtatttaa ggttttagaa tgcaaggaac agtgaattgg 1800 agttcgtctt gttataatta gcttcttggg gtatctttaa atactgtaga aaagaggaag 1860 gaaataataa atggctaaaa tgagaatatc accggaattg aaaaaactga tcgaaaaata 1920 ccgctgcgta aaagatacgg aaggaatgtc tcctgctaag gtatataagc tggtgggaga 1980 aaatgaaaac ctatatttaa aaatgacgga cagccggtat aaagggacca cctatgatgt 2040

ggaacgggaa aaggacatga tgctatggct ggaaggaaag ctgcctgttc caaaggtcct 2100 gcactttgaa cggcatgatg gctggagcaa tctgctcatg agtgaggccg atggcgtcct 2160 ttgctcggaa gagtatgaag atgaacaaag ccctgaaaag attatcgagc tgtatgcgga 2220 gtgcatcagg ctctttcact ccatcgacat atcggattgt ccctatacga atagcttaga 2280 cagccgctta gccgaattgg attacttact gaataacgat ctggccgatg tggattgcga 2340 aaactgggaa gaagacactc catttaaaga teegegegag etgtatgatt ttttaaagae 2400, ggaaaagccc gaagaggaac ttgtcttttc ccacggcgac ctgggagaca gcaacatctt 2460 tgtgaaagat ggcaaagtaa gtggctttat tgatcttggg agaagcggca gggcggacaa 2520 gtggtatgac attgccttct gcgtccggtc gatcagggag gatatcgggg aagaacagta 2580 tgtcgagcta ttttttgact tactggggat caagcctgat tgggagaaaa taaaatatta 2640 tattttactg gatgaattgt tttagtacct agatttagat gtctaaaaag ctttaactac 2700 aagettttta gacatetaat ettttetgaa gtacateege aactgteeat actetgatgt 2760 tttatatett ttetaaaagt tegetagata ggggteeega gegeetaega ggaatttgta 2820 togecattog coattoagge tgcgcaactg ttyggaaggg cgatcggtgc ggtcgactgg 2880 caygcaaaac aggacccaag gtcattgcga caggaggcct ggcgccgctc attgcgaacy 2940 aatcagattg tatagacatc gttgatccat tcttaaccct aaaagggctg gaattgattt 3000 atqaaagaaa ccgcgtagga agtgtatagg aggtttagta atggattatt tagtaaaagc 3060 acttycgtat gacggaaaag ttcgggctta tgcagcgaga acgactgata tggtaaatga 3120 ggggcagaga cgccatggta cgtggccgac agcatccgct gcactaggcc gtacaatgac 3180 agetteaett atgeteggeg etatgetgaa gggegatgat aagetgaeeg tgaaaatega 3240 gggcggaggt ccgatcggag ctattgtagc tgatgccaat gccaaaggag aagtcagagc 3300 ctatgtctct aacccgcaag ttcattttga tttaaatgaa caaggtaagc ttgatgtcag 3360 acgtgcggtt ggaacaaacg gaacgttaag tgtcgtaaaa gatttaggtt tgcgcgagtt 3420 cttcacagga caagtagaaa tcgtttcagg agaattagga gatgatttta cttactatct 3480 tgtgtcatct gagcaggttc cttcatcagt gggcgtaggt gtgctcgtaa atcctgacaa 3540 taccattctt gcggcagggg gctttattat tcagctgatg ccgggaacag atgatgaaac 3600 aatcacaaaa attgaacagc gtctatctca agtagagccg atttctaagc tcatccaaaa 3660 agggctgaca ccagaagaaa ttttagaaga agtcctaggc gagaaacctg agattttgga 3720 aacgatgeet gteagattee attgeeettg tteaaaagaa eggttegaaa eageeatttt 3780

aggactaggc aaaaaagaaa ttcaagatat gatagaagaa gatggacaag ccgaagcagt 3840 atgccatttt tgtaatgaaa agtacttatt tacaaaagaa gagctggaag ggcttcgtga 3900 ccaaactacc cgctaagctc tttagcgggt ttttaatttg agaaaagggg ctgaaagcag 3960 gtttgaaatc aagaacaatc tggacgcgtt ggatgcatag cttgagtatt ctatagtgtc 4020 acctaaatag cttggcgtaa tcatggtcat agctgtttcc tgtgtgaaat tgttatccgc 4080 tcacaattcc acacaacata cgagccggaa gcataaagtg taaagcctgg ggtgcctaat 4140 gagtgageta acteacatta attgegttge geteactgee egettteeag tegggaaace 4200 tgtcgtgcca gctgcattaa tgaatcggcc aacgcgcggg gagaggcggt ttgcgtattg 4260 ggcgctcttc cgcttcctcg ctcactgact cgctgcgctc ggtcgttcgg ctgcggcgag 4320 eggtateage teacteaaag geggtaatae ggttateeac agaateaggg gataaegeag 4380 gaaagaacat gtgagcaaaa ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc 4440 tygcgttttt cgataggete egececetg acgageatea caaaaatega egeteaagte 4500 agaggtggcg aaacccgaca ggactataaa gataccaggc gtttccccct ggaagctccc 4560 tegtgegete teetgtteeg accetgeege traceggata eetgteegee ttteteeett 4620 cyggzagcgt ggcgctttct catagctcac gctgtaggta tctcagttcg gtgtaggtcg 4680 ttogotocaa gotgggotgt gtgcacgaac coccegttca geoegaccgc tgcgccttat 4740 coggtaacta togtottgag tocaaccogg taagacacga ettatogoca otggcagcag 4800 ccactggtaa caggattagc agagcgaggt atgtaggcgg tgctacagag ttcttgaagt: 4860 ggtggcctaa ctacggctac actagaagga cagtatttgg tatctgcgct ctgctgaagc 4920 cagttacctt cggaaaaaga gttggtagct cttgatccgg caaacaaacc accgctggta 4980 gcggtggttt ttttgtttgc aagcagcaga ttacgcgcag aaaaaaagga tctcaagaag 5040 atcetttgat ettitetaeg gggtetgaeg eteagtggaa egaaaaetea egitaaggga 5100 ttttggtcat gagattatca aaaaggatct tcacctagat ccttttaaat taaaaatgaa 5160 gttttaaatc aatctaaagt atatatgagt aaacttggtc tgacagttac caatgcttaa 5220 tcagtgaggc acctatctca gcgatctgtc tatttcgttc atccatagtt gcctgactcc 5280 ccgtcgtgta gataactacg atacgggagg gcttaccatc tggccccagt gctgcaatga 5340 taccgcgaga cccacgctca ccggctccag atttatcagc aataaaccag ccagccggaa 5400 gggccgagcg cagaagtggt cctgcaactt tatccgcctc catccagtct attaattgtt 5460 gccgggaagc tagagtaagt agttcgccag ttaatagttt gcgcaacgtt gttggcattg 5520

ctacaggeat egtggtgtea egetegtegt ttggtatgge tteatteage teeggtteee 5580 aacgatcaag gcgagttaca tgatccccca tgttgtgcaa aaaagcggtt agctccttcg 5640 gtcctccgat cgttgtcaga agtaagttgg ccgcagtgtt atcactcatg gttatggcag 5700 cactgcataa ttetettaet gteatgeeat eegtaagatg ettteetgtg actggtgagt 5760 actcaaccaa gtcattctga gaataccgcg cccggcgacc gagttgctct tgcccggcgt 5820 caatacggga taatagtgta tgacatagca gaactttaaa agtgctcatc attggaaaac 5880 gttcttcggg gcgaaaactc tcaaggatct taccgctgtt gagatccagt tcgatgtaac 5940 ccactcgtgc acccaactga tcttcagcat cttttacttt caccagcgtt tctgggtgag 6000 caaaaacagg aaggcaaaat gccgcaaaaa agggaataag ggcgacacgg aaatgttgaa 6060 tactcatact cttccttttt caatattatt gaagcattta tcagggttat tgtctcatga 6120 gcggatacat atttgaatgt atttagaaaa ataaacaaat aggggttccg cgcacatttc 6180 cccgaaaagt gccacctgta tgcggtgtga aataccgcac agatgcgtaa ggagaaaata 6240 ccgcatcagg cgaaattgta aacgttaata ttttgttaaa attcgcgtta aatatttgtt 6300 aaatcagete attttttaac caataggeeg aaateggeaa aatceettat aaatcaaaag 6360 aatagaccga gatagggttg agtgttgttc cagtttggaa caagagtcca ctattaaaga 6420 acgtggactc caacgtcaaa gggcgaaaaa ccgtctatca gggcgatggc ccactacgtg 6480 aaccatcacc caaatcaagt tttttgcggt cgaggtgccg taaagctcta aatcggaacc 6540 ctaaagggag cccccgattt agagcttgac ggggaaagcc ggcgaacgtg gcgagaaagg 6600 aagggaagaa agcgaaagga gcgggcgcta gggcgctggc aagtgtagcg gtcacgctqc 6660 gcgtaaccac cacacccgcc gcgcttaa 6688

<210> 93

<211> 8503

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN004 plasmid

<400> 93

gaattttgcg gccgcttcga aagctgtaat ataaaaacct tcttcaacta acggggcagg 60 ttagtgacat tagaaaaccg actgtaaaaa gtacagtcgg cattatctca tattataaaa 120 gccagtcatt aggcctatct gacaattcct gaatagagtt cataaacaat cctgcatgat 180 aaccatcaca aacagaatga tgtacctgta aagatagcgg taaatatatt gaattacctt 240 tattaatgaa ttttcctgct gtaataatgg gtagaaggta attactatta ttattgatat 300 ttaagttaaa cccagtaaat gaagtccatg gaataataga aagagaaaaa gcattttcag 360 gtataggtgt tttgggaaac aatttccccg aaccattata tttctctaca tcagaaaggt 420 ataaatcata aaactetttg aagteattet ttacaggagt ecaaatacca gagaatgttt 480 tagatacacc atcaaaaatt gtataaagtg gctctaactt atcccaataa cctaactctc 540 cgtcgctatt gtaaccagtt ctaaaagctg tatttgagtt tatcaccctt gtcactaaga 600 aaataaatgc agggtaaaat ttatatcctt cttgttttat gtttcggtat aaaacactaa 660 tatcaatttc tgtggttata ctaaaagtcg tttgttggtt caaataatga ttaaatatct 720 cttttctctt ccaattqtct aaatcaattt tattaaagtt catttgatat gcctcctaaa 780 tttttatcta aagtgaattt aggaggetta ettgtetget ttetteatta gaatcaatee 840 ttttttaaaa gtcaatatta ctgtaacata aatatatatt ttaaaaaatat cccactttat 900 ccaattttcg tttgttgaac taatgggtgc tttagttgaa gaataaagac cacattaaaa 960 aatgtggtct tttgtgtttt tttaaaggat ttgagcgtag cgaaaaatcc ttttctttct 1020 tatcttgata ataagggtaa ctattgaatt cggtaccaag agtttgtaga aacgcaaaaa 1080 ggccatccgt caggatggcc ttctgcttaa tttgatgcct ggcagtttat ggcgggcgtc 1140 ctgcccgcca ccctccgggc cgttgcttcg caacgttcaa atccgctccc ggcggatttg; 1200 tectacteag gagagegtte accgacaaac aacagataaa acgaaaggee cagtettteg 1260 actgageett tegttttatt tgatgeetgg eagtteeeta etetegeatg gggagaeece 1320 acactaccat eggegetacg gegttteact tetgagtteg geatggggte aggtgggace 1380 accgcgctac tgccgccagg caaattctgt tttatcagac cgcttctgcg ttctgattta 1440 atotgtatca ggotgaaaat ottototoat cogocaaaac aggatootac ggaaatggag 1500 cggcaaaacc gttttactct caaaatctta aaagaaaacc cccgataaag ggggcttttc 1560 ttctacaaaa ttgtacgggc tggttcgttc cccagcattt gttcaatttt gttttgatca 1620 ttcagaacag ccactttcgg ctcatggctt gccgcttctt gatcagacat cattttgtag 1680 qaaataataa tgaccttatc tccttcctgc acaaggcgtg cggctgcacc gtttaagcat 1740 atgacgccgc ttccccgttt accaggaata atatacgttt caagacgtgc tccattatta 1800 ttattcacaa tttgtacttt ttcattagga agcattccca cagcatcaat gagatcttca 1860 tcaattgtaa tgcttcccac atagttcagg tttgcttccg taacagttgc cctgtgaagt 1920 ttqccqctca tcattqttcq atacatatta tattctctcc atttctcqaa tatcaataat 1980

gatattatct attaaacgcg cttttgaaaa agcaactgca acagcgagaa tcatctttcc 2040 agcaatttca ttcacaggct cgagttccgg ataggaataa agctctacat agtctatggt 2100 tecgetagte gttteadtga tatettttge agettttate accgetteag gatetettte 2160 accggcttgg acaagttccg cacttgtttg aagggcccga tacagcttag gcgcttcttt 2220 tettteetea getgttaagt atacattgeg agagettttg getaageegt etteetet 2280 gaccgtatcg acaggaacca attcaatatc catgaagaag tcgctgatta acccatcaac 2340 aacagctace tgctgcgcat cttttaaacc gaaataggca cgagtcggct tgactagatt 2400 gaaaagette gteagtaega tegegaeece gteaaaatgt eettetettg agegeeegea 2460 taacacgtct gtgcgtcttt ctacatgaat cgtgacattc ttttcaccgg gatacatatc 2520 atgagcatet ggegtaaaaa gaatategae teeggegttt tetgeadgag etgeateeeg 2580 ctcaatatcg cgcggatatg cttcaaaatc ttcattaggg ccgaattgtg caggattcac 2640 aaaaatactc ataataacgg cgtcgttttc ttgtcttgct ttgtctgcta aggttaaatg 2700 cccctcatgc agaaacccca tcgtcggaac aaatccgatt gacttgccct ctgaatggta 2760 ttgttttatg gettetttea getgtgaaat atcagtaate tgteteatet tatttteee 2820. cgtacaagcc gtcaagcact gtctggttca tttgaaagga atgcttttgt tcagggaaag 2880 cacgatgtct tacatcctga acatatccgc tgattgctgt ttcgatggtt tcatcaatgc 2940 gcgtatattg ctttacaaat ttaggtgttc tctcaacacc gtggccgata atatcatgat. 3000 aaacgagaac ttgtccgtcc gctttcacac cagccccgat tccaatgacc ggtatgctta 3060 gcgtctcggc aattttggct gtgagttctg ccggcacaca ttccagcaca agcatcatag 3120 ctcctgcttc ttcgcatttt atactgtctt ctattaattt tttggcgctt tgttcgtctt 3180 tgccctgtac tttatagccg cccagtacgc cgactgactg cggtgtcaaa cctaagtgac 3240 tgactactgg aatgcctcca agcgtcaatg cgcgaatgga ttcaaacacg ccttctccgc 3300 cctcaagctt cagtgcgtca gctccgcttt cctgaacgat agccgctgca tttttcagcg 3360 tatetteett agacaggtga taagacataa aeggeatate tgteacaata aaggtatteg 3420 gcgcacccct tttaacggct tttgtatgat ggatcatgtc cgcaactgtc acaccgacag 3480 ttgaatcaag gccgaggacg accattccaa gtgaatcacc gactaaaatc atgtcaactc 3540 ccgcttgttc agcaagttta gctgccggat aatcataagc ggtcagcatg acaatcggtt 3600 gagcgtcctg ctgttgttaa gattattata ccacaccttg tagataaagt caacaacttt 3720

ttgcaaaatt tttcaggaat tttagcagag gttgttctgg atgtagaaca aaacatcttt 3780 ccgctcttgt gctgttagga tatctttctt ggaagctagg taggcctcga gttatggcag 3840 ttggttaaaa ggaaacaaaa agaccgtttt cacacaaaac ggtcttttc gatttctttt 3900 tacagtcaca gccacttttg caaaaaccgg acagcttcat gccttataac tgctgtttcg 3960 gtcgacaagc ttcgcgaagc ggccgcaaaa ttcactggcc gtcgttttac aacgtcgtga 4020 ctgggaaaac cctggcgtta cccaacttaa tcgccttgca gcacatcccc ctttcgccag 4080 ctggcgtaat agcgaagagg cccgcaccga tcgcccttcc caacagttgc gcagcctgaa 4140 tggcgaatgg cgcctgatgc ggtattttct ccttacgcat ctgtgcggta tttcacaccg 4200 catatggtgc actotcagta caatotgoto tgatgccgca tagttaagcc agccccgaca 4260 cccgccaaca cccgctgact atgcttgtaa accgttttgt gaaaaaaattt ttaaaaataaa 4320 aaaggggacc tctagggtcc ccaattaatt agtaatataa tctattaaag gtcattcaaa 4380 aggtcatcca ccggatcagc ttagtaaagc cctcgctaga ttttaatgcg gatgttgcga 4440 ttacttcgcc aactattgcg ataacaagaa aaagccagcc tttcatgata tatctcccaa 4500 tttgtgtagg gcttattatg cacgcttaaa aataataaaa gcagacttga cctgatagtt 4560 tggctgtgag caattatgtg cttagtgcat ctaacgcttg agttaagccg cgccgcgaag 4620 cggcgtcggc ttgaacgaat tgttagacat tatttgccga ctaccttggt gatctcgcct 4680 ttcacgtagt ggacaaattc ttccaactga tctgcgcgcy aggccaagcg atcttcttct 4740 tgtccaagat aagcctgtct agcttcaagt atgacgggct gatactgggc cggcaggcgc 4800 tccattgccc agtcggcagc gacatcettc ggcgcgattt tgccggttac tgcgctgtac 4860 caaatgeggg acaaegtaag cactacattt egeteatege eageocagte gggeggegag 4920 ttccatagcg ttaaggtttc atttagcgcc tcaaatagat cctgttcagg aaccggatca 4980 aagagtteet eegeegetgg acetaceaag geaaegetat gttetettge tittgteage 5040 aagatagcca gatcaatgtc gatcgtggct ggctcgaaga tacctgcaag aatgtcattg 5100 cgctgccatt ctccaaattg cagttcgcgc ttagctggat aacgccacgg aatgatgtcg 5160 tcgtgcacaa caatggtgac ttctacagcg cggagaatct cgctctctcc aggggaagcc 5220 gaagtttcca aaaggtcgtt gatcaaagct cgccgcgttg tttcatcaag ccttacggtc 5280 accytaacca gcaaatcaat atcactytyt gycttcagyc cyccatccac tycgygayccy 5340 tacaaatgta cggccagcaa cgtcggttcg agatggcgct cgatgacgcc aactacctct 5400 gatagttgag tegataette ggegateace getteeetea tgatgtttaa etttgtttta 5460

gggcgactgc cctgctgcgt aacatcgttg ctgctccata acatcaaaca tcgacccacg 5520 gcgtaacgcg cttgctgctt ggatgcccga ggcatagact gtaccccaaa aaaacagtca 5580 taacaagcca tgaaaaccgc cactgcgccg ttaccaccgc tgcgttcggt caaggttctg 5640 gaccagttgc gtgagcgcat acgctacttg cattacagct tacgaaccga acaggcttat 5700 gtccactggg ttcgtgcctt catccgtttc cacggtgtgc gtcacccggc aaccttgggc 5760 agcagcgaag tegaggeatt tetgteetgg etggegaaeg agegeaaggt tteggtetee 5820 acgeategte aggeattgge ggeettgetg ttettetaeg geaaggtget gtgeaeggat 5880 ctgccctggc ttcaggagat cggaagacct cggccgtcgc ggcgcttgcc ggtggtgctg 5940 acceeggatg aagtggtteg cateeteggt tttetggaag gegageateg tttgttegee 6000 cagettetgt atggaacggg catgeggate agtgagggtt tgcaactgeg ggtcaaggat 6060 ctggatttcg atcacggcac gatcatcgtg cgggagygca agggctccaa ggatcgggcc 6120 ttgatgttac ccgagagett ggcacccage ctgcgcgage aggggaattg atccggtgga 6180 tgaccttttg aatgaccttt aatagattat attactaatt aattggggac cctagaggtc 6240 contitue terraaaaat tettecanaa aanggettan aagnataang ggettegetg 6300 cccgcaaacg ggctgttctg gtgttgctag tttgttatca gaatcgcaga tccggcttca 6360 ggtttgccgg ctgaaagcgc tatttcttcc agaattgcca tgattttttc cccacgggag 6420 gegteactgg etceegtgtt gteggeaget ttgattegat aageageate geetgtttea. 6480 ggctgtctat gtgtgactgt tgagctgtaa caagttgtct caggtgttca atttcatgtt 6540 ctagttgctt tgttttactg gtttcacctg ttctattagg tgttacatgc tgttcatctg 6600 ttacattgtc gatctgttca tggtgaacag ctttaaatgc accaaaaact cgtaaaagct 6660 ctgatgtatc tatctttttt acaccgtttt catctgtgca tatggacagt tttccctttg 6720 atatctaacg gtgaacagtt gttctacttt tgtttgttag tcttgatgct tcactgatag 6780 atacaagage cataagaace teagateett eegtatttag eeagtatgtt etetagtgtg 6840 gttcgttgtt tttgcgtgag ccatgagaac gaaccattga gatcatgctt actttgcatg 6900 tcactcaaaa attttgcctc aaaactggtg agctgaattt ttgcagttaa agcatcgtgt 6960 agtgtttttc ttagtccgtt acgtaggtag gaatctgatg taatggttgt tggtattttg 7020 tcaccattca tttttatctg gttgttctca agttcggtta cgagatccat ttgtctatct 7080 agttcaactt ggaaaatcaa cgtatcagtc gggcggcctc gcttatcaac caccaatttc 7140 atattgctgt aagtgtttaa atctttactt attggtttca aaacccattg gttaagcctt 7200

ttaaactcat ggtagttatt ttcaagcatt aacatgaact taaattcatc aaggctaatc 7260 tctatatttg ccttgtgagt tttcttttgt gttagttctt ttaataacca ctcataaatc 7320 ctcatagagt atttgttttc aaaagactta acatgttcca gattatattt tatgaatttt 7380 tttaactgga aaagataagg caatatotot toactaaaaa ctaattotaa tttttogott 7440 gagaacttgg catagtttgt ccactggaaa atctcaaagc ctttaaccaa aggattcctg 7500 atttccacag ttctcgtcat cagctctctg gttgctttag ctaatacacc ataagcattt 7560 tecetactga tgtteateat etgagegtat tggttataag tgaacgatae egteegttet 7620 ttccttgtag ggttttcaat cgtggggttg agtagtgcca cacagcataa aattagcttg 7680 qtttcatqct ccgttaagtc atagcgacta atcgctagtt catttgcttt gaaaacaact 7740 aattcagaca tacatctcaa ttggtctagg tgattttaat cactatacca attgagatgg 7800 gctagtcaat gataattact agtccttttc ctttgagttg tgggtatctg taaattctgc 7860 tagacettig etggaaaact igtaaattei getagaeeet eigtaaatte egetagaeet 7920 ttgtgtgttt tttttgttta tattcaagtg gttataattt atagaataaa gaaagaataa 7980 aaaaagataa aaagaataga toocagooot gtgtataaot cactaottta gtcagttoog 8040 cactattaca aaaggatgtc gcaaacgctg tttgctcctc tacaaaacag accttaaaac 8100 cctaaaggct taagtagcac cctcgcaagc tcgggcaaat cgctgaatat tccttttgtc 3160 tecgaccate aggeacetga gtegetgtet tittegtgae atteagtteg etgegeteae 8220 ggctctggca gtgaatgggg gtaaatggca ctacaggcgc cttttatgga ttcatgcaag 8280 gaaactaccc ataatacaag aaaagcccgt cacgggcttc tcagggcgtt ttatggcggg 8340 tctgctatgt ggtgctatct gactttttgc tgttcagcag ttcctgccct ctgattttcc 8400 agtotgacca ottoggatta tocogtgaca ggtoattoag actggotaat gcacccagta 8460 8503 aggrageget atcatcaaca ggcttacceg tettactgte aac

<210> 94

<211> 7381

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:recombinant pAN006 plasmid

<400> 94
ttqcqqccqc ttcqaaagct gtaatataaa aaccttcttc aactaacggg gcaggttagt 60

gacattagaa aaccgactgt aaaaagtaca gtcggcatta tctcatatta taaaagccag 120 tcattaggcc tatctgacaa ttcctgaata gagttcataa acaatcctgc atgataacca 180 tcacaaacag aatgatgtac ctgtaaagat agcggtaaat atattgaatt acctttatta 240 atgaattttc ctgctgtaat aatgggtaga aggtaattac tattattatt gatatttaag 300 ttaaacccag taaatgaagt ccatggaata atagaaagag aaaaagcatt ttcaggtata 360 ggtgttttgg gaaacaattt ccccgaacca ttatatttct ctacatcaga aaggtataaa 420 tcataaaact ctttgaagtc attctttaca ggagtccaaa taccagagaa tgttttagat 480 acaccatcaa aaattgtata aagtggctct aacttatccc aataacctaa ctctccgtcg 540 ctattgtaac cagttctaaa agctgtattt gagtttatca cccttgtcac taagaaaata 600 aatgcagggt aaaatttata toottottgt tttatgttto ggtataaaac actaatatca 660 atttctgtgg ttatactaaa agtcgtttgt tggttcaaat aatgattaaa tatctctttt 720 ctcttccaat tgtctaaatc aattttatta aagttcattt gatatgcctc ctaaattttt 780 atctaaagtg aatttaggag gcttacttgt ctgctttctt cattagaatc aatccttttt 840 taaaagtcaa tattactgta acataaatat atattttaaa aatatcccac tttatccaat 900 tttcgtttgt tgaactaatg ggtgctttag ttgaagaata aagaccacat taaaaaatgt 960 ggtcttttgt gttttttaa aggatttgag cgtagcgaaa aatccttttc tttcttatct 1020 tgataataag ggtaactatt gaatteggta ceaagagttt gtagaaaege aaaaaggeea 1080 tccgtcagga tggccttctg cttaatttga tgcctggcag tttatggcgg gcgtcctgcc 1140 cgccaccctc cgggccgttg cttcgcaacg ttcaaatccg ctcccggcgg atttgtccta 1200 ctcaggagag cgttcaccga caaacaacag ataaaacgaa aggcccagtc tttcgactga 1260 gcctttcgtt ttatttgatg cctggcagtt ccctactctc gcatggggag accccacact 1320 accateggeg ctaeggegtt teacttetga gtteggeatg gggteaggtg ggaceaeege 1380 gctactgccg ccaggcaaat tctgttttat cagaccgctt ctgcgttctg atttaatctg 1440 tatcaggctg aaaatcttct ctcatccgcc aaaacaggat cctacggaaa tggagcggca 1500 aaaccgtttt actctcaaaa tcttaaaaga aaacccccga taaagggggc ttttcttcta 1560 caaaattgta cgggctggtt cgttccccag catttgttca attttgtttt gatcattcag 1620 aacagccact ttcggctcat ggcttgccgc ttcttgatca gacatcattt tgtaggaaat 1680 aataatgace ttateteett eetgeacaag gegtgegget geacegttta ageatatgae 1740 gccgcttccc cgtttaccag gaataatata cgtttcaaga cgtgctccat tattattatt 1800

cacaattigt actititeat taggaageat teccaeagea teaatgagat etteateaat 1860 tgtaatgctt cccacatagt tcaggtttgc ttccgtaaca gttgccctgt gaagtttgcc 1920 gctcatcatt gttcgataca tattatattc tctccatttc tcgaatatca ataatgatat 1980 tatetattaa aegegetttt gaaaaageaa etgeaaeage gagaateate ttteeageaa 2040 tttcattcac aggetegagt teeggatagg aataaagete tacatagtet atggtteege 2100. tagtcgtttc aatgatatct tttgcagctt ttatcaccgc ttcaggatct ctttcaccgg 2160 cttggacaag ttccgcactt gtttgaaggg cccgatacag cttaggcgct tctttcttt 2220 cctcagctgt taagtataca ttgcgagagc ttttggctaa gccgtcttcc tctctgaccg 2280 tatcgacagg aaccaattca atatccatga agaagtcgct gattaaccca tcaacaacag 2340 ctacctgctg cgcatctttt aaaccgaaat aggcacgagt cggcttgact agattgaaaa 2400 gettegteag tacgategeg acceegteaa aatgteette tettgagege eegeataaca 2460 cgtctgtgcg tctttctaca tgaatcgtga cattcttttc accgggatac atatcatgag 2520 catotggogt aaaaagaata togactoogg ogttttotgc aagagotgca tocogotcaa 2580 tategegegy atatgettea aaatetteat tagggeegaa ttgtgeagga tteacaaaaa 2640 tactcataat aacggcgtcg tittcttgtc tigctitgic igctaaggit aaaigcccci 2700 catgoagaaa coccatogto ggaacaaato ogattgactt goodtotgaa tggtattgtt 2760 ttatggette tttcagetgt gaaatateag taatetgtet catettattt teeceegtae 2820 aageegteaa geactgtetg gtteatttga aaggaatget tttgtteagg gaaageacga 2880 tgtcttacat cctgaacata tccgctgatt gctgtttcga tggtttcatc aatgcgcgta 2940 tattgcttta caaatttagg tgttctctca acaccgtggc cgataatatc atgataaacg 3000 agaacttgtc cgtccgcttt cacaccagcc ccgattccaa tgaccggtat gcttagcgtc 3060 teggeaattt tggetgtgag ttetgeegge acacatteea geacaageat catageteet 3120 gettettege attituatet gtettetatt aattitttgg egettigtte gtetttgeee 3180 tgtactttat agccgcccag tacgccgact gactgcggtg tcaaacctaa gtgactgact 3240 actggaatgc ctccaagcgt caatgcgcga atggattcaa acacgccttc tccgccctca 3300 agetteagtg egteagetee gettteetga acgatageeg etgeattttt eagegtatet 3360 tccttagaca ggtgataaga cataaacggc atatctgtca caataaaggt attcggcgca 3420 coccttttaa cggcttttgt atgatggatc atgtccgcaa ctgtcacacc gacagttgaa 3480 tcaaggccga ggacgaccat tccaagtgaa tcaccgacta aaatcatgtc aactcccgct 3540

tgttcagcaa gtttagctgc cggataatca taagcggtca gcatgacaat cggttcttca 3600 gactccttca tttttagaaa atccagtttt gttttcatgt tttctcctcc tctagagcgt 3660 cctgctgttg ttaagattat tataccacac cttgtagata aagtcaacaa ctttttgcaa 3720 aatttttcag gaattttagc agaggttgtt ctggatgtag aacaaaacat ctttccgctc 3780 ttgtgctgtt aggatatett tettggaage taggtaggee tegagttatg geagttggtt 3840 aaaaggaaac aaaaagaccg ttttcacaca aaacggtctt tttcgatttc tttttacagt 3900 cacagecaet titgeaaaaa eeggaeaget teatgeetta taaetgetgt tieggiegae 3960 ctgcaggcat gcaagctteg cgaageggee geegaegega ggetggatgg cetteeceat 4020 tatgattett etegetteeg geggeategg gatgeeegeg ttgeaggeea tgetgteeag 4080 gcaygtagat gacgaccatc agggacagct tcaaggatcg ctcgcgqctc ttaccagcct 4140 aacttegate aetggaeege tgategteae ggegatttat geegeetegg egageaeatg 4200 gaacgggttg gcatggattg taggcgccgc cctatacctt gtctgcctcc ccgcgttgcg 4260 tegeggtgca tggageeggg ceaectegae etgaatggaa geeggeggea eetegetaae 4320 gyatteacea etecaagaat tggageeaat eaattetige ggagaactgt gaatgegeaa 4380 accaaccett ggcagaacat atccategeg teegecatet ccageageeg caegeggege 4440 atotogggca goyttgggto otggocaogg gtgogoatga togtgotoot gtogttgaggg 4500 acceggetag getggegggg ttgeettact ggttageaga atgaateace gataegegag 4560 egaaegtyaa gegaetgetg etgeaaaaeg tetgegaeet gageaacaae atgaatggte 4620 ttcggtttcc gtgtttcgta aagtctggaa acgcggaagt cagcgccctg caccattatg 4680 ttccggatct gcatcgcagg atgctgctgg ctaccctgtg gaacacctac atctgtatta 4740 acgaageget ggeattgace etgagtgatt tttetetggt ecegeegeat ccatacegee 4800 agttgtttac cctcacaacg ttccagtaac cgggcatgtt catcatcagt aacccgtatc 4860 gtgagcatcc tetetegttt categgtate attacececa tgaacagaaa tteeceetta 4920 cacggaggca tcaagtgacc aaacaggaaa aaaccgccct taacatggcc cgctttatca 4980 gaagccagac attaacgctt ctggagaaac tcaacgagct ggacgcggat gaacaggcag 5040 acatetgtga ategetteae gaceaegetg atgagettta eegeagetge etegegegtt 5100 teggtgatga eggtgaaaae etetgaeaea tgeageteee ggagaeggte acagettgte 5160 tgtaagcgga tgccgggagc agacaagccc gtcagggcgc gtcagcgggt gttggcgggt 5220 gtcggggcgc agccatgacc cagtcacgta gcgatagcgg agtgtatact ggcttaacta 5280

tgeggcatca gagcagattg tactgagagt gcaccatatg eggtgtgaaa tacegcacag 5340 atgcgtaagg agaaaatacc gcatcaggcg ctcttccgct tcctcgctca ctgactcgct 5400 gcgctcggtc gttcggctgc ggcgagcggt atcagctcac tcaaaggcgg taatacggtt 5460 atccacagaa tcaggggata acgcaggaaa gaacatgtga gcaaaaggcc agcaaaaggc 5520 caggaaccgt aaaaaggeeg egttgetgge gttttteeat aggeteegee eeeetgaega 5580 gcatcacaaa aatcgacgct caagtcagag gtggcgaaac ccgacaggac tataaagata 5640 ccaggogttt ccccctggaa getecctegt gegetetect gtteegaeee tgeegettae 5700 cggatacetg tecgeettte tecetteggg aagegtggeg ettteteata geteaegetg 5760 taggtatete agtteggtgt aggtegtteg etceaagetg ggetgtgtge acgaacecec 5820 cgttcagccc gaccgctgcg ccttatccgg taactatcgt cttgagtcca acccggtaag 5880 acacgaetta tegecaetgg cageageeae tggtaacagg attagcagag egaggtatgt 5940 aggoggtgot acagagttot tgaagtggtg gootaactac ggotacacta gaaggacagt 6000 atttggtatc tgcgctctgc tgaagccagt taccttcgga aaaagagttg gtagctcttg 6060 atcoggoama camaccaccy ctygtagogg tygttttttt gtttgcmmagc agcagattac 6120 gegeagaaaa aaaggatete aagaagatee titgatetit tetaeggggt etgaegetea 6180 gtggaacqaa aactcacgtt aagggatttt ggtcatgaga ttatcaaaaa ggatcttcac 6240 ctagateett ttaaattaaa aatgaagttt taaateaate taaagtatat atgagtaaae 6300 ttggtctgac agttaccaat gcttaatcag tgaggcacct atctcagcga tctgtctatt 6360 togttoatoc atagttgoot gactocoogt ogtgtagata actaogatac gggagggott 6420 accatctggc cccagtgctg caatgatacc gcgagaccca cgctcaccgg ctccagattt 64'80 atcagcaata aaccagccag ccggaagggc cgagcgcaga agtggtcctg caactttatc 6540 cgcctccatc cagtctatta attgttgccg ggaagctaga gtaagtagtt cgccagttaa 6600 tagtttgcgc aacgttgttg ccattgctgc aggcatcgtg gtgtcacgct cgtcgtttgg 6660 tatggcttca ttcagctccg gttcccaacg atcaaggcga gttacatgat cccccatgtt 6720 gtgcaaaaaa gcggttagct ccttcggtcc tccgatcgtt gtcagaagta agttggccgc 6780 agtgttatca ctcatggtta tggcagcact gcataattct cttactgtca tgccatccgt 6840 aagatgettt tetgtgaetg gtgagtaete aaccaagtea ttetgagaat agtgtatgeg 6900 gcgaccgagt tgctcttgcc cggcgtcaat acgggataat accgcgccac atagcagaac 6960 tttaaaagtg ctcatcattg gaaaacgttc ttcggggcga aaactctcaa ggatcttacc 7020

gctgttgaga	tccagttcga	tgtaacccac	tcgtgcaccc	aactgatctt	cagcatcttt	7080
tactttcacc	agcgtttctg	ggtgagcaaa	aacaggaagg	cadaatgccg	caaaaaaggg	7140
aataagggcg	acacggaaat	gttgaatact	catactcttc	ctttttcaat	attattgaag	7200
catttatcag	ggttattgtc	tcatgagcgg	atacatattt	gaatgtattt	agaaaaataa	7260
acaaataggg	gttccgcgca	catttccccg	aaaagtgcca	cctgacgtct	aagaaaccat	7320
tattatcatg	acattaacct	ataaaaatag	gcgtatcacg	aggccctttc	gtcttcaaga	7380
a						7381

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 29 March 2001 (29.03.2001)

(10) International Publication Number WO 01/21772 A3

C12N 15/52, (51) International Patent Classification7: 15/53, 15/54, 15/60, 15/75, 9/00, 9/02, 9/10, 9/12, 9/88, C12P 7/42, 13/02, 13/06

PCT/US00/25993 (21) International Application Number:

(22) International Filing Date:

21 September 2000 (21.09.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

21 September 1999 (21.09.1999) US 09/400,494 US 60/210,072 7 June 2000 (07.06.2000) US 28 July 2000 (28.07.2000) 60/221,836 24 August 2000 (24.08.2000) 60/227,860

US (71) Applicant: OMNIGENE BIOPRODUCTS [US/US];

(72) Inventors: YOCUM, Rogers, R.; 4 Orchard Lane, Lexington, MA 02420 (US). PATTERSON, Thomas, A.; 89

763(d) Concord Avenue, Cambridge, MA 02138 (US).

Church Street, North Attleboro, MA 02760 (US). HER-MANN, Theron; 18 Chilhowie Drive, Kinnelon, NJ 07405 (US). PERO, Janice, G.; 20 Solomon Pierce Road, Lexington, MA 02420 (US).

- (74) Agents: HANLEY, Elizabeth, A. et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).
- (81) Designated States (national): AU, BR, CA, CN, CZ, HU, ID, IL, IN, JP, KR, LT, LV, MX, NO, PL, RU, SG. SK, TR, UA, ZA.
- (84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).


Published:

with international search report

(88) Date of publication of the international search report: 7 March 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS



(57) Abstract: The present invention features methods of producing panto-compounds (<i>e.g.</i>, pantothenate) using microorganisms in which the pantothenate biosynthetic pathway and/or the isoleucine-valine biosynthetic pathway and/or the coenzymeA biosynthetic pathway has been manipulated. Methods featuring ketopantoate reductase overexpressing microorganisms as well as aspartate g(a)-decarboxylase overexpressing microorganisms are provided. Methods of producing panto-compounds in a precursor-independent manner and in high yield are described. Recombinant microorganisms, vectors, isolated nucleic acid molecules, genes and gene products useful in practicing the above methodologies are also provided. The present invention also features a previously microbial pantothenate kinase gene, <i>coaX</i>, as well as methods of producing panto-compounds utilizing microorganisms having modified pantothenate kinase activity. Recombinant microorganisms, vectors, isolated <i>coaX</i> nucleic acid molecules and purified CoaX proteins are featured. Also featured are methods for identifying pantothenate kinase modulators utilizing the recombinant microorganisms and/or purified CoaX proteins of the present invention.

n: :tional Application No PCT/US 00/25993

4 01 105	TICATION OF CUR LECT MATTER			
A. CLASSIF IPC 7	C12N15/52 C12N15/53 C12N15/5 C12N9/00 C12N9/02 C12N9/10 C12P7/42 C12P13/02 C12P13/0	C12N9/12		
According to	International Patent Classification (IPC) or to both national classification			
B. FIELDS				
Minimum do IPC 7	cumentation searched (classification system followed by classification C12N C12P	symbols)		
	ion searched other than minimum documentation to the extent that su		,	
	ata base consulted during the international search (name of data base		rms used)	
EMBL,	EPO-Internal, WPI Data, PAJ, BIOSIS,	MEDLINE, EMBASE	•	
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT			
Category °	Citation of document, with indication, where appropriate, of the relevant	vant passages	Relevant to claim No.	
х	EP 0 590 857 A (TAKEDA CHEMICAL I LTD) 6 April 1994 (1994-04-06) cited in the application	NDUSTRIES	12,13, 24,26, 27,48, 51,55, 59,60, 71,76	
Y	the whole document		1-6, 33-35, 54, 56-58, 62-64, 78-82	
	page 14, line 1-3	·/		
X Further documents are listed in the continuation of box C. Patent family members are listed in annex.				
*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention lifting date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention lifting date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention cannot be considered to priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention cannot be considered to priority date and not in conflict with the application but cited to understand the principle or theory underlying the cited to understand the princi				
Date of the actual completion of the international search 10 July 2001 2 2. 10, ns				
		Authorized officer		
Name and	Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016 Van de Kamp, M			

Form PCT/ISA/210 (second sheet) (July 1992)

0121772A2 1 -

tnt tional Application No PCT/US 00/25993

0.40	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category *	entry and the relevant passages	Relevant to claim No.
Υ .	SAHM H ET AL.: "D-Pantothenate synthesis in Corynebacterium glutamicum and use of panBC and genes encoding L-valine synthesis for D-pantothenate overproduction" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 65, no. 5, May 1999 (1999-05), pages 1973-1979, XP002169517 cited in the application the whole document	12,13, 24,26, 27,48, 51,54, 56,59, 60,71, 76,78,79
x	SOROKIN A ET AL.: "Sequence analysis of the Bacillus subtilis chromosome region between the serA and kdg loci cloned in a yeast artificial chromosome" MICROBIOLOGY, vol. 142, no. 8, 1996, pages 2005-2016, XP000910121 ISSN: 1350-0872	83-86, 92,93, 95, 97-100, 102,103
Y	page 2011, right-hand column, line 14-20; table 1	1-6, 33-35, 54, 56-58, 62-64, 78-82
X	DATABASE EM_PRO [Online] EMBL; ID BSYPIA, AC L47709, 23 January 1996 (1996-01-23) HENNER D ET AL.: "Bacillus subtilis (clone YAC15-6B) ypiABF genes, qcrABC genes, ypjABCDEFGHI genes, birA gene, panBCD genes, dinG gene, ypmB gene, aspB gene, asnS gene, dnaD gene, nth gene and ypoC gene, complete cds." XP002171539 page 5, line 43-60 page 10	83-86, 98-100, 102,103
A	BAIGORI M ET AL.: "Isolation and characterization of Bacillus subtilis mutants blocked in the syntehsis of pantothenic acid" JOURNAL OF BACTERIOLOGY, vol. 173, no. 13, July 1991 (1991-07), pages 4240-4242, XP001002216 abstract	99,100, 102,103

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

In ational Application No
PCT/US 00/25993

Category °	Citation of document, with indication, where appropriate, of the relevant passages	
Jaley Ury	Chanch of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 18954 A (MONSANTO CO) 7 May 1998 (1998-05-07) the whole document	
Α	EP 0 224 294 A (GIST BROCADES NV) 3 June 1987 (1987-06-03) the whole document	
Ρ,Χ	EP 1 006 192 A (DEGUSSA) 7 June 2000 (2000-06-07)	12,13, 24,26, 27,48, 51, 54-56, 59,60, 71,76, 78,79
	examples 2-5	
P,X	EP 1 006 189 A (DEGUSSA; KERNFORSCHUNGSANLAGE JUELICH (DE)) 7 June 2000 (2000-06-07)	12,13, 24,26, 27,48, 51,54, 56,59, 60,71, 76,78,79
	examples 1,7,9	76,78,79
	·	

4

..ernational application No. PCT/US 00/25993

INTERNATIONAL SEARCH REPORT

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows:
see additional sheet
As all required additional search fees were timely paid by the applicant, this International Search Report-covers all searchable claims.
As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-6,12,13,24,26-28,33-35,48,51,54-64,71,76,78-86,92,93,95,97100,102,103 (all Partially)
Remark on Protest The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1998)

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-6,12,13,24,26-28,33-35,48,51,54-64,71,76,78-86, 92,93,95,97-100,102,103 (all partially)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L $^{\circ}$ at least), pantoate, or ketopantoate, e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses a ketopantoate hydroxymethyltransferase-encoding gene, e.g. the panB gene, e.g., from Bacillus, under conditions such that said panto-compound is produced, and possibly further recovering the compound. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) which overexpresses a Bacillus (subtilis) ketopantoate hydroxymethyltransferase-encoding gene. A recombinant vector encoding a Bacillus (subtilis) ketopantoate hydroxymethyltransferase-encoding gene operably linked to regulatory sequences, e.g., comprising a nucleic acid sequence according to SEQ ID NO:23 or part of SEQ ID NO:59. An isolated nucleic acid molecule encoding a Bacillus (subtilis) ketopantoate hydroxymethyltransferase, and said isolated ketopantoate hydroxymethyltransferase polypeptide.

2. Claims: 1-6,12,13,24,26-28,33-35,48,51,54-64,71,76,78-86, 92,93,95,97-100,102,103 (all partially)

A method of producing panthotenate (e.g., 2 g/L up to 40 g/L at least), pantoate, or ketopantoate, e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses a pantothenate synthetase-encoding gene, e.g. the panC gene. e.g., from Bacillus, under conditions such that said panto-compound is produced, and possibly further recovering the compound. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) which overexpresses a Bacillus (subtilis) pantothenate synthetase-encoding gene. A recombinant vector encoding a Bacillus (subtilis) pantothenate synthetase-encoding gene operably linked to regulatory sequences, e.g., comprising a nucleic acid sequence according to SEQ ID NO:25 or part of SEQ ID NO:59. An isolated nucleic acid molecule encoding a Bacillus (subtilis) pantothenate synthetase, and said isolated pantothenate synthetase polypeptide.

3. Claims: 1-6,12-14,24,26-28,33-35,48,49,51,54-64,66,71,76, 78-86,92,93,95,97-100,102,103 (all partially); 15, 17,19,23,32,106,107 (both completely)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), pantoate, or ketopantoate, e.g., independent of aspartate or beta-alanine feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses an aspartate-alpha-decarboxylase-encoding gene, e.g., from Bacillus, e.g., the aspartate-alpha-decarboxylase-encoding panD gene from Bacillus subtilis, under conditions such that said panto-compound is produced, and possibly further recovering the compound. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) which overexpresses a Bacillus (subtilis) aspartate-alpha-decarboxylase-encoding gene. A recombinant vector encoding a Bacillus (subtilis) aspartate-alpha-decarboxylase-encoding gene operably linked to regulatory sequences, e.g., comprising a nucleic acid sequence according to SEQ ID NO:27 or part of SEQ ID NO:59. An isolated nucleic acid molecule encoding a Bacillus (subtilis) aspartate-alpha-decarboxylase, and said isolated aspartate-alpha-decarboxylase polypeptide.

4. Claims: 1-6,24,26-28,33-35,48,49,51,54-64,71,76,78-87,92, 93,95,97-100,102,103 (all partially); 7-11,65,101, 104,105 (all completely)

> A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), pantoate, or ketopantoate, e.g., independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses a ketopantoate reductase-encoding gene, e.g., from Bacillus, e.g., the ketopantoate reductase-encoding panEl gene from Bacillus subtilis, under conditions such that said panto-compound is produced, and possibly further recovering the compound. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) which overexpresses a Bacillus (subtilis) ketopantoate reductase-encoding gene. A recombinant vector encoding a Bacillus (subtilis) ketopantoate reductase-encoding gene operably linked to regulatory sequences, e.g., comprising a nucleic acid sequence according to SEQ ID NO:29. An isolated nucleic acid molecule encoding a Bacillus (subtilis)

ketopantoate reductase, and said isolated ketopantoate reductase polypeptide.

5. Claims: 14,16,18,28,48,54-61,66,77-82, 97 (all partially); 20,29 (completely)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of valine or alpha-ketoisovalerate feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a deregulated isoleucine-valine (ilv) pathway, wherein the microorganism overexpresses acetohydroxyacid synthase or is transformed with a vector comprising an ilvBN nucleic acid sequence or an alsS sequence, e.g., from Bacillus, under conditions such that pantothenate is produced, and possibly further recovering the pantothenate. Said microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)), and said vector.

6. Claims: 14,16,18,28,48,54-61,66,77-82, 97 (all partially): 21,30 (completely)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of valine or alpha-ketoisovalerate feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococi or Streptomyces) or Gram-negative) having a deregulated isoleucine-valine (ilv) pathway, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase or is transformed with a vector comprising an ilvC nucleic acid sequence, e.g., from Bacillus, under conditions such that pantothenate is produced, and possibly further recovering the pantothenate. Said microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)), and said vector.

7. Claims: 14,16,18,28,48,54-61,66,77-82,97 (partially); 22, 31 (completely)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of valine or alpha-ketoisovalerate feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci

or Streptomyces) or Gram-negative) having a deregulated isoleucine-valine (ilv) pathway, wherein the microorganism overexpresses dihydroxyacid dehydratase or is transformed with a vector comprising an ilvD nucleic acid sequence, e.g., from Bacillus, under conditions such that pantothenate is produced, and possibly further recovering the pantothenate. Said microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)), and said vector.

8. Claims: 25,28,50,54-61 (all partially)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant avtA gene under conditions such that pantothenate is produced, and possibly further recovering the pantothenate.

9. Claims: 25,28,50,54-61 (all partially)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant ilv£ gene under conditions such that pantothenate is produced, and possibly further recovering the pantothenate.

10. Claims: 25,28,50,54-61 (all partially)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant ans8 gene under conditions such that pantothenate is produced, and possibly further recovering the pantothenate.

11. Claims: 25,28,50,54-61 (all partially)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative)

having a mutant alsD gene under conditions such that pantothenate is produced, and possibly further recovering the pantothenate.

12. Claims: 36,37,54-59,61 (all partially)

A method of producing beta-alanine, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses an aspartate-alpha-decarboxylase-encoding gene, under conditions such that beta-alanine is produced, wherein the asparte-alpha-decarboxylase-overexpressing microorganism has a mutation in a nucleic acid sequence encoding ketopantoate hydroxymethyltransferase (panB), and possibly further recovering the compound.

13. Claims: 36,37,54-59,61 (all partially)

A method of producing beta-alanine, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses an aspartate-alpha-decarboxylase-encoding gene, under conditions such that beta-alanine is produced, wherein the asparte-alpha-decarboxylase-overexpressing microorganism has a mutation in a nucleic acid sequence encoding ketopantoate reductase (panE), and possibly further recovering the compound.

14. Claims: 36,37,54-59,61 (all partially)

A method of producing beta-alanine, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses an aspartate-alpha-decarboxylase-encoding gene, under conditions such that beta-alanine is produced, wherein the asparte-alpha-decarboxylase-overexpressing microorganism has a mutation in a nucleic acid sequence encoding pantothenate synthetase (panC), and possibly further recovering the compound.

15. Claim : 38 (completely)

A method of producing beta-alanine comprising contacting a composition comprising aspartate with an isolated Bacillus aspartate-alpha-decarboxylase enzyme under conditions such that beta-alanine is produced.

16. Claims: 41,44-47,51,53,54-61,69,71,72,75,78-81, 97 (all partially); 39,43,52,67,70,74,88-91, 108-110 (all completely)

A method for producing or for enhancing production of ketopantoate, pantoate, or pantothenate (e.g., 10, 20 or 40 g/L at least), comprising culturing a mutant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant pantothenate kinase-encoding coaX gene, under conditions such that said panto-compound is produced or that production is enhanced, and possibly further recovering the compound. A method for identifying compounds which modulate pantothenate kinase activity comprising contacting a recombinant cell expressing the coax gene, possibly further comprising a mutant coaA gene encoding a pantothenate kinase with reduced activity, with a test compound and determining the ability of the test compound to modulate pantothenate kinase activity in said cell. A recombinant microorganism having a mutant coax gene encoding a pantothenate kinase with reduced activity. A vector comprising a mutant coaX gene encoding a pantothenate kinase with reduced activity, possibly further comprising regulatory sequences. A recombinant microorganism comprising a vector comprising an isolated coax gene (e.g., from Bacillus (subtilis)), and said vector, possibly further comprising regulatory sequences, e.g., a constitutively active promoter. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) that overproduces a panto-compound having a mutation in a coaX gene that results in a reduced level of pantothenate kinase activity, resulting in a decrease in the capacity of the microorganism to synthesize coenzyme A. An isolated nucleic acid molecule comprising a (mutant) coax gene, and an isolated pantothenate kinase protein encoded by a coaX gene.

17. Claims: 41,44-47,51,53-61,69,71,72,75,78-81, 97 (all partially); 40,42,68,73 (all completely)

A method for producing or for enhancing production of a panto-compound, e.g., ketopantoate, pantoate or pantothenate (e.g., 10, 20 or 40 g/L at least), comprising culturing a mutant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant pantothenate kinase-encoding coaA gene, under conditions such that the panto-compound is produced or that production is enhanced, and possibly further recovering the panto-compound. A recombinant microorganism having a mutant coaA gene encoding a pantothenate kinase with reduced

activity. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) that overproduces a panto-compound having a mutation in a coaA gene that results in a reduced level of pantothenate kinase activity, resulting in a decrease in the capacity of the microorganism to synthesize coenzyme A. A vector containing a (mutated) coaA gene.

18. Claim: 94 (partially)

A vector containing regulatory sequences comprising the constitutively active promoter Pveg (SEQ ID NO:41).

19. Claim : 94 (partially)

A vector containing regulatory sequences comprising the constitutively active promoter P15 (SEQ ID NO:39).

20. Claim: 94 (partially)

A vector containing regulatory sequences comprising the constitutively active promoter P26 (SEQ ID NO:40).

21. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEO ID NO:49.

22. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:50.

23. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:51.

24. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:52.

25. Claim: 96 (partially)

A vector containing regulatory sequences comprising an

artificial RBS according to SEQ 10 NO:53.

26. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:54.

27. Claim : 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ 1D NO:55.

28. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:56.

29. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:57.

Information on patent family members

Inc. ...tional Application No PCT/US 00/25993

	701/03 00/25993		00/23793
Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0590857	A 06-04-1994	CN 1095105 A,B CN 1225394 A CN 1225388 A CN 1225392 A CN 1225358 A JP 6261772 A US 5518906 A	16-11-1994 11-08-1999 11-08-1999 11-08-1999 11-08-1999 20-09-1994 21-05-1996
WO 9818954	A 07-05-1998	US 5834259 A EP 0934426 A JP 2001503979 T	10-11-1998 11-08-1999 27-03-2001
EP 0224294	A 03-06-1987	AT 56989 T DE 3674548 D DK 533286 A ES 2031818 T FI 864547 A JP 62181789 A PT 83714 A,B US 5024943 A	15-10-1990 31-10-1990 09-05-1987 01-01-1993 09-05-1987 10-08-1987 01-12-1986 18-06-1991
EP 1006192	A 07-06-2000	DE 19855313 A BR 9905776 A CN 1256314 A HU 9904447 A JP 2000228990 A SK 163399 A US 6184007 B	08-06-2000 24-04-2001 14-06-2000 28-11-2000 22-08-2000 11-07-2000 06-02-2001
EP 1006189	A 07-06-2000	DE 19855312 A BR 9905783 A CN 1256313 A HU 9904448 A JP 2000166580 A SK 164099 A US 6177264 B	08-06-2000 24-04-2001 14-06-2000 28-11-2000 20-06-2000 11-07-2000 23-01-2001

Form PCT/ISA/210 (patent tamely annex) (July 1992)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date 29 March 2001 (29.03.2001)

PCT

(10) International Publication Number WO 01/021772 A3

(74) Agents: HANLEY, Elizabeth, A. et al.; Lahive & Cock-

field, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (national): AU, BR, CA, CN, CZ, HU, ID, IL, IN, JP, KR, LT, LV, MX, NO, PL, RU, SG, SK, TR,

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,

- (51) International Patent Classification⁷: C12N 15/52, 15/53, 15/54, 15/60, 15/75, 9/00, 9/02, 9/10, 9/12, 9/88, C12P 7/42, 13/02, 13/06
- (21) International Application Number: PCT/US00/25993
- (22) International Filing Date:

21 September 2000 (21.09.2000)

(25) Filing Language:

English

(26) Publication Language:

English

Published:

UA, ZA.

NL, PT, SE).

- with international search report

(30) Priority Data:

 09/400,494
 21 September 1999 (21.09.1999)
 US

 60/210,072
 7 June 2000 (07.06.2000)
 US

 60/221,836
 28 July 2000 (28.07.2000)
 US

 60/227,860
 24 August 2000 (24.08.2000)
 US

- (71) Applicant: BASF Aktiengesellschaft [DE/DE]; 67056 Ludwigshafen, Rheinland-Pfalz (DE).
- (72) Inventors: YOCUM, Rogers, R.; 4 Orchard Lane, Lexington, MA 02420 (US). PATTERSON, Thomas, A.; 89 Church Street, North Attleboro, MA 02760 (US). HERMANN, Theron; 18 Chilhowie Drive, Kinnelon, NJ 07405 (US). PERO, Janice, G.; 20 Solomon Pierce Road, Lexington, MA 02420 (US).

- (88) Date of publication of the international search report: 7 March 2002
- (48) Date of publication of this corrected version:
 '5 December 2002
 - 5 December 2002
- (15) Information about Correction: see PCT Gazette No. 49/2002 of 5 December 2002, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

7

(54) Title: METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS

(57) Abstract: The present invention features methods of producing panto-compounds e.g., pantothenate) using microorganisms in which the pantothenate biosynthetic pathway and/or the isoleucine-valine biosynthetic pathway and/or the coenzymeA biosynthetic pathway has been manipulated. Methods featuring ketopantoate reductase overexpressing microorganisms as well as aspartate g(a)-decarboxylase overexpressing microorganisms are provided. Methods of producing panto-compounds in a precursor-independent manner and in high yield are described. Recombinant microorganisms, vectors, isolated nucleic acid molecules, genes and gene products useful in practicing the above methodologies are also provided. The present invention also features a previously microbial pantothenate kinase gene, coal, as well as methods of producing panto-compounds utilizing microorganisms having modified pantothenate kinase activity. Recombinant microorganisms, vectors, isolated coal nucleic acid molecules and purified Coal proteins are featured. Also featured are methods for identifying pantothenate kinase modulators utilizing the recombinant microorganisms and/or purified Coal proteins of the present invention.

10

15

20

25

35

METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS

Background of the Invention

Pantothenate, also known as pantothenic acid or vitamin B5, is a member of the B complex of vitamins and is a nutritional requirement for mammals, including livestock and humans (e.g., from food sources, as a water soluble vitamin supplement or as a feed additive). In cells, pantothenate is used primarily for the biosynthesis of coenzyme A (CoA) and acyl carrier protein (ACP). These coenzymes function in the metabolism of acyl moieties which form thioesters with the sulfhydryl group of the 4'-phosphopantetheine portion of these molecules. These coenzymes are essential in all cells, participating in over 100 different intermediary reactions in cellular metabolism.

The conventional means of synthesizing pantothenate (in particular, the bioactive D isomer) is via chemical synthesis from bulk chemicals, a process which is hampered by excessive substrate cost as well as the requirement for optical resolution of racemic intermediates (e.g., resolution of DL-pantolactone to obtain D-pantolactone for chemical condensation with β-alanine). Accordingly, researchers have recently looked to bacterial or microbial systems that produce enzymes useful in pantothenate biosynthesis processes (as bacteria are themselves capable of synthesizing pantothenate). In particular, bioconversion processes have been evaluated as a means of favoring production of the D isomer of pantothenic acid, e.g., using microorganisms which selectively hydrolyze a DL-pantothenic acid ester to D-pantothenic acid; microorganisms which selectively decompose L-pantolactone resulting in D-pantolactone alone; and microorganisms which selectively hydrolyze DL-pantolactone to D-pantoic acid.

There is still, however, significant need for improved pantothenate production processes, in particular, for processes requiring reduced quantities of substrates and/or less expensive substrates. To this end, methods of direct microbial synthesis have recently been examined as a means of improving D-pantothenate production. In microbes, pantothenate biosynthetis is a multistep pathway resulting in condensation of pantoate (derived from α -ketoisovalerate) and β -alanine to form D-pantothenate. The isoleucine-valine (ilv) pathway biosynthetic enzymes, acetohydroxyacid synthetase (the ilvBN or alsS gene product), acetohydroxyacid isomeroreductase (the ilvC gene product) and dihydroxyacid dehydratase (the ilvD gene product) catalyze the conversion of pyruvate to α -ketoisovalerate. The reactions are further catalyzed by the pantothenate (pan) pathway biosynthetic enzymes ketopantoate hydroxymethyltransferase (the panB gene product), ketopantoate reductase (the panE gene product), aspartate- α -

15

20

25

decarboxylase (the panD gene product) and pantothenate synthetase (the panC gene product).

The genes encoding the enzymes involved in the biosynthesis of pantothenic acid in Salmonella typhimurium and Escherichia coli have recently been identified and characterized (Frodyma and Downs (1998) J. Biol. Chem. 273:5572-5576 and Jackowski (1996) pp. 687-694, In Neidhardt et al (ed.) Escherichia coli and Salmonella: Cellular and Molecular Biology, 2^{nd} ed. Am. Soc. Microbiol. Wash, D.C). In E. coli, for example, the biosynthesis of pantothenic acid consists of four key steps. The first reaction is catalyzed by the panB gene product, ketopantoate hydroxymethyltransferase, and uses the L-valine intermediate α -ketoisovalerate to generate ketopantoate, which is subsequently reduced to pantoate by the panE gene product, ketopantoate reductase. The panD gene product, aspartate- α -decarboxylase, generates β -alanine from aspartate. The panC gene product, pantothenate synthetase, subsequently ligates β -alanine with pantoate to yield D-pantothenate.

The authors Dusch et al. described the identification of the Corynebacterium glutamicum panD gene and reported that expression of the C. glutamicum panD gene in E. coli yielded a strain producing pantothenate with a specific productivity of 140 ng of pantothenate per mg (dry weight) per hour. (Dusch et al. (1999) Appl. Environ. Microbiol. 65:1530-1539).

The authors Sahm and Eggeling have further identified the Corynebacterium glutamicum panB and pan C genes and have described a genetically engineered strain of C. glutamicum which overexpresses the panBC genes (Sahm and Eggeling (1999) Appl. Environ. Microbiol. 65:1973-1979). The engineered strain produces pantothenate, however, it was necessary to overexpress the genes responsible for α -ketoisovalerate production in the host organism in order that pantothenic acid production could be detected. Moreover, without the addition of β -alanine, no substantial amounts of pantothenate accumulated with the strain constructed.

Likewise, a method of producing D-pantothenic acid has been described that takes advantage of a sodium salicylate resistant mutant strain of E. coli which produces D-pantothenic acid when cultured in the presence of β -alanine (U.S. Patent No. 5,518,906). Generation of E. coli strains resistant to α -ketoisovaleric acid and/or α -ketobutyric acid, and/or α -aminobutyric acid, and/or β -hydroxyaspartic acid and/or O-methyl-threonine, in addition to salicylic acid, further increased pantothenic acid production. Moreover, transformation of a plasmid DNA carrying the panB, panC and panD genes into the salicylic acid resistant mutant strain resulted in increased pantothenate production, however, up to 20 g/L β -alanine or more was fed in the examples given. The panB-panC-panD genes are clustered on the E. coli chromosome.

20

Finally, a method of producing D-pantothenic acid has been described which utilizes a salicylic acid-resistant, α -ketoisovalerate-resistant, α -ketobutyrate-resistant, β hydroxyaspartate-resistant, o-methylthreonine-resistent E. coli strain transformed with pantothenate biosynthesis gene-containing DNA fragments and/or branched amino acid biosynthesis gene-containing DNA fragments and cultured in the presence of β-alanine (U.S. Patent No. 5,932,457).

Pantothenate production in bacteria results from the condensation of pantoate and β-alanine and involves the pantothenate biosynthetic enzymes ketopantoate hydroxymethyltransferase (the panB gene product), ketopantoate reductase (the panEgene product), aspartate- α -decarboxylase (the panD gene product) and pantothenate synthetase (the panC gene product). Although pantothenate is biologically active as a vitamin, it is further metabolized in all cells to Coenzyme A (CoA) which participates as an acyl group carrier in the tricarboxylic acid (TCA) cycle, fatty acid metabolism and numerous other reactions of intermediary metabolism. The initial (and possibly rate-15. controlling) step in the conversion of pantothenate to Coenzyme A (CoA) is phosphorylation of pantothenate by pantothenate kinase. A pantothenate kinase activity was first identified in Salmonella typhimurium by screening for temperature-sensitive mutants which synthesized CoA at permissive temperatures but excreted pantothenate at non-permissive temperatures. The mutations were mapped in the Salmonella chromosome and the genetic locus was designated coaA. The gene encodes the enzyme that catalyzes the first step in the biosynthesis of coenzyme A from pantothenate (Dunn and Snell (1979) J. Bacteriol. 140:805-808). Escherichia coli temperature sensitive mutants have also been isolated and characterized (Vallari and Rock (1987) J. Bacteriol. 169:5795-5800). These mutants (named coaA15(Ts)) are defective in the conversion of pantothenate to CoA and further exhibit a temperature-sensitive growth phenotype, indicating that pantothenate kinase activity is essential for growth. Moreover, it was noted that CoA inhibited pantothenate kinase activity to the same degree in the mutant as compared to the wild-type enzyme.

Feedback resistant E. coli mutants (named coaA16(Fr)) have also been isolated that posses a pantothenate kinase activity that is refractory to feedback inhibition by 30 CoA (Vallari and Jackowski (1988) J. Bacteriol. 170:3961-3966). The mutation responsible for the reversion is, suprisingly, not genetically linked to the coal gene by transduction. Additional data described therein support the view that the total cellular CoA content is controlled by both modulation of biosynthesis at the pantothenate kinase step and possibly by degradation of CoA to 4'-phosphopantetheine. 35

20

25

30

35

The wild-type E. coli coaA gene was cloned by functional complementation of E. coli temperature-sensitive mutants. The sequence of the wild-type gene was determined (Song and Jackowski (1992) J. Bacteriol. 174:6411-6417 and Flamm et al. (1988) Gene (Amst.) 74:555-558). Strains containing multiple copies of the coaA gene possessed 76-fold higher specific activity of pantothenate kinase, however, there was only a 2.7-fold increase in the steady state level of CoA (Song and Jackowski, supra). It has further been reported that the prokaryotic enzyme (encoded by coaA in E.coli and a variety of other microorganisms) is feedback inhibited by CoA both in vivo and in vitro with CoA being about five times more potent than acetyl-CoA in inhibiting the enzyme (Song and Jackowski, supra and Vallari et al., supra). Moreover, it has been reported that the panB gene product in E. coli is inhibited by CoA (Powers and Snell (1976) J. Biol. Chem. 251:3786-3793). These data further support the view that feedback inhibition of pantothenate kinase activity is a critical factor controlling intracellular CoA concentration.

Using standard search and alignment tools, coal homologues have been identified in Hemophilus influenzae, Mycobacterium tuberculosis, Vibrio cholerae, Streptococcus pyogenes and Bacillus subtilis. By contrast, proteins with significant similarity could not be identified in eukaryotic cells including Saccharomyces cerevisiae or in mammalian expressed sequence tag (EST) databases. Using a genetic selection strategy, a cDNA encoding pantothenate kinase activity has recently been identified from Aspergillus nidulans (Calder et al. (1999) J. Biol. Chem. 274:2014-2020). The eukaryotic pantothenate kinase gene (panK) has distinct primary structure and unique regulatory properties that clearly distinguish it from its prokaryotic counterpart. A mammalian pantothenate kinase gene (mpanKla) has also been isolated which encodes a protein having homology to the A. nidulans PanK protein and to the predicted gene product of GenBankTM Accession Number 927798 identified in the S. cerevisiae genome (Rock et al. (2000) J. Biol. Chem. 275:1377-1383).

Summary of the Invention

The present invention is based, at least in part, on the discovery of key enzyme-encoding genes of the pantothenate biosynthetic pathway in Bacillus subtilis. In particular, the present inventors have identified the panE gene of B. subtilis. Overexpression or deregulation of the panE gene in B. subtilis results in enhanced production of the panE gene product, ketopantoate reductase, further resulting in increased production of pantothenate. Likewise, mutations in this gene reduce pantothenate production in B. subtilis >90%. The present inventors have further identified the presumptive panBCD operon in B. subtilis, overexpression or

15.

20

25

30

35

deregulation of which results in increased pantothenate production. The present inventors have further demonstrated that overexpression or deregulation of the panD gene in B. subtilis (resulting in enhanced production of the panD gene product, aspartate- α -decarboxylase) further results in increased production of pantothenate, in particular, in combination with deregulation of genes encoding key enzymes of the isoleucine-valine (ilv) biosynthetic pathway.

Accordingly, the present invention features methods of producing pantothenate, as well as other compounds of the pantothenate biosynthetic pathway (e.g., ketopantoate, pantoate and β-alanine), termed "panto-compounds" herein, using microorganisms in which the pantothenate biosynthetic pathway and/or isoleucinevaline biosynthetic pathway has been manipulated such that pantotheriate or other desired panto-compounds are produced. In one embodiment, the invention features a method of producing a panto-compound (e.g., pantothenate or pantoate) that involves culturing a microorganism which overexpresses the panE gene product, ketopantoate reductase, also referred to herein as a ketopantoate reductase-overexpressing or "KPAR-O" microorganism, under conditions such that the panto-compound (e.g., pantothenate or pantoate) is produced. In another embodiment, the present invention features a method of producing panto-compounds (e.g., pantothenate or pantoate) which includes culturing a microorganism which overexpresses at least one pantothenate biosynthetic enzyme (e.g., at least one of the panB, panC or panD gene products), preferably in a KPAR-O microorganism, under conditions such that the panto-compound (e.g., pantothenate or pantoate) is produced.

Yet another aspect of the invention features methods of producing panto-compounds which are independent of the need to feed precursors (e.g., β -alanine or aspartate and/or α -ketoisovalerate or valine). In one embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed that includes culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism having a deregulated isoleucine-valine (iiv) pathway under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed that includes culturing an A α D-O microorganism having a deregulated pantothenate (pan) pathway and a deregulated isoleucine-valine (iiv) pathway, under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of aspartate or β -alanine feed that includes culturing an A α D-O microorganism under conditions such that pantothenate is produced. In another embodiment, the invention such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of valine or α -ketoisovalerate feed that includes

15

20

25

30

35

culturing a microorganism having a deregulated isoleucine-valine (*ilv*) biosynthetic pathway under conditions such that pantothenate is produced. In yet another embodiment, the invention features a high yield production method for producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly high yield (*e.g.*, at a level greater than 10 g/L, 20 g/L, 30 g/L or 40g/L).

The methods of the present invention further feature microorganisms that overexpresses acetohydroxyacid synthetase or acetohydroxyacid isomeroreductase (e.g., microorganisms transformed with a vector that includes an ilvBNC nucleic acid sequence), microorganisms that overexpresses dihydroxyacid dehydratase (e.g., microorganisms transformed with a vector that includes an ilvD nucleic acid sequence), microorganisms that overexpresses aspartate-α-decarboxylase (e.g., microorganisms transformed with a vector that includes a panD nucleic acid sequence), microorganisms having a deregulated isoleucine-valine (ilv) biosynthetic pathway and microorganisms having a deregulated pantothenate biosynthetic pathway (e.g., microorganisms that overexpress any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate-a-decarboxylase, for example, microorganisms transformed with a vector comprising a panBCD nucleic acid sequence or a vector comprising a panE1 nucleic acid sequence). In one embodiment, the recombinant microorganism is Gram positive (e.g., microorganisms belonging to the genus Bacillus. Cornyebacterium, Lactobacillus, Lactococci or Streptomyces). In another embodiment, the recombinant microorganism is Gram negative. Particularly preferred is a Bacillus recombinant microorganism (e.g., Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus subtilis, Bacillus pumilus, Bacillus halodurans, and the like). Recombinant vectors that contain the genes encoding Bacillus pantothenate and/or isoleucine-valine biosynthetic enzymes (e.g., B. subtilis pantothenate and/or isoleucine-valine biosynthetic enzymes) are also described.

Also featured are methods of producing β -alanine that include culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that β -alanine is produced and methods of producing β -alanine that involve contacting a composition comprising aspartate with an isolated *Bacillus* aspartate- α -decarboxylase enzyme under conditions such that β -alanine is produced.

The production methods of the present invention further can include recovering the panto-compound (e.g., pantothenate or pantoate).

The present invention further features recombinant microorganisms (e.g., AaD-O microorganisms, microorganisms having a deregulated isoleucine-valine (ilv) pathway, microorganisms overexpressing at least one of ketopantoate

10

20

25

30

35

hydroxymethyltransferase (the panB gene product), pantothenate synthetase (the panC gene product), aspartate-a-decarboxylase (the panD gene product), ketopantoate reductase (the panE1 gene product) and microorganisms having a deregulated panBCD operon. Also featured are panB, panC, panD, panE, ilvB, ilvN, alsS, ilvC, and/or ilvD nucleic acid molecules, as well as vectors including such nucleic acid molecules and gene products encoded by such nucleic acid molecules.

The methodology of the present invention further includes, for example in addition to overexpressing at least one pantothenate biosynthetic enzyme, deleting or mutating a second pantothenate biosynthetic enzyme, said second pantothenate biosynthetic enzyme preferably being downstream of the desired product in the pantothenate biosynthetic pathway. For example, mutating panC, in addition to overexpressing the panE gene product, results in even further enhanced or increased production of pantoate. Accordingly, in one embodiment, the invention features a method of producing pantoate which includes culturing a microorganism which 15 overexpresses the panE gene product and which has a deletion in the panC gene. In another embodiment, the invention features a method of producing pantoate which includes culturing a microorganism which overexpresses the panE gene product and/or panB gene product and which has a deletion in the panC gene. Other exemplary embodiments include a method of producing ketopantoate which includes culturing a microorganism which overexpresses the panB gene product and which has a deletion in the panE gene and a method of producing \beta-alanine which includes culturing a microorganism which overexpresses the panD gene product and which has a deletion in the panC gene. Also included are methods of producing panto-compounds which include overexpressing at least one valine biosynthetic enzyme in a microorganism which has at least one pantothenate biosynthetic enzyme deleted.

The present invention is also based at least in part, on the identification and characterization of a previously unidentified microbial pantothenate kinase gene, coaX. CoaX was first identified in Bacillus subtilis and corresponds to an open reading frame in a portion of the chromosomal DNA that includes the 5' end of the ftsH gene, and all of the yacB, yacC, yacD, cysK and pabB genes. The present inventors have demonstrated that the yacB open reading frame encodes a novel pantothenate kinase activity, the gene being unrelated by homology to any previously known pantothenate kinase gene. The gene has been renamed coaX, as it encodes the enzyme which catalyzes the first step in the pathway from pantothenate to CoaA.

Accordingly, the present invention features new and improved methods of producing pantothenate and other key compounds of the pantothenate biosynthetic pathway (e.g., panto-compounds) utilizing microorganisms having modified

10

15

20

25

30

pantothenate kinase activity. In particular, the present invention features recombinant microorganisms that contain the coaX gene or that contain a mutant coaX gene, having reduced pantothenate kinase activity. In one embodiment, the invention features such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway. In another embodiment, the invention features such recombinant microorganisms further having a deregulated isoleucine-valine (ilv) pathway. In a preferred embodiment, the microorganisms belong to the genus Bacillus (e.g., B. subtilis).

The present invention also features recombinant microorganisms (e.g., microorganisms belonging to the genus Bacillus, for example, B. subtilis) that contain the coaA gene or that contain a mutant coaA gene, optionally including a coaX gene or mutant thereof, having reduced pantothenate kinase activity. In one embodiment, the invention features such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway or having a deregulated isoleucine-valine (ilv) pathway.

Also featured are vectors that contain isolated coaX or coaA genes as well as mutant coaX and/or coaA genes. Isolated nucleic acid molecules that contain isolated coaX genes or mutant coaX genes are featured in addition to isolated CoaX proteins and mutant CoaX proteins.

The nucleic acids, vectors and recombinant microorganisms described above are particularly useful in the methodologies of the present invention. In particular, the invention features methods of enhancing panto-compound production (e.g., ketopantoate, pantoate and or pantothenate production) that include culturing a recombinant microorganism having a mutant coaX gene under conditions such that panto-compound production is enhanced. In one embodiment, the recombinant microorganism further includes a mutant coaA gene. In another embodiment, the recombinant microorganism further includes a mutant avtA and/or mutant ilvE gene and/or mutant ansB gene and/or mutant alsD gene. Also featured are methods for identifying pantothenate modulators utilizing the recombinant microorganisms and purified CoaX proteins of the present invention.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1 is a schematic representation of the pantothenate biosynthetic pathway. Figure 2 is a schematic representation of the plasmid pAN240, containing sequences ligated upstream of the $P_{26}panBCD$ cassette, equivalent to the integrated

5 'version in strain PA221.

Figure 3A is a schematic representation of the plasmid pAN004, containing the panBCD operon expressed from P_{26} and RBS1.

Figure 3B is a schematic representation of the plasmid pAN006, containing the panBCD operon expressed from P_{26} and RBS2.

Figure 4 is a schematic representation of the plasmid pAN236, containing an integratable and amplifiable P_{26} -RBS2-panE1 expression cassette.

Figure 5 is a schematic representation of the construction of plasmid pAN423.

Figure 6 is a schematic representation of the construction of plasmids pAN426 and pAN427.

Figure 7 is a schematic representation of the construction of plasmids pAN428 and pAN429.

Figure 8 is a schematic representation of the construction of plasmid pAN431.

Figure 9 is a schematic representation of the construction of plasmid pAN441.

Figure 10 is a schematic representation of the construction of plasmid pAN440.

Figure 11 is a schematic representation of the plasmid pAN251 designed to integrate a single copy of a P_{26} -panE1 cassette at the panE1 locus by double crossover.

Figure 12 is a schematic representation of the plasmid pAN267 designed to integrate a single copy of a P_{26} -ilvBNC cassette at the amyE locus.

Figure 13 is a schematic representation of the plasmid pAN257, a clone of Bacillus subtilis ilvD in a low copy vector.

Figure 14 is a schematic representation of the plasmid pAN263, designed to integrate a single copy of a P_{26} -ilvD cassette at the ilvD locus.

Figure 15 is a schematic representation of the plasmid pAN261, designed to disrupt the Bacillus subtilis ilvD gene with the cat gene.

Figure 16 is a schematic representation of the Coenzyme A biosynthetic pathway in E. coli.

Figure 17 is a schematic representation of the structure of pAN296, a plasmid designed to delete most of the B. subtilis coaA gene and substitute a chloramphenicol resistance gene.

Figure 18 is a schematic representation of the structure of the Bacillus subtilis genome in the region of the coaA gene. The scale is in base pairs and the significant open reading frames are shown by open arrows.

20

Figure 19 is a schematic representation of the plasmid pAN281, a plasmid for expressing Bacillus subtilis coaA after integration at the bpr locus.

Figure 20A-B depicts a multiple sequence alignment (MSA) of the amino acid sequences encoded by six known or predicted microbial coaA genes. SEQ ID NOs:4-6 and 1-3 correspond to the amino acid sequences of Mycobacterium leprae (SwissProtTM Accession No. Q9X795), Mycobacterium tuberculosis (SwissProtTM Accession No. O53440), Streptomyces coelicolor (SwissProtTM Accession No. O86799), Haemophilus influenzae (SwissProtTM Accession No. P44793), Escherichia coli SwissProtTM Accession No. P15044) and Bacillus subtilis (SwissProtTM Accession No. P54556), respectively. The alignment was generated using ClustalW MSA software at the GenomeNet CLUSTALW Server at the Institute for Chemical Research, Kyoto University. The following parameters were used: Pairwise Alignment, K-tuple (word) size = 1, Window size = 5, Gap Penalty = 3, Number of Top Diagonals = 5, Scoring Method = Percent; Multiple Alignment, Gap Open Penalty = 10, Gap Extension Penalty = 0.0, Weight Transition = No, Hydrophilic residues = Gly, Pro, Ser, Asn, Asp, Gln, Glu, Arg and Lys, Hydrophobic Gaps = Yes; and Scoring Matrix = BLOSUM.

Figure 21 is a schematic representation of the structure of the Bacillus subtilis genome in the region of the coaX (yacB) gene. The scale is in base pairs, the significant open reading frames are shown by open arrows and certain predicted restriction fragments are indicated by thick bars.

Figure 22 is a schematic representation of the structure of pAN341 and pAN342, two independent PCR-derived clones of B. subtilis yacB (remaned herein as coaX).

Figure 23A-D depicts a multiple sequence alignment (MSA) of the amino acid sequences encoded by fourteen known or predicted microbial coax genes. SEQ ID NOs:9, 74, 7-8, 75, 11, 10 and 12-18 correspond to the amino acid sequences of Bacillus 25 subtilis (SwissProt™ Accession No. P37564), Clostridium acetobulyticum (WIT™ Accession No. RCA03301, Argonne National Laboratories), Streptomyces coelicolor (PIR™ Accession No. T36391), Mycobacterium tuberculosis (SwissProt™ Accession No. O06282), Rhodobacter capsulatus (WIT™ Accession No. RRC02473), Desulfovibrio vulgaris (DBJTM Accession No. BAA21476.1), Deinococcus radiodurans 30 (SwissProt™ Accession No. Q9RX54), Thermotoga maritima (GenBank™ Accession No. AAD35964.1), Treponema pallidum (SwissProt™ Accession No. O83446). Borrelia burgdorferi (SwissProt™ Accession No.O51477), Aquifex aeolicus (SwissProt™ Accession No. O67753), Synechocystis sp. (SwissProt™ Accession No. P74045), Helicobacter pylori (SwissProt™ Accession No. O25533), and Bordetella 35

pertussis (SwissProt[™] Accession No. Q45338), respectively. The alignment was

generated using ClustalW MSA software at the GenomeNet CLUSTALW Server at the

20

Institute for Chemical Research, Kyoto University. The following parameters were used: Pairwise Alignment, K-tuple (word) size = 1, Window size = 5, Gap Penalty = 3, Number of Top Diagonals = 5, Scoring Method = Percent; Multiple Alignment, Gap Open Penalty = 10, Gap Extension Penalty = 0.0, Weight Transition = No, Hydrophilic residues = Gly, Pro, Ser, Asn, Asp, Gln, Glu, Arg and Lys, Hydrophobic Gaps = Yes; and Scoring Matrix = BLOSUM.

Figure 24 depicts a multiple sequence alignment of a portion of the protein sequences of the coaA gene products from the following microorganisms: Bacillus subtilis, Escherichia coli, Haemophilus influenzae, Mycobacterium leprae, Mycobacterium tuberculosis, and Streptomyces coelicolor. The residues that are mutated in E. coli coaA15(Ts) and B. subtilis coaA282A are indicated below and above the alignment, respectively. The portions correspond to amino acid residues 168-187 of SEQ ID NO:3, 167-186 of SEQ ID NO:2, 165-184 of SEQ ID NO:1, 169-188 of SEQ ID NO:4, 169-188 of SEQ ID NO:5 and 179-198 of SEQ ID NO:6, respectively.

Figure 25 is a schematic representation of the structure of pAN294, a plasmid for integrating mutagenized B. subtilis coaA at its native locus.

Figure 26 is a schematic representation of the structure of pAN336, a plasmid designed to delete B. subtilis coaX from its chromosomal locus and replace it with a kanamycin resistence gene.

20

30

35

15

Detailed Description of the Invention

The present invention features new and improved methods of producing pantothenate and other key compounds of the pantothenate biosynthetic pathway (referred to herein as "panto-compounds", for example, pantothenate, ketopantoate, pantoate and β -alanine) using microorganisms in which the pantothenate biosynthetic pathway has been manipulated such that pantothenate or other desired panto-compounds are produced.

The new and improved methodologies of the present invention include methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one enzyme of the pantothenate biosynthetic pathway manipulated such that pantothenate or other desired panto-compounds are produced (e.g., produced at an increased level). For example, the invention features methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase or aspartate- α -decarboxylase manipulated such that pantothenate or other desired panto-compounds are produced. The methodologies of the present invention also include methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one

valine-isoleucine biosynthetic enzyme, described herein, manipulated such that pantothenate or other desired panto-compounds are produced. For example, the invention features methods of producing panto-compounds (e.g., pantothenate) in microorganisms having at least one of acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase or dihydroxyacid dehydratase manipulated such that pantothenate or other desired panto-compounds are produced.

The invention also features methods of producing panto-compounds that involve culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism under conditions such that the panto-compound is produced. The invention also features methods of producing pantothenate in a manner independent of precursor feed that involve culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism under conditions such that pantothenate is produced. Also featured are β-alanine independent high yield pantothenate production methods as well as methods of producing β-alanine. The present invention also features methods for enhancing production of panto-compounds that involve culturing pantothenate kinase mutants. In particular, the present invention features new and improved methods of producing pantothenate and other key compounds of the pantothenate biosynthetic pathway (e.g., panto-compounds) utilizing microorganisms having modified pantothenate kinase activity, for example, microorganisms that include the coaX gene or that include a mutant coaX gene, having reduced pantothenate kinase activity.

In order that the present invention may be more readily understood, certain terms are first defined herein.

The term "pantothenate biosynthetic pathway" includes the biosynthetic pathway involving pantothenate biosynthetic enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of pantothenate. The term "pantothenate biosynthetic pathway" includes the biosynthetic pathway leading to the synthesis of pantothenate in a microorganisms (e.g., in vivo) as well as the biosynthetic pathway leading to the synthesis of pantothenate in vitro. Figure 1 includes a schematic representation of the pantothenate biosynthetic pathway. Pantothenate biosynthetic enzymes are depicted in bold and their corresponding genes indicated in italics.

The term "pantothenate biosynthetic enzyme" includes any enzyme utilized in the formation of a compound (e.g., intermediate or product) of the pantothenate biosynthetic pathway. According to Figure 1, synthesis of pantoate from α -ketoisovalerate (α -KIV) proceeds via the intermediate, ketopantoate. Formation of ketopantoate is catalyzed by the pantothenate biosynthetic enzyme ketopantoate

20

25

30

35

15

20

30

hydroxymethyltransferase (the panB gene product). Formation of pantoate is catalyzed by the pantothenate biosynthetic enzyme ketopantoate reductase (the panE gene product). Synthesis of β -alanine from aspartate is catalyzed by the pantothenate biosynthetic enzyme aspartate- α -decarboxylase (the panD gene product). Formation of pantothenate from pantoate and β -alanine (e.g., condensation) is catalyzed by the pantothenate biosynthetic enzyme pantothenate synthetase (the panC gene product).

The term "isoleucine-valine biosynthetic pathway" includes the biosynthetic pathway involving isoleucine-valine biosynthetic enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of conversion of pyruvate to valine or isoleucine. The term "isoleucine-valine biosynthetic pathway" includes the biosynthetic pathway leading to the synthesis of valine or isoleucine in a microorganisms (e.g., in vivo) as well as the biosynthetic pathway leading to the synthesis of valine or isoleucine in vitro. Figure 1 includes a schematic representation of the isoleucine-valine biosynthetic pathway. Isoleucine-valine biosynthetic enzymes are depicted in bold italics and their corresponding genes indicated in italics

The term "isoleucine-valine biosynthetic enzyme" includes any enzyme utilized in the formation of a compound (e.g., intermediate or product) of the isoleucine-valine biosynthetic pathway. According to Figure 1, synthesis of valine from pyruvate proceeds via the intermediates, acetolactate, α,β -dihydroxyisovalerate (α,β -DHIV) and α -ketoisovalerate (α -KIV). Formation of acetolactate from pyruvate is catalyzed by the isoleucine-valine biosynthetic enzyme acetohydroxyacid synthetase (the ilvBN gene product, or alternatively, the alsS gene product). Formation of α,β -DHIV from acetolactate is catalyzed by the isoleucine-valine biosynthetic enzyme acetohydroxyacidisomero reductase (the ilvC gene product). Synthesis of α -KIV from α,β -DHIV is catalyzed by the isoleucine-valine biosynthetic enzyme dihydroxyacid dehydratase (the ilvD gene product). Moreover, valine and isoleucine can be interconverted by branched chain amino acid transaminases.

As used herein, each of ketopantoate, pantoate, β -alanine and pantothenate are "panto-compounds". The term "panto-compound" includes a compound (e.g., a substrate, intermediate or product) in the pantothenate biosynthetic pathway which is downstream from a particular pantothenate biosynthetic enzyme. In one example, a panto-compound is downstream of the pantothenate biosynthetic enzyme ketopantoate hydroxymethyltransferase (the panB gene product) and can include ketopantoate, pantoate and/or pantothenate. In another example, a panto-compound is downstream of the pantothenate biosynthetic enzyme ketopantoate reductase (the panE gene product)

20

30

and can include pantoate and/or pantothenate. In yet another example, a panto-compound is downstream of the pantothenate biosynthetic enzyme pantothenate synthetase (the panC gene product) and can include pantothenate. In yet another example, a panto-compound is downstream of the pantothenate biosynthetic enzyme aspartate- α -decarboxylase (the panD gene product) and can include β -alanine and/or pantothenate.

Preferred panto-compounds include pantothenate and pantoate. The term "pantothenate" includes the free acid form of pantothenate, also referred to as "pantothenic acid" as well as any salt thereof (e.g., derived by replacing the acidic hydrogen of pantothenate or pantothenic acid with a cation, for example, calcium, sodium, potassium, ammonium), also referred to as a "pantothenate salt". The term "panto-compound" also includes alcohol derivatives of pantothenate. Preferred pantothenate salts are calcium pantothenate or sodium pantothenate. A preferred alcohol derivative is pantothenol. Pantothenate salts and/or alcohols of the present invention include salts and/or alcohols prepared via conventional methods from the free acids described herein. In another embodiment, calcium pantothenate is synthesized directly by a microorganism of the present invention. A pantothenate salt of the present invention can likewise be converted to a free acid form of pantothenate or pantothenic acid by conventional methodology.

The term "pantoate" includes the free acid form of pantoate, also referred to as "pantoic acid" as well as any salt thereof (e.g., derived by replacing the acidic hydrogen of pantoate or pantoic acid with a cation, for example, calcium, sodium, potassium, ammonium), also referred to as a "pantoate salt". Preferred pantoate salts are calcium pantoate or sodium pantoate. Pantoate salts of the present invention include salts prepared via conventional methods from the free acids described herein. A pantoate salt of the present invention can likewise be converted to a free acid form of pantoate or pantoic acid by conventional methodology. Moreover, a free acid form of pantoate or pantoic acid can be converted to pantolactone by conventional methodology.

The term "CoA biosynthetic pathway" includes the biosynthetic pathway involving CoA biosynthetic enzymes (e.g., polypeptides encoded by biosynthetic enzyme-encoding genes), compounds (e.g., precursors, substrates, intermediates or products), cofactors and the like utilized in the formation or synthesis of CoA from pantothenate. A schematic representation of the CoA biosynthetic pathway in E. coli is set forth as Figure 16. (The pathway depicted is also presumed to be that utilized by other microorganisms.) The term "CoA biosynthetic pathway" includes the biosynthetic pathway leading to the synthesis of CoA in microorganisms (e.g., in vivo) as well as the biosynthetic pathway leading to the synthesis of CoA in vitro. The term "Coenzyme A

or CoA biosynthetic enzyme" includes any enzyme utilized in the formation of a compound (e.g., intermediate or product) of the CoA biosynthetic pathway, for example, the coaA, panK or coaX gene product which catalyzes the phosphorylation of pantothenate to form 4'-phosphopantothenate, or the coaD gene product which catalyzes the conversion of 4'-phosphopantetheine to dephosphocoenzyme A.

I. Recombinant Microorganisms and Methods for Culturing Microorganisms Such That Panto-Compounds are Produced

The methodologies of the present invention feature microorganisms, e.g., recombinant microorganisms, preferably including vectors or genes (e.g., wild-type 10 and/or mutated genes) as described herein and/or cultured in a manner which results in the production of a desired product (e.g. a panto-compound or panto-compounds). The term "recombinant" microorganism includes a microorganism (e.g., bacteria, yeast cell, fungal cell, etc.) which has been genetically altered, modified or engineered (e.g., genetically engineered) such that it exhibits an altered, modified or different genotype 15 and/or phenotype (e.g., when the genetic modification affects coding nucleic acid sequences of the microorganism) as compared to the naturally-occurring microorganism. from which it was derived. Preferably, a "recombinant" microorganism of the present invention has been genetically engineered such that it overexpresses at least one bacterial gene or gene product (e.g., a pantothenate or isoleucine-valine biosynthetic 20 enzyme encoding-gene) as described herein, preferably a biosynthetic enzyme encodinggene included within a recombinant vector as described herein and/or a biosynthetic enzyme expressed from a recombinant vector. The ordinary skilled will appreciate that a microorganism expressing or overexpressing a gene product produces or overproduces the gene product as a result of expression or overexpression of nucleic acid sequences 25 and/or genes encoding the gene product.

The term "manipulated microorganism" includes a microorganism that has been engineered (e.g., genetically engineered) or modified such that the microorganism has at least one enzyme of the pantothenate biosynthetic pathway and/or at least one enzyme of the isoleucine-valine biosynthetic pathway modified such that pantothenate or other desired panto-compounds are produced. Modification or engineering of such microorganisms can be according to any methodology described herein including, but not limited to, deregulation of a biosynthetic pathway and/or overexpression of at least one biosynthetic enzyme. A "manipulated" enzyme (e.g., a "manipulated" biosynthetic enzyme) includes an enzyme, the expression or production of which has been altered or modified such that at least one upstream or downstream precursor, substrate or product

30

35

of the enzyme is altered or modified, for example, as compared to a corresponding wildtype or naturally occurring enzyme.

The term "overexpressed" or "overexpression" includes expression of a gene product (e.g., a pantothenate biosynthetic enzyme or isoleucine-valine biosynthetic enzyme) at a level greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. In one embodiment, the microorganism can be genetically manipulated (e.g., genetically engineered) to overexpress a level of gene product greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. Genetic manipulation can include, but is not limited to, altering or modifying regulatory sequences or sites associated with expression of a particular gene (e.g., by adding strong promoters, inducible promoters or multiple promoters or by removing regulatory sequences such that expression is constitutive), modifying the chromosomal location of a particular gene, altering nucleic acid sequences adjacent to a particular gene such as a ribosome binding site or transcription terminator, increasing the copy number of a particular gene, modifying proteins (e.g., regulatory proteins, suppressors, enhancers, transcriptional activators and the like) involved in transcription of a particular gene and/or translation of a particular gene product, or any other conventional means of deregulating expression of a particular-gene routine in the art (including but not limited to use of antisense nucleic acid molecules, for example, to block expression of repressor proteins).

In another embodiment, the microorganism can be physically or environmentally manipulated to overexpress a level of gene product greater than that expressed prior to manipulation of the microorganism or in a comparable microorganism which has not been manipulated. For example, a microorganism can be treated with or cultured in the presence of an agent known or suspected to increase transcription of a particular gene and/or translation of a particular gene product such that transcription and/or translation are enhanced or increased. Alternatively, a microorganism can be cultured at a temperature selected to increase transcription of a particular gene and/or translation of a particular gene product such that transcription and/or translation are enhanced or increased.

The term "deregulated" or "deregulation" includes the alteration or modification of at least one gene in a microorganism that encodes an enzyme in a biosynthetic pathway, such that the level or activity of the biosynthetic enzyme in the microorganism is altered or modified. Preferably, at least one gene that encodes an enzyme in a biosynthetic pathway is altered or modified such that the gene product is enhanced or increased. The phrase "deregulated pathway" can also include a biosynthetic pathway in

25

30

20

25

30

35

which more than one gene that encodes an enzyme in a biosynthetic pathway is altered or modified such that the level or activity of more than one biosynthetic enzyme is altered or modified. The ability to "deregulate" a pathway (e.g., to simultaneously deregulate more than one gene in a given biosynthetic pathway) in a microorganism arises from the particular phenomenon of microorganisms in which more than one enzyme (e.g., two or three biosynthetic enzymes) are encoded by genes occurring adjacent to one another on a contiguous piece of genetic material termed an "operon".

The term "operon" includes a coordinated unit of gene expression that contains a promoter and possibly a regulatory element associated with one or more, preferably at least two, structural genes (e.g., genes encoding enzymes, for example, biosynthetic enzymes). Expression of the structural genes can be coordinately regulated, for example, by regulatory proteins binding to the regulatory element or by anti-termination of transcription. The structural genes can be transcribed to give a single mRNA that encodes all of the structural proteins. Due to the coordinated regulation of genes included in an operon, alteration or modification of the single promoter and/or regulatory element can result in alteration or modification of each gene product encoded by the operon. Alteration or modification of the regulatory element can include, but is not limited to removing the endogenous promoter and/or regulatory element(s), adding strong promoters, inducible promoters or multiple promoters or removing regulatory sequences such that expression of the gene products is modified, modifying the chromosomal location of the operon, altering nucleic acid sequences adjacent to the operon or within the operon such as a ribosome binding site, increasing the copy number of the operon, modifying proteins (e.g., regulatory proteins, suppressors, enhancers, transcriptional activators and the like) involved in transcription of the operon and/or translation of the gene products of the operon, or any other conventional means of deregulating expression of genes routine in the art (including but not limited to use of antisense nucleic acid molecules, for example, to block expression of repressor proteins). Deregulation can also involve altering the coding region of one or more genes to yield, for example, an enzyme that is feedback resistant or has a higher or lower specific activity.

A particularly preferred "recombinant" microorganism of the present invention has been genetically engineered to overexpress a bacterially-derived gene or gene product. The term "bacterially-derived" or "derived-from", for example bacteria, includes a gene which is naturally found in bacteria or a gene product (e.g., ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase, aspartate- α -decarboxylate, acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase or

dihydroxyacid dehydratase) which is encoded by a bacterial gene (e.g., encoded by panB, panE, panC, panD, ilvB, ilvN, alsS, ilvC, or ilvD).

The methodologies of the present invention feature recombinant microorganisms which overexpress at least one of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase or aspartate-a-decarboxylase. A particularly preferred recombinant microorganism of the present invention has been genetically engineered to overexpress a Bacillus (e.g., Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus, etc.) biosynthetic enzyme (e.g., has been engineered to overexpress at least one of B. subtilis ketopantoate reductase (the panE gene product) (e.g., ketopantoate reductase having the amino acid sequence of SEQ ID NO:30 or encoded by the nucleic acid sequence of SEQ ID NO:29), B. subtilis ketopantoate hydroxymethyltransferase (the panB gene product) (e.g., ketopantoate hydroxymethyltransferase having the amino acid sequence of SEQ ID NO:24 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:23), B. subtilis pantothenate synthetase (the panC gene product) (e.g., 15 pantothenate synthetase having the amino acid sequence of SEQ ID NO:26 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:25) and/or B. subtilis aspartate-a-decarboxylase (the panD gene product) (e.g., aspartate-adecarboxylase having the amino acid sequence of SEQ ID NO:28 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:27). 20

In an exemplary embodiment, the invention features a microorganism (e.g., a KPAR-O microorganism) that has been transformed with a vector comprising a panE nucleic acid sequence (e.g., a panE nucleic acid sequence as set forth in SEQ ID NO:29). In another embodiment, the invention features a microorganism that has been transformed with a vector comprising a panB nucleic acid sequence (e.g., a panB nucleic acid sequence as set forth in SEQ ID NO:23), a vector comprising a panC nucleic acid sequence (e.g., a panC nucleic acid sequence as set forth in SEQ ID NO:25) or a vector comprising a panD nucleic acid sequence (e.g., a panD nucleic acid sequence as set forth in SEQ ID NO:27). In yet another embodiment, the invention features a microorganism having a deregulated panBCD operon (e.g., SEQ ID NO:59).

Other preferred "recombinant" microorganisms of the present invention have a deregulated isoleucine-valine (ilv) pathway. The phrase "microorganism having a deregulated isoleucine-valine (ilv) pathway" includes a microorganism having an alteration or modification in at least one gene encoding an enzyme of the isoleucine-valine (ilv) pathway or having an alteration or modification in an operon including more than one gene encoding an enzyme of the isoleucine-valine (ilv) pathway. A preferred "microorganism having a deregulated isoleucine-valine (ilv) pathway" has been

25

30

20

25

30

genetically engineered to overexpress a Bacillus (e.g., B. subtilis) ilv biosynthetic enzyme (e.g., has been engineered to overexpress at least one of acetohydroxyacid synthetase (the ilvBN gene products or the alsS gene product) (e.g., acetohydroxyacid synthetase having subunits having the amino acid sequences of SEQ ID NO:32 and SEQ ID NO:34 or encoded by nucleic acid molecules having the nucleotide sequence of SEQ ID NO:31 and SEQ ID NO:33 or the nucleotide sequence of SEQ ID NO:58 from nucleotides 1-2246 or acetohydroxyacid synthetase encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:86), acetohydroxyacid isomeroreductase (the ilvC gene product) (e.g., acetohydroxyacid isomeroreductase having the amino acid sequence of SEQ ID NO:36 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:35), dihydroxyacid dehydratase (the ilvD gene product) (e.g., dihydroxyacid dehydratase having the amino acid sequence of SEQ ID NO:38 or encoded by a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:37), and/or has been transformed with a vector comprising an ilvBNC nucleic 15 acid sequence (SEQ ID NO:58, coding regions from nucleotides 1-1725, 1722-2246 and 2263-3291) and/or an ilvD nucleic acid sequence (SEQ ID NO:37).

In another preferred embodiment, a recombinant microorganism is designed or engineered such that a mutant CoaA and/or CoaX biosynthetic enzyme is expressed and at least one pantothenate biosynthetic enzyme and/or at least one isoleucine-valine biosynthetic enzyme is overexpressed or deregulated.

In another preferred embodiment, a microorganism of the present invention overexpresses or is mutated for a gene or biosynthetic enzyme (e.g., a CoA biosynthetic enzyme, pantothenate biosynthetic enzyme or isoleucine-valine biosynthetic enzyme) which is bacterially-derived. The term "bacterially-derived" or "derived-from", for example bacteria, includes a gene product (e.g., ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase, aspartate-α-decarboxylate, acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase, dihydroxyacid dehydratase or pantothenate kinase) which is encoded by a bacterial gene (e.g., panB, panE, panC, panD, ilvBN (or alsS), ilvC, ilvD, or encoded by coaA or coaX).

Still other preferred recombinant microorganisms of the present invention are mutant microorganisms. As used herein, the term "mutant microorganism" includes a recombinant microorganism that has been genetically engineered to express a mutated gene or protein that is normally or naturally expressed by the microorganism. Preferably, a mutant microorganism expresses a mutated gene or protein such that the microorganism exhibits an altered, modified or different phenotype (e.g., has been engineered to express a mutated CoaA biosynthetic enzyme, for example, pantothenate kinase). In one embodiment, a mutant microorganism is designed or engineered such

that it includes a mutant coaX gene, as defined herein. In another embodiment, a recombinant microorganism is designed or engineered such that it includes a mutant coal gene, as defined herein. In another embodiment, a mutant microorganism is designed or engineered such that a coaX gene has been deleted (i.e., the protein encoded by the coaX gene is not produced). In another embodiment, a mutant microorganism is designed or engineered such that a coaA gene has been deleted (i.e., the protein encoded by the coal gene is not produced). Preferably, a mutant microorganism has a mutant coaX gene or a mutant coaA gene, or has been engineered to have a coaX gene and/or coaA deleted, such that that the mutant microorganism encodes a "reduced pantothenate kinase activity". In the context of a whole microorganism, a "reduced pantothenate kinase activity" can be determined by measuring or assaying for a decrease in an intermediate or product of the CoA biosynthetic pathway, for example, measuring or assaying for 4'-phosphopantothenate, 4'-phosphopantothenylcysteine, 4'phosphopantetheine, dephosphocoenzyme A, Coenzyme A, apo-acyl carrier protein (apo-ACP) or holo-acyl carrier protein (ACP) in the microorganism (e.g., in a lysate 15 isolated or derived from the microorganism) or in the medium in which the microorganism is cultured (see e.g., Figure 16). Alternatively, a "reduced pantothenate ... kinase activity" can be determined by measuring or assaying for decreased growth of the microorganism. Alternatively, a "reduced pantothenate kinase activity" can be determined by measuring or assaying for an increase in a panto-compound (e.g., 20 pantothenate) in the microorganism or surrounding media, as panto-compounds lie upstream of the CoA biosynthetic pathway, the first step of which is catalyzed by pantothenate kinase. The invention also features recombinant microorganisms that, in addition to having reduced pantothenate kinase activity (e.g., expressing mutant coal and/or mutant coaX genes) have a deregulated pantothenate biosynthesis pathway and/or 25 a deregulated isoleucine-valine (ilv) biosynthetic pathway.

In one embodiment, a recombinant microorganism of the present invention is a Gram positive organism (e.g., a microorganism which retains basic dye, for example, crystal violet, due to the presence of a Gram-positive wall surrounding the microorganism). In a preferred embodiment, the recombinant microorganism is a microorganism belonging to a genus selected from the group consisting of Bacillus, Cornyebacterium, Lactobacillus, Lactococci and Streptomyces. In a more preferred embodiment, the recombinant microorganism is of the genus Bacillus. In another preferred embodiment, the recombinant microorganism is selected from the group consisting of Bacillus subtilis, Bacillus lentimorbus, Bacillus lentus, Bacillus firmus, Bacillus pantothenticus, Bacillus amyloliquefaciens, Bacillus cereus, Bacillus circulans, Bacillus coagulans, Bacillus licheniformis, Bacillus megaterium, Bacillus pumilus,

25

30

35

0101770A2 1A.

Bacillus thuringiensis, and other Group 1 Bacillus species, for example, as characterized by 16S rRNA type (Priest (1993) in Bacillus subtilis and Other Gram-Positive Bacteria eds. Sonenshein et al., ASM, Washington, D.C., p. 6). In another preferred embodiment, the recombinant microorganism is Bacillus brevis or Bacillus stearothermophilus. In another preferred embodiment, the recombinant microorganism is selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus.

In another embodiment, the recombinant microorganism is a Gram negative (excludes basic dye) organism. In a preferred embodiment, the recombinant microorganism is a microorganism belonging to a genus selected from the group consisting of Salmonella, Escherichia, Klebsiella, Serratia, and Proteus. In a more preferred embodiment, the recombinant microorganism is of the genus Escherichia. In an even more preferred embodiment, the recombinant microorganism is Escherichia coli. In another embodiment, the recombinant microorganism is Saccharomyces (e.g., S. 15 cerevisiae).

An important aspect of the present invention involves culturing the recombinant microorganisms described herein, such that a desired compound (e.g., a desired pantocompound) is produced. The term "culturing" includes maintaining and/or growing a living microorganism of the present invention (e.g., maintaining and/or growing a culture or strain). In one embodiment, a microorganism of the invention is cultured in liquid media. In another embodiment, a microorganism of the invention is cultured in solid media or semi-solid media. In a preferred embodiment, a microorganism of the invention is cultured in media (e.g., a sterile, liquid media) comprising nutrients essential or beneficial to the maintenance and/or growth of the microorganism (e.g., carbon sources or carbon substrate, for example complex carbohydrates such as bean or grain meal, starches, sugars, sugar alcohols, hydrocarbons, oils, fats, fatty acids, organic acids and alcohols; nitrogen sources, for example, vegetable proteins, peptones, peptides and amino acids derived from grains, beans and tubers, proteins, peptides and amino acids derived form animal sources such as meat, milk and animal byproducts such as peptones, meat extracts and casein hydrolysates; inorganic nitrogen sources such as urea, ammonium sulfate, ammonium chloride, ammonium nitrate and ammonium phosphate; phosphorus sources, for example, phosphoric acid, sodium and potassium salts thereof; trace elements, for example, magnesium, iron, manganese, calcium. copper, zinc, boron, molybdenum, and/or cobalt salts; as well as growth factors such as amino acids, vitamins, growth promoters and the like).

Preferably, microorganisms of the present invention are cultured under controlled pH. The term "controlled pH" includes any pH which results in production of the desired product (e.g., a panto-compound). In one embodiment, microorganisms are cultured at a pH of about 7. In another embodiment, microorganisms are cultured at a pH of between 6.0 and 8.5. The desired pH may be maintained by any number of methods known to those skilled in the art.

Also preferably, microorganisms of the present invention are cultured under controlled aeration. The term "controlled aeration" includes sufficient aeration (e.g., oxygen) to result in production of the desired product (e.g., panto-compound). In one embodiment, aeration is controlled by regulating oxygen levels in the culture, for example, by regulating the amount of oxygen dissolved in culture media. Preferably, aeration of the culture is controlled by agitating the culture. Agitation may be provided by a propeller or similar mechanical agitation equipment, by revolving or shaking the growth vessel (e.g., fermentor) or by various pumping equipment. Aeration may be further controlled by the passage of sterile air or oxygen through the medium (e.g., through the fermentation mixture). Also preferably, microorganisms of the present invention are cultured without excess foaming (e.g., via addition of antifoaming agents).

Moreover, microorganisms of the present invention can be cultured under controlled temperatures. The term "controlled temperature" includes any temperature which results in production of the desired product (e.g., a panto-compound). In one embodiment, controlled temperatures include temperatures between 15°C and 95°C. In another embodiment, controlled temperatures include temperatures between 15°C and 70°C. Preferred temperatures are between 20°C and 55°C, more preferably between 30°C and 45°C or between 30°C and 50°C.

Microorganisms can be cultured (e.g., maintained and/or grown) in liquid media and preferably are cultured, either continuously or intermittently, by conventional culturing methods such as standing culture, test tube culture, shaking culture (e.g., rotary shaking culture, shake flask culture, etc.), aeration spinner culture, or fermentation. In a preferred embodiment, the microorganisms are cultured in shake flasks. In a more preferred embodiment, the microorganisms are cultured in a fermentor (e.g., a fermentation process). Fermentation processes of the present invention include, but are not limited to, batch, fed-batch and continuous methods of fermentation. The phrase "batch process" or "batch fermentation" refers to a closed system in which the composition of media, nutrients, supplemental additives and the like is set at the beginning of the fermentation and not subject to alteration during the fermentation, however, attempts may be made to control such factors as pH and oxygen concentration to prevent excess media acidification and/or microorganism death. The phrase "fed-

20

25

30

batch process" or "fed-batch" fermentation refers to a batch fermentation with the exception that one or more substrates or supplements are added (e.g., added in increments or continuously) as the fermentation progresses. The phrase "continuous process" or "continuous fermentation" refers to a system in which a defined fermentation media is added continuously to a fermentor and an equal amount of used or "conditioned" media is simultaneously removed, preferably for recovery of the desired product (e.g., panto-compound). A variety of such processes have been developed and are well-known in the art.

The phrase "culturing under conditions such that a desired compound (e.g., a panto-compound, for example, pantothenate) is produced" includes maintaining and/or growing microorganisms under conditions (e.g., temperature, pressure, pH, duration, etc.) appropriate or sufficient to obtain production of the desired compound or to obtain desired yields of the particular compound being produced. For example, culturing is continued for a time sufficient to produce the desired amount of a panto-compound (e.g., 15 pantothenate, pantoate or β -alanine). Preferably, culturing is continued for a time sufficient to substantially reach maximal production of the panto-compound. In one embodiment, culturing is continued for about 12 to 24 hours. In another embodiment, culturing is continued for about 24 to 36 hours, 36 to 48 hours, 48 to 72 hours, 72 to 96 hours, 96 to 120 hours, 120 to 144 hours, or greater than 144 hours. In another embodiment, culturing is continued for a time sufficient to reach production yields of panto-compound, for example, cells are cultured such that at least about 15 to 20 g/L of panto-compound are produced, at least about 20 to 25 g/L panto-compound are produced, at least about 25 to 30 g/L panto-compound are produced, at least about 30 to 35 g/L panto-compound are produced, at least about 35 to 40 g/L panto-compound are produced (e.g., at least about 37 g/L panto-compound) or at least about 40 to 50 g/L panto compound are produced. In yet another embodiment, microorganisms are cultured under conditions such that a preferred yield of panto-compound, for example, a yield within a range set forth above, is produced in about 24 hours, in about 36 hours, in about 48 hours, in about 72 hours, or in about 96 hours.

The methodology of the present invention can further include a step of recovering a desired compound (e.g., a panto-compound). The term "recovering" a desired compound (e.g., a panto-compound) includes extracting, harvesting, isolating or purifying the compound from culture media. Recovering the compound can be performed according to any conventional isolation or purification methodology known in the art including, but not limited to, treatment with a conventional resin (e.g., anion or cation exchange resin, non-ionic adsorption resin, etc.), treatment with a conventional adsorbent (e.g., activated charcoal, silicic acid, silica gel, cellulose, alumina, etc.),

10

20

25

30

15

20

30

alteration of pH, solvent extraction (e.g., with a conventional solvent such as an alcohol, ethyl acetate, hexane and the like), dialysis, filtration, concentration, crystallization, recrystallization, pH adjustment, lyophilization and the like. For example, a compound (e.g., a panto-compound) can be recovered from culture media by first removing the microorganisms from the culture. Media is then passed through or over a cation exchange resin to remove unwanted cations and then through or over an anion exchange resin to remove unwanted inorganic anions and organic acids having stronger acidities than the panto-compound of interest (e.g., pantothenate). The resulting panto-compound (e.g., pantothenate) can subsequently be converted to a pantothenate salt (e.g., calcium pantothenate) as described herein.

Preferably, a desired compound of the present invention is "extracted", "isolated" or "purified" such that the resulting preparation is substantially free of other components (e.g., free of media components and/or fermentation byproducts). The language "substantially free of other components" includes preparations of desired compound in which the compound is separated (e.g., purified or partially purified) from media components or fermentation byproducts of the culture from which it is produced. In one embodiment, the preparation has greater than about 80% (by dry weight) of the desired compound (e.g., less than about 20% of other media-components or fermentation byproducts), more preferably greater than about 90% of the desired compound (e.g., less than about 10% of other media components or fermentation byproducts), still more preferably greater than about 95% of the desired compound (e.g., less than about 5% of other media components or fermentation byproducts), and most preferably greater than about 98-99% desired compound (e.g., less than about 1-2% other media components or fermentation byproducts). When the desired compound is a panto-compound that has been derivatized to a salt (e.g. a pantothenate salt or pantoate salt), the panto-compound is preferably further free (e.g., substantially free) of chemical contaminants associated with the formation of the salt. When the desired compound is a panto-compound that has been derivatized to an alcohol, the panto-compound is preferably further free (e.g., substantially free) of chemical contaminants associated with the formation of the alcohol.

In an alternative embodiment, the desired panto-compound is not purified from the microorganism, for example, when the microorganism is biologically non-hazardous (e.g., safe). For example, the entire culture (or culture supernatant) can be used as a source of product (e.g., crude product). In one embodiment, the culture (or culture supernatant) supernatant is used without modification. In another embodiment, the culture (or culture supernatant) is concentrated. In yet another embodiment, the culture (or culture supernatant) is dried or lyophilized.

Panto-Compound Production Methodologies Featuring Ketopantoate Reductase-Overexpressing Microorganisms

One aspect of the invention features methods of producing a panto-compounds that involve culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism under conditions such that the panto-compound is produced. The term "ketopantoate reductase-overexpressing (KPAR-O) microorganism" includes a microorganism which has been manipulated such that ketopantoate reductase is overexpressed (e.g., a B. subtilis ketopantoate reductase protein having the amino acid sequence of SEQ ID NO:30) and/or has been transformed with a vector comprising a panEl nucleic acid sequence (e.g., a B. subtilis panEl nucleic acid sequence as set forth in SEQ ID NO:29). In one embodiment, the panto-compound is pantothenate. In another embodiment, the panto-compound is pantoate. In another embodiment, the ketopantoate reductase is bacterial-derived. In another embodiemnt, the ketopantoate reductase is derived from Bacillus (e.g., is derived from Bacillus subtilis). In yet 15 another embodiment, the KPAR-O microorganism is Gram positive. In yet another embodiment, the KPAR-O microorganism is a microorganism belonging to a genus selected from the group consisting of Bacillus, Cornyebacterium, Lactobacillus, Lactococci and Streptomyces. In a preferred embodiemnt, the KPAR-O microorganism is of the genus Bacillus. In a more preferred embodiment, the KPAR-O microorganism is selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis and Bacillus pumilus. In a particularly preferred embodiemnt, the KPAR-O microorganism is Bacillus subtilis.

In still other embodiments, the KPAR-O microorganism further overexpresses at least one pantothenate biosynthetic enzyme in addition to ketopantoate reductase. In an exemplary embodiment, the KPAR-O microorganism further overexpresses at least one of ketopantoate hydroxymethyltransferase, pantothenate synthetase and aspartate-αdecarboxylase. Also featured are methods of producing panto-compounds, for example, methods that involve culturing a KPAR-O microorganism, which further include the step of recovering the panto-compound.

30

35

20

25

Methods of Producing Panto-Compounds Independent of Precursor Feed III. Requirements

Depending on the biosynthetic enzyme or combination of biosynthetic enzymes manipulated, it may be desirable or necessary to provide (e.g., feed) microorganisms of the present invention at least one pantothenate biosynthetic precursor such that pantothenate or other desired panto-compounds are produced. The term "pantothenate biosynthetic precursor" or "precursor" includes an agent or compound which, when

20

25

35

provided to, brought into contact with, or included in the culture medium of a microorganism, serves to enhance or increase pantothenate biosynthesis. In one embodiment, the pantothenate biosynthetic precursor or precursor is aspartate. In another embodiment, the pantothenate biosynthetic precursor or precursor is \(\beta-alanine.

The amount of aspartate or β-alanine added is preferably an amount that results in a concentration in the culture medium sufficient to enhance productivity of the microorganism (e.g., a concentration sufficient to enhance production of a pantocompound, for example, β-alanine, ketopantoate, pantoate or pantothenate).

Pantothenate biosynthetic precursors of the present invention can be added in the form of a concentrated solution or suspension (e.g., in a suitable solvent such as water or buffer) or in the form of a solid (e.g., in the form of a powder). Moreover, pantothenate biosynthetic precursors of the present invention can be added as a single aliquot, continuously or intermittently over a given period of time.

In yet another embodiment, the pantothenate biosynthetic precursor is valine, see e.g., Example III. In yet another embodiment, the pantothenate biosynthetic precursor is α -ketoisovalerate. Preferably, valine or α -ketoisovalerate is added in an amount that results in a concentration in the medium sufficient for production of the desired product (e.g., panto-compound) to occur. Pantothenate biosynthetic precursors are also referred to herein as "supplemental pantothenate biosynthetic substrates".

Providing pantothenate biosynthetic precursors in the pantothenate biosynthetic methodologies of the present invention, can be associated with high costs, for example, when the methodologies are used to produce high yields of panto-compounds. Accordingly, preferred methodologies of the present invention feature microorganisms having at least one biosynthetic enzyme or combination of biosynthetic enzymes (e.g., at least one pantothenate biosynthetic enzyme and/or valine-isoleucine biosynthetic enzyme) manipulated such that pantothenate or other desired panto-compounds are produced in a manner independent of precursor feed. The phrase "a manner independent of precursor feed", for example, when referring to a method for producing a desired compound (e.g., a panto-compound), includes an approach to or a mode of producing the desired compound that does not depend or rely on precursors being provided (e.g., fed) to the microorganism being utilized to produce the desired compound. For example, microorganisms featured in the methodologies of the present invention can be used to produce panto-compounds in a manner requiring no feeding of the precursors aspartate, β-alanine, valine and/or α-KIV.

Alternative preferred methodologies of the present invention feature microorganisms having at least one biosynthetic enzyme or combination of biosynthetic enzymes manipulated such that pantothenate or other desired panto-compounds are

20

25

30

35

produced in a manner substantially independent of precursor feed. The phrase "a manner substantially independent of precursor feed" includes an approach to or a method of producing the desired compound that depends or relies to a lesser extent on precursors being provided (e.g., fed) to the microorganism being utilized. For example, microorganisms featured in the methodologies of the present invention can be used to produce panto-compounds in a manner requiring feeding of substantially reduced amounts of the precursors aspartate, β-alanine, valine and/or α-KIV. In one embodiment, the invention features methods of producing panto-compounds (e.g., pantothenate) in a manner that requires feeding of less than 5%-10% of the amount of precursor required by a control microorganism (e.g., a microorganism that is dependent, for example is wholly dependent, on precursor feed to efficiently produce the desired compound). In another embodiment, the invention features methods of producing panto-compounds in a manner that requires feeding of less than 15-20% of the amount of precursor required by a control microorganism. In another embodiment, the 15 invention features methods of producing panto-compounds in a manner that requires feeding of less than 25-30%, 35-40%, 45-50% or 55-60% of the amount of precursor required by a control microorganism. As described in Examples I-III herein, particular microorganisms featured in the methodologies of the present invention require, for example, 5 g/L of aspartate, β -alanine, valine or α -KIV (e.g., in test tube or in shake flask cultures). Accordingly, in a preferred embodiment, the present invention features methods of producing panto-compounds (e.g., pantothenate) in a manner requiring feeding of less than 0.25 g/L, 0.5 g/L, 0.75 g/L, 1 g/L, 1.25 g/L, 1.5 g/L, 1.75 g/L, 2 g/L, 2.25 g/L, 2.5 g/L, 2.75 g/L or 3 g/L.

Preferred methods of producing desired compounds (e.g., panto-compounds) in a manner independent of precursor feed or alternatively, in a manner substantially independent of precursor feed, involve culturing microorganisms which have been manipulated (e.g., designed or engineered, for example, genetically engineered) such that expression of at least one pantothenate biosynthetic enzyme, and/or at least one isoleucine-valine biosynthetic enzyme, is modified. For example, in one embodiment, a microorganism is manipulated (e.g., designed or engineered) such that the production of at least one pantothenate biosynthetic enzyme, and/or at least one isoleucine/valine biosynthetic enzyme is deregulated. In a preferred embodiment, a microorganism is manipulated (e.g., designed or engineered) such that it has a deregulated biosynthetic pathway, for example, a deregulated pantothenate biosynthesis pathway and/or a deregulated isoleucine-valine biosynthetic pathway, as defined herein. In another preferred embodiment, a microorganism is manipulated (e.g., designed or engineered)

such that at least one pantothenate biosynthetic enzyme, and/or at least one isoleucinevaline biosynthetic enzyme is overexpressed.

Preferred methods of producing desired compounds (e.g., panto-compounds) in a manner independent of precursor feed or alternatively, in a manner substantially independent of precursor feed, are as follows. In one embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism having a deregulated isoleucine-valine (ilv) pathway under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate-a-decarboxylase-overexpressing (AaD-O) microorganism having a deregulated pantothenate (pan) pathway and a deregulated isoleucine-valine (ilv) pathway, under conditions such that pantothenate is produced. In another embodiment, the invention features a method of producing pantothenate in a manner independent of aspartate or β-alanine feed comprising culturing an aspartate-αdecarboxylase-overexpressing (AaD-O) microorganism under conditions such that pantothenate is produced. In yet another embodiment, the invention features a method of producing pantothenate in a manner independent of valine or a-ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced.

The term "aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism" includes a microorganism which has been manipulated such that aspartate- α -decarboxylase is overexpressed. A preferred "aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism" has been transformed with a vector comprising a *B. subtilis panD* nucleic acid sequence (*e.g.*, a *panD* nucleic acid sequence that encodes an aspartate- α -decarboxylase protein having the amino acid sequence of SEQ ID NO:28, for example, a *panD* nucleic acid sequence as set forth in SEQ ID NO:27).

The phrase "microorganism having a deregulated isoleucine-valine (ilv) pathway" includes a microorganism having an alteration or modification in at least one gene encoding an enzyme of the isoleucine-valine (ilv) pathway or having an alteration or modification in an operon including more than one gene encoding an enzyme of the isoleucine-valine (ilv) pathway. A preferred "microorganism having a deregulated isoleucine-valine (ilv) pathway" overexpresses acetohydroxyacid synthetase (e.g., acetohydroxyacid synthetase having subunits having the amino acid sequences of SEQ ID NO:32 and SEQ ID NO:34 or acetohydroxyacid synthetase having the amino acid sequence of SEQ ID NO:87), acetohydroxyacid isomeroreductase (having the amino acid sequence of SEQ ID NO:36), or dihydroxyacid dehydratase [having the amino acid

15

20

25

. ;..

sequence of SEQ ID NO:38) and/or has been transformed with a vector comprising *ilvB*, *ilvN*, *ilvC*, *ilvBN*, *ilvBNC* or *alsS* nucleic acid sequences (SEQ ID NO:31, SEQ ID NO:33, SEQ ID NO:35, nucleotides 1-2246 of SEQ ID NO:58, SEQ ID NO:58 having coding regions from nucleotides 1-1725, 1722-2246 and 2263-3291, or SEQ ID NO:86, respectively) and/or an *ilvD* nucleic acid sequence (SEQ ID NO:37).

IV. High Yield Production Methodologies

A particularly preferred embodiment of the present invention is a high yield production method for producing pantothenate comprising culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly 10 high yield. The phrase "high yield production method", for example, a high yield production method for producing a desired compound (e.g., for producing a pantocompound) includes a method that results in production of the desired compound at a level which is elevated or above what is usual for comparable production methods. 15 Preferably, a high yield production method results in production of the desired compound at a significantly high yield. The phrase "significantly high yield" includes a level of production or yield which is sufficiently elevated or above what is usual for comparable production methods, for example, which is elevated to a level sufficient for commercial production of the desired product (e.g., production of the product at a 20 commercially feasible cost). In one embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 2 g/L. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism 25 under conditions such that pantothenate is produced at a level greater than 10 g/L. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 20 g/L. In yet another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 30 g/L. In yet another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 40 g/L.

The invention further features a high yield production method for producing a desired compound (e.g., for producing a panto-compound) that involves culturing a manipulated microorganism under conditions such that a sufficiently elevated level of

compound is produced within a commercially desireable period of time. In an exemplary embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under 'conditions such that pantothenate is produced at a level greater than 15-20 g/L in 36 hours. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 25-30 g/L in 48 hours. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level-greater than 35-40 g/L in 72 hours, for example, greater that 37 g/L in 72 hours. In another embodiment, the invention features a high yield production method of producing pantothenate that includes culturing a manipulated microorganism under conditions such that pantothenate is produced at a level greater than 30-40 g/L in 60 hours, for example, greater that 30, 35 or 40 g/L in 60 hours. Values and ranges included and/or intermediate within the ranges set forth herein are also intended to be within the scope of the present invention. For example, pantothenate production at levels of at least 31, 32, 33, 34, 35, 36, 37, 38 and 39 g/L in 60 hours are intended to be included within the range of 30-40 g/L in 60 hours. In another example, ranges of 30-35 g/L or 35-40 g/L are intended to be included within the range of 30-40 g/L in 60 hours. Moreover, the skilled artisan will appreciate that culturing a manipulated microorganism to achieve a production level of, for example, "30-40 g/L in 60 hours" includes culturing the microorganism for additional time periods (e.g., time periods longer than 60 hours), optionally resulting in even higher yields of pantothenate being produced.

25

30

35

20

V. Panto-Compound Production Methodologies Featuring Pantothenate Kinase Mutant Microorganisms

The present invention relates to methods of producing pantothenate using microorganisms engineered to produce high yields of pantothenate as well as other panto-compounds. Cells overproducing pantothenate result in high intracellular pantothenate levels that could overcome the feedback inhibition of pantothenate kinase by CoA, leading to overproduction of CoA. Besides consuming pantothenate, increased synthesis of CoA may cause increased feedback inhibition of the PanB, PanD, PanE or PanC reaction, thereby limiting pantothenate production. Accordingly, a reduction in pantothenate kinase activity may lead to a decrease in CoA levels with resulting increases in PanB, PanD, PanE or PanC activity and pantothenate production.

10

15

20

25

30

.,

Thus, certain methodologies of the present invention are based, at least in part, on the identification and characterization of the *B. subtilis coaA* gene and the demonstration that the gene is neither essential for *B. subtilis* growth (*i.e.*, deletion of the coaA gene from the chromosome of *B. subtilis* is not a lethal event) nor for pantothenate kinase activity in *B. subtilis*. A second pantothenate kinase-encoding gene has been identified and characterized in *B. subtilis*, and is termed "coaX". This gene complements an *E. coli* mutant that contains a temperature sensitive pantothenate kinase and is not related by homology to any previously known pantothenate kinase gene.

In one aspect, the methodologies of the invention feature recombinant microorganisms that include the coaX gene or that include a mutant coaX gene, having reduced pantothenate kinase activity. In one embodiment, the methodologies feature such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway. In another embodiment, the methodologies feature such recombinant microorganisms further having a deregulated isoleucine-valine (ilv) pathway. In a preferred embodiment, the microorganisms belong to the genus Bacillus (e.g., B. subtilis).

The methodologies of the invention also feature recombinant microorganisms (e.g., microorganisms belong to the genus Bacillus, for example, B. subtilis) that include the coal gene or that include a mutant coal gene, optionally including a coal gene or mutant thereof, having reduced pantothenate kinase activity. In one embodiment, the methodologies feature such recombinant microorganisms further having a deregulated pantothenate biosynthetic pathway or having a deregulated isoleucine-valine (ilv) pathway. Also featured are vectors that include isolated coal or coal genes as well as mutant coal and/or coal genes. Isolated nucleic acid molecules that include isolated coal genes or mutant coal genes are features in addition to isolated Coal proteins and mutant Coal proteins.

The above-described nucleic acid molecules (e.g., genes), proteins, vectors, and recombinant microorganisms (e.g., mutant microorganisms), are particularly suited for use in methods of producing panto-compounds and/or methods of enhancing panto-compound production. In one embodiment, the invention features a method for producing a panto-compound (e.g., pantothenate) that includes culturing a pantothenate kinase mutant (e.g., a recombinant microorganism that misexpresses, e.g., is mutated for, pantothenate kinase, as defined herein) under conditions such that panto-compound is produced. In another embodiment, the invention features a method for enhancing production of a panto-compound (e.g., pantothenate) that includes culturing a pantothenate kinase mutant (e.g., a recombinant microorganism that misexpresses, e.g., is mutated for, pantothenate kinase, as defined herein) under conditions such that

20

25

30

35

Production

production of the panto-compound is produced. As used herein, the term "enhancing" (for example, in the context of the phrase "enhancing production") includes increasing the level or rate of production of panto-compound (e.g., pantothenate) as compared to the level or rate of production in a non-mutant microorganism (e.g., a microorganism having a normal pantothenate kinase gene(s) and/or having normal pantothenate production rates and/or levels.

Preferably, the level of panto-compound produced in methodologies featuring the pantothenate kinase mutants of the present invention is increased by at least 5% as compared to the level produced by a non-mutant (e.g., a recombinant microorganism expressing non-mutated pantothenate kinase). Even more preferably, the level of panto-compound is increased 10% as compared to methodologies featuring non-mutants. Even more preferably, panto-compound levels (e.g., pantothenate levels) are increased 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, are increased 2-fold, 5-fold, 10-fold, 50-fold, 100-fold or more as compared to methodologies featuring non-mutants.

VI. Additional Mutations Resulting in Enhanced Panto-Compound

The methodologies of the present invention further can include, for example in addition to overexpressing or deregulating a pantothenate biosynthetic enzyme and/or an isoleucine-valine biosynthetic enzyme, or in addition to mutating a pantothenate-kinase encoding gene, deleting or mutating an enzyme that catalyzes the conversion of key pantothenate biosynthesis substrates or precursors to unwanted or undesirable products. For example, mutating the *ilvE* gene (Kuramitsu et al. (1985) J. Biochem. 97:993-999) or a homologue thereof (SEQ ID NO:62 or SEQ ID NO:64), thereby limiting the conversion of a-ketoisovalerate to valine, in addition to mutating a pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or increased production of panto-compound. Alternatively, mutating the ansB gene (Sun and Seflow (1991) J. Bacteriol. 173:3831-3845) or a homologue thereof (SEQ ID NO:66), thereby limiting the degradation of aspartate, in addition to mutating a pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or increased production of pantocompound. Alternatively, mutating the alsD gene (Renna et al. (1993) J. Bacteriol. 175:3863-3875) or a homologue thereof (SEQ ID NO:68), thereby limiting the conversion of acetolactate to acetoin, in addition to mutating a pantothenate kinase encoding enzyme, is predicted to result in even further enhanced or increased production of panto-compound. Alternatively, mutating the avtA gene encoding alanine-valine transaminase or a homologue thereof, thereby limiting the conversion of aketoisovalerate to valine, in addition to mutating a pantothenate kinase encoding

20

25

30

35

enzyme, is predicted to result in even further enhanced or increased production of panto-compound. Mutating the avtA gene can include mutating, for example, an avtA gene having the nucleotide sequence of SEQ ID NO:70 (e.g., the E. coli avtA gene), or a structural homolog thereof (e.g., a homologue encoding a protein having 30-40%, 40-50%, 50-60%, 60-70%, 70-80%, 80-90%, 90-95% or more identity with the amino acid sequence of SEQ ID NO:71) or a functional homologue (e.g., a gene encoding a structurally unrelated protein having alanine-valine transmainase activity. Such mutations can be accomplished using the methodologies as exemplified in the Examples (e.g., Examples XIII, XV, XVI and XVII).

Accordingly, in one embodiment, the invention features a method of producing a panto-compound which includes culturing a microorganism having a mutant pantothenate kinase-encoding gene and which further has a deletion or mutation in an avtA, ilvE, ansB, and/or alsD gene, or homologue thereof. In another embodiment, the invention features a method of producing a panto-compound which includes culturing a microorganism having a mutant pantothenate-kinase encoding gene and a deregulated pantothenate biosynthetic pathway enzyme and which further has a deletion or mutation in an avtA, ilvE, ansB, and/or alsD gene, or homologue thereof. In another embodiment, the invention features a method of producing a panto-compound which includes culturing a microorganism having a mutant pantothenate-kinase encoding gene and a deregulated isoleucine-valine biosynthetic pathway enzyme and which further has a deletion or mutation in an avtA, ilvE, ansB, and/or alsD gene, or homologue thereof.

Mutating the alsD gene can be particularly useful when accomplished in conjunction with overexpression or deregulation of the alsS gene, for example, to prevent carbon (e.g., acetolactate) from being drawn away from the precursor pool utilized for α-KIV production. Accordingly, to maximize the contribution of the als locus to panto-compound production, it is desirable to disrupt the alsD gene in addition to overexpressing the alsS gene. To disrupt the alsD gene, appropriate fragments of the als operon, flanking the alsD gene, are amplified by PCR and cloned to provide homology for creating the disruptions. A drug resistance gene, such as the cat gene, is cloned between the flanking DNA fragments in place of the alsD gene, and the linearized DNA is transformed into a pantothenate production strain such as PA824, selecting for drug-resistance. To overexpress alsS, the alsS coding sequence (e.g., an alsS coding sequence that has been engineered by PCR for expression) is cloned into an expression vector. Vectors which express alsS (or alternatively, vectors which express alsS plus ilvC) are the introduced into panto-compound production strains (e.g., the pantothenate producing strain PA824).

20

25

30

The methodologies of the present invention further can include, for example in addition to overexpressing or deregulating a pantothenate biosynthetic enzyme and/or an isoleucine-valine biosynthetic enzyme, or in addition to mutating a pantothenate-kinase encoding gene, deleting or mutating an enzyme that catalyzes the conversion of desired panto-compounds to unwanted or undesireable downstream products.

VII. Isolated Nucleic Acid Molecules and Genes

Another aspect of the present invention features isolated nucleic acid molecules that encode Bacillus proteins (e.g., B. subtilis proteins), for example, Bacillus pantothenate biosynthetic enzymes (e.g., B. subtilis pantothenate biosynthetic enzymes) or Bacillus valine-isoleucine biosynthetic enzymes (e.g., B. subtilis valine-isoleucine biosynthetic enzymes). Also featured are isolated coaX and/or coaA nucleic acid molecules (e.g., isolated coaX and/or coaA genes) as well as isolated nucleic acid molecules that include such coaX and/or coaA nucleic acid molecules or genes.

The term "nucleic acid molecule" includes DNA molecules (e.g., linear, circular, cDNA or chromosomal DNA) and RNA molecules (e.g., tRNA, rRNA, mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA. The term "isolated" nucleic acid molecule includes a nucleic acid molecule which is free of sequences which naturally flank the nucleic acid molecule (i.e., sequences located at the 5' and 3' ends of the nucleic acid molecule) in the chromosomal DNA of the organism from which the nucleic acid is derived. In various embodiments, an isolated nucleic acid molecule can contain less than about 10 kb, 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb, 0.1 kb, 50 bp, 25 bp or 10 bp of nucleotide sequences which naturally flank the nucleic acid molecule in chromosomal DNA of the microorganism from which the nucleic acid molecule is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular materials when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

The term "gene", as used herein, includes a nucleic acid molecule (e.g., a DNA molecule or segment thereof), for example, a protein or RNA-encoding nucleic acid molecule, that in an organism, is separated from another gene or other genes, by intergenic DNA (i.e., intervening or spacer DNA which naturally flanks the gene and/or separates genes in the chromosomal DNA of the organism). A gene may direct synthesis of an enzyme or other protein molecule (e.g., may comprise coding sequences, for example, a contiguous open reading frame (ORF) which encodes a protein) or may itself be functional in the organism. A gene in an organism, may be clustered in an

operon, as defined herein, said operon being separated from other genes and/or operons by the intergenic DNA. Individual genes contained within an operon may overlap without intergenic DNA between said individual genes. An "isolated gene", as used herein, includes a gene which is essentially free of sequences which naturally flank the gene in the chromosomal DNA of the organism from which the gene is derived (i.e., is free of adjacent coding sequences which encode a second or distinct protein or RNA molecule, adjacent structural sequences or the like) and optionally includes 5' and 3' regulatory sequences, for example promoter sequences and/or terminator sequences. In one embodiment, an isolated gene includes predominantly coding sequences for a protein (e.g., sequences which encode Bacillus proteins). In another embodiment, an isolated gene includes coding sequences for a protein (e.g., for a Bacillus protein) and adjacent 5' and/or 3' regulatory sequences from the chromosomal DNA of the organism from which the gene is derived (e.g., adjacent 5' and/or 3' Bacillus regulatory sequences). Preferably, an isolated gene contains less than about 10 kb, 5 kb, 2 kb, 1 kb, 0.5 kb, 0.2 kb, 0.1 kb, 50 bp, 25 bp or 10 bp of nucleotide sequences which naturally flank the gene in the chromosomal DNA of the organism from which the gene is derived.

In one aspect, the present invention features isolated panB nucleic acid sequences or genes, isolated panC nucleic acid sequences or genes, isolated panD nucleic acid sequences or genes, isolated panE nucleic acid sequences or genes, isolated ilvB, ilvN, ilvBN nucleic acid sequences or genes, isolated alsS nucleic acid sequences or genes, isolated ilvC nucleic acid sequences or genes and/or isolated ilvD nucleic acid sequences or genes.

In a preferred embodiment, the nucleic acid or gene is derived from Bacillus (e.g., is Bacillus-derived). The term "derived from Bacillus" or "Bacillus-derived" includes a nucleic acid or gene which is naturally found in microorganisms of the genus Bacillus. Preferably, the nucleic acid or gene is derived from a microorganism selected from the group consisting of Bacillus subtilis, Bacillus lentimorbus, Bacillus lentus, Bacillus firmus, Bacillus pantothenticus, Bacillus amyloliquefaciens, Bacillus cereus, Bacillus circulans, Bacillus coagulans, Bacillus licheniformis, Bacillus megaterium, Bacillus pumilus, Bacillus thuringiensis, and other Group 1 Bacillus species, for example, as characterized by 16S rRNA type (Priest, supra). In another preferred embodiment, the nucleic acid or gene is derived from Bacillus brevis or Bacillus stearothermophilus. In another preferred embodiment, the nucleic acid molecules and/or genes of the present invention are derived from a microorganism selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens, Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus. In a particularly preferred

WO 01/021772

10

15

20

embodiment, the nucleic acid or gene is derived from Bacillus subtilis (e.g., is Bacillus subtilis-derived). The term "derived from Bacillus subtilis" or "Bacillus subtilis-derived" includes a nucleic acid or gene which is naturally found in Bacillus subtilis. In yet another preferred embodiment, the nucleic acid or gene is a Bacillus gene homologue (e.g., is derived from a species distinct from Bacillus but having significant homology to a Bacillus gene of the present invention, for example, a Bacillus pan gene or Bacillus ilv gene).

Included within the scope of the present invention are bacterial-derived nucleic acid molecules or genes and/or Bacillus-derived nucleic acid molecules or genes (e.g., B. subtilis-derived nucleic acid molecules or genes), for example, the genes identified by 10 the present inventors, for example, Bacillus or B. subtilis coaX genes, coaA genes, pan genes and/or ilv genes. Further included within the scope of the present invention are bacterial-derived nucleic acid molecules or genes and/or Bacillus-derived nucleic acid molecules or genes (e.g., B. subtilis-derived nucleic acid molecules or genes) (e.g., B. subtilis nucleic acid molecules or genes) which differ from naturally-occurring bacterial 15 and/or Bacillus nucleic acid molecules or genes (e.g., B. subtilis nucleic acid molecules or genes), for example, nucleic acid molecules or genes which have nucleic acids that are substituted, inserted or deleted, but which encode proteins substantially similar to the naturally-occurring gene products of the present invention. In one embodiment, an isolated nucleic acid molecule comprises at least one of the nucleotide sequences set 20 forth as SEO ID NO:23, SEO ID NO:25, SEO ID NO:27, SEO ID NO:29, SEO ID NO 31, SEO ID NO:33, SEO ID NO:86, SEQ ID NO:35 or SEQ ID NO:37. In another preferred embodiment, an isolated nucleic acid molecule comprises at least two, three or four of the nucleotide sequences set forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEO ID NO:29, SEO ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID 25 NO:35 or SEO ID NO:37. For example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequences of SEQ ID NO:23, SEQ ID NO:25 and SEQ ID NO:27, preferably linked such that the proteins encoded by the nucleotide sequences of SEQ ID NO:23, SEQ ID NO:25 and SEQ ID NO:27 are each produced when the isolated nucleic acid molecule is expressed in a microorganism (e.g., SEQ ID NO:59). In another example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequences of SEQ ID NO:31 and SEQ ID NO:33, preferably linked such that the proteins encoded by the nucleotide sequences of SEQ ID NO:31 and SEQ ID NO:33 are each produced when the isolated nucleic acid molecule is expressed in a microorganism (e.g., nucleotides 1-2246 of SEQ ID NO:58). In another example, a preferred isolated nucleic acid molecule of the present invention can include the nucleotide sequence of SEQ ID NO:86. In another example, a preferred

35

isolated nucleic acid molecule of the present invention can include the nucleotide sequences of SEQ ID NO:31, SEQ ID NO:33 and SEQ ID NO:35, preferably linked such that the proteins encoded by the nucleotide sequences of SEQ ID NO:31, SEQ ID NO:33 and SEQ ID NO:35 are each produced when the isolated nucleic acid molecule is expressed in a microorganism (e.g., SEQ ID NO:58).

In another embodiment, an isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which is at least about 60-65%, preferably at least about 70-75%, more preferable at least about 80-85%, and even more preferably at least about 90-95% or more identical to a nucleotide sequence set forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37. In another embodiment, an isolated nucleic acid molecule hybridizes under stringent conditions to a nucleic acid molecule having a nucleotide sequence set forth as SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37. Such stringent conditions are known to those skilled in 15 the art and can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent (e.g. high stringency) hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that 20 hybridizes under stringent conditions to the sequence of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature. 25

A nucleic acid molecule of the present invention (e.g., a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37 can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, nucleic acid molecules can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) or can be isolated by the polymerase chain reaction using synthetic oligonucleotide primers designed based upon the sequence of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO 31, SEQ ID NO:33, SEQ ID NO:88, SEQ ID NO:35 or SEQ ID NO:37. A nucleic acid of the invention can be

amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:29, SEQ ID NO:33, SEQ ID NO:33, SEQ ID NO:35.

Additional panC nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:25, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:26 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:26 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:25 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:26, or are complementary to a panC nucleotide sequence as set forth herein.

Aditional panD nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:27, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:28 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:28 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:27 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:28, or are complementary to a panD nucleotide sequence as set forth herein.

Additional panE nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:29, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:30 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:30 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:29 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:30, or are complementary to a panE nucleotide sequence as set forth herein.

20

25

30

15

20

30

35

012177243 145

Additional *ilvB* nucleic acid sequences are those that comprise the nucleotide sequence of SEQ ID NO:31, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:32 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:32 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:31 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:32, or are complementary to an *ilvB* nucleotide sequence as set forth herein.

Additional *ilvN* nucleic acid sequences are those that comprise the nucleotide sequence of SEQ ID NO:33, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:34 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:34 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:33 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:34, or are complementary to an *ilvN* nucleotide sequence as set forth herein.

Additional *ilvC* nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:35, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:36 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:36 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:35 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:36, or are complementary to an *ilvC* nucleotide sequence as set forth herein.

Additional *ilvD* nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:37, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:38 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:38 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:37 or

10

15

20

25

30

to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:38, or are complementary to an *ilvD* nucleotide sequence as set forth herein.

Additional alsS nucleic acid sequences include those that comprise the nucleotide sequence of SEQ ID NO:86, encode a homologue of the polypeptide having the amino acid sequence set forth in SEQ ID NO:87 (e.g., encode a polypeptide having at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or more identity to the polypeptide having the amino acid sequence as set forth in SEQ ID NO:87 and a substantially identical activity as said polypeptide), hybridize under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:86 or to all or a portion of a nucleic acid molecule that encodes a polypeptide having the amino acid sequence of SEQ ID NO:87, or are complementary to an alsS nucleotide sequence as set forth herein.

In another embodiment, an isolated nucleic acid molecule is or includes a coaX gene, or portion or fragment thereof. In one embodiment, an isolated coaX nucleic acid molecule or gene comprises the nucleotide sequence as set forth in SEQ ID NO:19 (e.g., comprises the B. subtilis coaX nucleotide sequence). In another embodiment, an isolated coaX nucleic acid molecule or gene comprises a nucleotide sequence that encodes the amino acid sequence as set forth in SEQ ID NO:9 (e.g., encodes the B. subtilis CoaX amino acid sequence). In yet another embodiment, an isolated coaX nucleic acid molecule or gene encodes a homologue of the CoaX protein having the amino acid sequence of SEQ ID NO:9. As used herein, the term "homologue" includes a protein or polypeptide sharing at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identity with the amino acid sequence of a wild-type protein or polypeptide described herein and having a substantially equivalent functional or biological activity as said wild-type protein or polypeptide. For example, a CoaX homologue shares at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identity with the protein having the amino acid sequence set forth as SEQ ID NO:9 and has a substantially equivalent functional or biological activity (i.e., is a functional equivalent) of the protein having the amino acid sequence set forth as SEQ ID NO:9 (e.g., has a substantially equivalent pantothenate kinase activity). In a preferred embodiment, an isolated coaX nucleic acid molecule or gene comprises a nucleotide sequence that encodes a polypeptide as set forth in any one of SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID

NO:74 or SEQ ID NO:75. In another embodiment, an isolated coaX nucleic acid molecule hybridizes to all or a portion of a nucleic acid molecule having the nucleotide sequence set forth in SEQ ID NO:19 or hybridizes to all or a portion of a nucleic acid molecule having a nucleotide sequence that encodes a polypeptide having the amino 5 acid sequence of any of SEQ ID NOs:7-18, 74 or 75. Such hybridization conditions are known to those skilled in the art and can be found in Current Protocols in Molecular Biology, Ausubel et al., eds., John Wiley & Sons, Inc. (1995), sections 2, 4 and 6. Additional stringent conditions can be found in Molecular Cloning: A Laboratory Manual, Sambrook et al., Cold Spring Harbor Press, Cold Spring Harbor, NY (1989), chapters 7, 9 and 11. A preferred, non-limiting example of stringent hybridization 10 conditions includes hybridization in 4X sodium chloride/sodium citrate (SSC), at about 65-70°C (or hybridization in 4X SSC plus 50% formamide at about 42-50°C) followed by one or more washes in 1X SSC, at about 65-70°C. A preferred, non-limiting example of highly stringent hybridization conditions includes hybridization in 1X SSC, at about 65-70°C (or hybridization in 1X SSC plus 50% formamide at about 42-50°C) followed by one or more washes in 0.3X SSC, at about 65-70°C. A preferred, nonlimiting example of reduced stringency hybridization conditions includes hybridization in 4X SSC, at about 50-60°C (or alternatively hybridization in 6X SSC plus 50% formamide at about 40-45°C) followed by one or more washes in 2X SSC, at about 50-60°C. Ranges intermediate to the above-recited values, e.g., at 65-70°C or at 42-50°C 20 are also intended to be encompassed by the present invention. SSPE (1X SSPE is 0.15 M NaCl, 10mM NaH₂PO₄, and 1.25 mM EDTA, pH 7.4) can be substituted for SSC (1X SSC is 0.15 M NaCl and 15 mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes each after hybridization is complete. The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length 25 should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, $T_m(^{\circ}C) = 2(\# \text{ of } A + T \text{ bases}) + 4(\# \text{ of } G + C \text{ bases})$. For hybrids between 18 and 49 base pairs in length, $T_m(^{\circ}C) = 81.5 + 16.6(\log_{10}[Na^{+}]) + 0.41(\%G+C) - (600/N)$, where N is the number of bases in the hybrid, and [Na⁺] is the concentration of sodium ions in the hybridization buffer ([Na⁺] for 1X SSC = 0.165 M). It will also be recognized by the skilled practitioner that additional reagents may be added to hybridization and/or wash buffers to decrease non-specific hybridization of nucleic acid molecules to membranes, for example, nitrocellulose or nylon membranes, including but not limited to blocking agents (e.g., BSA or salmon or herring sperm carrier DNA), detergents (e.g., SDS), chelating agents (e.g., EDTA), Ficoll, PVP and the like. When using nylon membranes, in particular, an additional preferred, non-limiting example of

stringent hybridization conditions is hybridization in 0.25-0.5M NaH₂PO₄, 7% SDS at about 65°C, followed by one or more washes at 0.02M NaH₂PO₄, 1% SDS at 65°C, see e.g., Church and Gilbert (1984) Proc. Natl. Acad. Sci. USA 81:1991-1995, (or, alternatively, 0.2X SSC, 1% SDS). In another preferred embodiment, an isolated nucleic acid molecule comprises a nucleotide sequence that is complementary to a coaX nucleotide sequence as set forth herein (e.g., is the full complement of the nucleotide sequence set forth as SEQ ID NO:19).

In another preferred embodiment, an isolated nucleic acid molecule is or includes a coaA gene, for example, a Bacillus (e.g., B. subtilis) coaA gene, or portion or fragment thereof. Exemplary isolated coal nucleic acid molecules and/or genes include (1) an isolated coal nucleic acid molecule or gene comprising the nucleotide sequence as set forth in any one of SEQ ID NOs:20-22; (2) an isolated coal nucleic acid molecule or gene comprising a nucleotide sequence that encodes the amino acid sequence as set forth in SEQ ID NO:3; (3) an isolated coaA nucleic acid molecule or gene comprising a nucleotide sequence which encodes a CoaA homologue (e.g., a polypeptide having an amino acid sequence at least about 30-35%, preferably at least about 35-40%, more preferably at least about 40-50%, and even more preferably at least about 60%, 70%, 80%, 90% or more identical to the amino acid sequence set forth as SEQ ID NO:3 and having a substantially equivalent enzymatic activity; (4) an isolated coaA nucleic acid molecule or gene comprising a nucleotide sequence that encodes a polypeptide as set forth in any one of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6; (5) an isolated nucleic acid molecule that hybridizes under stringent conditions to all or a portion of a nucleic acid molecule having the nucleotide sequence set forth in SEQ ID NO:20, SEQ ID NO:21 or SEQ ID NO:22 or hybridizes to all or a portion of a nucleic acid molecule having a nucleotide sequence that encodes a polypeptide having the amino acid sequence of SEQ ID NO:3; and (6) an isolated nucleic acid molecule comprising a nucleotide sequence that is complementary to a coal nucleotide sequence as set forth herein (e.g., is the full complement of the nucleotide sequence set forth in SEQ ID NO:20, SEQ ID NO:21 or SEQ ID NO:22).

A nucleic acid molecule of the present invention (e.g., a coaX nucleic acid molecule or gene or a coaA nucleic acid molecule or gene), can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, nucleic acid molecules can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) or can be isolated by the polymerase chain reaction using synthetic oligonucleotide primers designed

25

30

PCT/US00/25993

based upon the coaX or coaA nucleotide sequences set forth herein, or flanking sequences thereof. A nucleic acid of the invention (e.g., a coaX nucleic acid molecule or gene or a coaA nucleic acid molecule or gene), can be amplified using cDNA, mRNA or alternatively, chromosomal DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques.

Yet another embodiment of the present invention features mutant coaX and coaA nucleic acid molecules or genes. The phrase "mutant nucleic acid molecule" or "mutant gene" as used herein, includes a nucleic acid molecule or gene having a nucleotide sequence which includes at least one alteration (e.g., substitution, insertion, deletion) such that the polypeptide or protein that may be encoded by said mutant exhibits an activity that differs from the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene. Preferably, a mutant nucleic acid molecule or mutant gene (e.g., a mutant coaA or coaX gene) encodes a polypeptide or protein having a reduced activity (e.g., having a reduced pantothenate kinase activity) as compared to the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene, for example, when assayed under similar conditions (e.g., assayed in microorganisms cultured at the same temperature). A mutant gene also can encode no polypeptide or have a reduced level of production of the wild-type polypeptide.

As used herein, a "reduced activity" or "reduced enzymatic activity" is one that is at least 5% less than that of the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene, preferably at least 5-10% less, more preferably at least 10-25% less and even more preferably at least 25-50%, 50-75% or 75-100% less than that of the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene. Ranges intermediate to the above-recited values, e.g., 75-85%, 85-90%, 90-95%, are also intended to be encompassed by the present invention. As used herein, a "reduced activity" or "reduced enzymatic activity" also includes an activity that has been deleted or "knocked out" (e.g., approximately 100% less activity than that of the polypeptide or protein encoded by the wild-type nucleic acid molecule or gene). Activity can be determined according to any well accepted assay for measuring activity of a particular protein of interest. Activity can be measured or assayed directly, for example, measuring an activity of a protein isolated or purified from a cell. Alternatively, an activity can be measured or assayed within a cell or in an extracellular medium. For example, assaying for a mutant coal gene or a mutant coal gene (i.e., said mutant encoding a reduced pantothenate kinase activity) can be accomplished by expressing the mutated gene in a microorganism, for example, a mutant microorganism which expresses pantothenate kinase in a temperature-sensitive manner, assaying the mutant gene for the ability to complement a temperature sensitive (Ts) mutant for

5

10

20

10

15

20

25

35

pantothenate kinase activity. A coaX mutant gene or coaA mutant gene that encodes a "reduced pantothenate kinase activity" is one that complements the Ts mutant less effectively than, for example, a corresponding wild-type coaX gene or coaA gene.

It will be appreciated by the skilled artisan that even a single substitution in a nucleic acid or gene sequence (e.g., a base substitution that encodes an amino acid change in the corresponding amino acid sequence) can dramatically affect the activity of an encoded polypeptide or protein as compared to the corresponding wild-type polypeptide or protein. A mutant nucleic acid or mutant gene (e.g., encoding a mutant polypeptide or protein), as defined herein, is readily distinguishable from a nucleic acid or gene encoding a protein homologue, as described above, in that a mutant nucleic acid or mutant gene encodes a protein or polypeptide having an altered activity, optionally observable as a different or distinct phenotype in a microorganism expressing said mutant gene or nucleic acid or producing said mutant protein or polypeptide (i.e., a mutant microorganism) as compared to a corresponding microorganism expressing the wild-type gene or nucleic acid or producing said mutant protein or polypeptide. By contrast, a protein homologue has an identical or substantially similar activity, optionally phenotypically indiscernable when produced in a microorganism, as compared to a corresponding microorganism expressing the wild-type gene or nucleic acid. Accordingly it is not, for example, the degree of sequence identity between nucleic acid molecules, genes, protein or polypeptides that serves to distinguish between homologues and mutants, rather it is the activity of the encoded protein or polypeptide that distinguishes between homologues and mutants: homologues having, for example, low (e.g., 30-50% sequence identity) sequence identity yet having substantially equivalent functional activities, and mutants, for example sharing 99% sequence identity yet having dramatically different or altered functional activities. Exemplary homologues are set forth in Figure 20 (i.e., CoaA homologues) and in Figure 23 (i.e., CoaX homologues). Exemplary mutants are described in Examples XV and XVIII herein.

30 VIII. Recombinant Nucleic Acid Molecules and Vectors

The present invention further features recombinant nucleic acid molecules (e.g., recombinant DNA molecules) that include nucleic acid molecules and/or genes described herein (e.g., isolated nucleic acid molecules and/or genes), preferably Bacillus genes, more preferably Bacillus subtilis genes, even more preferably Bacillus subtilis pantothenate kinase genes (e.g., coaX genes or coaA genes), pantothenate biosynthetic genes (e.g., genes encoding pantothenate biosynthetic enzymes, for example, panB genes encoding ketopantoate hydroxymethyltransferase, panE genes encoding

20

25

30

35

0101770A3 IA

ketopantoate reductase, panC genes encoding pantothenate synthetase, and/or panD genes encoding aspartate-α-decarboxylase) and/or isoleucine-valine (ilv) biosynthetic genes (e.g., ilvBN or alsS genes encoding acetohydroxyacid synthetase, ilvC genes encoding acetohydroxyacid isomeroreductase and/or ilvD genes encoding dihydroxyacid dehydratase).

The present invention further features vectors (e.g., recombinant vectors) that include nucleic acid molecules (e.g., isolated or recombinant nucleic acid molecules and/or genes) described herein. In particular, recombinant vectors are featured that include nucleic acid sequences that encode bacterial gene products as described herein, preferably Bacillus gene products, more preferably Bacillus subtilis gene products, even more preferably Bacillus subtilis pantothenate biosynthetic gene products (e.g., pantothenate biosynthetic enzymes, for example, ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase, and/or aspartate-α-decarboxylase) and/or isoleucine-valine biosynthetic gene products (e.g., 15 acetohydroxyacid synthetase, acetohydroxyacid isomeroreductase and/or dihydroxyacid dehydratase).

The term "recombinant nucleic acid molecule" includes a nucleic acid molecule (e.g., a DNA molecule) that has been altered, modified or engineered such that it differs . in nucleotide sequence from the native or natural nucleic acid molecule from which the recombinant nucleic acid molecule was derived (e.g., by addition, deletion or substitution of one or more nucleotides). Preferably, a recombinant nucleic acid molecule (e.g., a recombinant DNA molecule) includes an isolated nucleic acid molecule or gene of the present invention (e.g., an isolated coaX, coaA, pan or ilv gene) operably linked to regulatory sequences.

The term "recombinant vector" includes a vector (e.g., plasmid, phage, phasmid, virus, cosmid or other purified nucleic acid vector) that has been altered, modified or engineered such that it contains greater, fewer or different nucleic acid sequences than those included in the native or natural nucleic acid molecule from which the recombinant vector was derived. Preferably, the recombinant vector includes a coaX, coaA, pan or ilv gene or recombinant nucleic acid molecule including such coaX, coaA, pan or ilv gene, operably linked to regulatory sequences, for example, promoter sequences, terminator sequences and/or artificial ribosome binding sites (RBSs), as defined herein.

The phrase "operably linked to regulatory sequence(s)" means that the nucleotide sequence of the nucleic acid molecule or gene of interest is linked to the regulatory sequence(s) in a manner which allows for expression (e.g., enhanced, increased, constitutive, basal, attenuated, decreased or repressed expression) of the

10

15

20

25

30

35

nucleotide sequence, preferably expression of a gene product encoded by the nucleotide sequence (e.g., when the recombinant nucleic acid molecule is included in a recombinant vector, as defined herein, and is introduced into a microorganism).

The term "regulatory sequence" includes nucleic acid sequences which affect (e.g., modulate or regulate) expression of other nucleic acid sequences. In one embodiment, a regulatory sequence is included in a recombinant nucleic acid molecule or recombinant vector in a similar or identical position and/or orientation relative to a particular gene of interest as is observed for the regulatory sequence and gene of interest as it appears in nature, e.g., in a native position and/or orientation. For example, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence which accompanies or is adjacent to the gene of interest in the natural organism (e.g., operably linked to "native" regulatory sequences, for example, to the "native" promoter). Alternatively, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence which accompanies or is adjacent to another (e.g., a different) gene in the natural organism. Alternatively, a gene of interest can be included in a recombinant nucleic acid molecule or recombinant vector operably linked to a regulatory sequence from another organism. For example, regulatory sequences from other microbes (e.g., other bacterial regulatory sequences, bacteriophage regulatory sequences and the like) can be operably linked to a particular gene of interest.

In one embodiment, a regulatory sequence is a non-native or non-naturallyoccurring sequence (e.g., a sequence which has been modified, mutated, substituted, derivatized, deleted including sequences which are chemically synthesized). Preferred regulatory sequences include promoters, enhancers, termination signals, anti-termination signals and other expression control elements (e.g., sequences to which repressors or inducers bind and/or binding sites for transcriptional and/or translational regulatory proteins, for example, in the transcribed mRNA). Such regulatory sequences are described, for example, in Sambrook, J., Fritsh, E. F., and Maniatis, T. Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989. Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in a microorganism (e.g., constitutive promoters and strong constitutive promoters), those which direct inducible expression of a nucleotide sequence in a microorganism (e.g., inducible promoters, for example, xylose inducible promoters) and those which attenuate or repress expression of a nucleotide sequence in a microorganism (e.g., attenuation signals or repressor sequences). It is also within the scope of the present invention to regulate expression of a gene of interest by removing or deleting regulatory sequences. For

35

example, sequences involved in the negative regulation of transcription can be removed such that expression of a gene of interest is enhanced.

In one embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes a nucleic acid sequence or gene that encodes at least one bacterial gene product (e.g., a pantothenate biosynthetic enzyme, an isoleucinevaline biosynthetic enzyme, or a CoaA biosynthetic enzyme, for example CoaA or CoaX) operably linked to a promoter or promoter sequence. Preferred promoters of the present invention include Bacillus promoters and/or bacteriophage promoters (e.g., bacteriophage which infect Bacillus). In one embodiment, a promoter is a Bacillus promoter, preferably a strong Bacillus promoter (e.g., a promoter associated with a biochemical housekeeping gene in Bacillus or a promoter associated with a glycolytic pathway gene in Bacillus). In another embodiment, a promoter is a bacteriophage promoter. In a preferred embodiment, the promoter is from the bacteriophage SP01. In a particularly preferred embodiment, a promoter is selected from the group consisting of 15 P_{15} , P_{26} or P_{veg} , for example, the promoters set forth in SEQ ID NO:39, SEQ ID NO:40 or SEQ ID NO:41. Additional preferred promoters include tef (the translational elongation factor (TEF) promoter) and pyc (the pyruvate carboxylase (PYC) promoter), which promote high level expression in Bacillus (e.g., Bacillus subtilis). Additional preferred promoters, for example, for use in Gram positive microorganisms include, but are not limited to, the amyE promoter or phage SP02 promoters. Additional preferred 20 promoters, for example, for use in Gram negative microorganisms include, but are not limited to tac, trp, tet, trp-tet, lpp, lac, lpp-lac, lacIq, T7, T5, T3, gal, trc, ara, SP6, λ-P_R or λ - P_L .

In another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes a terminator sequence or terminator sequences (e.g., transcription terminator sequences). The term "terminator sequences" includes regulatory sequences which serve to terminate transcription of a gene. Terminator sequences (or tandem transcription terminators) can further serve to stabilize mRNA (e.g., by adding structure to mRNA), for example, against nucleases.

In yet another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes sequences which allow for detection of the vector containing said sequences (i.e., detectable and/or selectable markers), for example, sequences that overcome auxotrophic mutations, for example, ura3 or ilvE, fluorescent markers, and/or colorimetric markers (e.g., lacZ/β-galactosidase), and/or antibiotic resistance genes (e.g., amp or tet).

30

In yet another embodiment, a recombinant nucleic acid molecule or recombinant vector of the present invention includes an artificial ribosome binding site (RBS). The term "artificial ribosome binding site (RBS)" includes a site within an mRNA molecule (e.g., coded within DNA) to which a ribosome binds (e.g., to initiate translation) which differs from a native RBS (e.g., a RBS found in a naturally-occurring gene) by at least one nucleotide. Preferred artificial RBSs include about 5-6, 7-8, 9-10, 11-12, 13-14, 15-16, 17-18, 19-20, 21-22, 23-24, 25-26, 27-28, 29-30 or more nucleotides of which about 1-2, 3-4, 5-6, 7-8, 9-10, 11-12, 13-15 or more differ from the native RBS (e.g., the native RBS of a gene of interest). Preferably, nucleotides which differ are substituted such that they are identical to one or more nucleotides of an ideal RBS (e.g., SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47 or SEQ ID NO:48), when optimally aligned for comparisons. Artificial RBSs can be used to replace the naturallyoccurring or native RBS associated with a particular gene. Artificial RBSs preferably increase translation of a particular gene. Preferred artificial RBSs (e.g., RBSs for increasing the translation of panB, for example, of B. subtilis panB) are depicted in Table IA (e.g., SEQ ID NO:49 and SEQ ID NO:50).

Table 1A: Preferred panB Ribosome Binding Sites

```
20
10 20

-----AGAAAGGAGGTGA ideal RBS (SEQ ID NO:44)

CCCTCT-AG-AAGGAGGAGAAAACATG RBS1 (SEQ ID NO:49)

CCCTCT-AG--AGGAGGAGAAAACATG RBS2 (SEQ ID NO:50)

25 TAAACAT-G--AGGAGGAGAAAACATG panB native RBS (SEQ ID NO:42)
```

Additional preferred artificial RBSs (e.g., RBSs for increasing the translation of panD, for example, of B. subtilis panD) are depicted in Table 1B (e.g., SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53 and SEQ ID NO:54).

Table 1B: Preferred panD Ribosome Binding Sites

	10 20			
	1 1	·		
35	CTAGAAAAGGAGGAATTTAAATG	pAN423 RBS (SEQ ID NO:88)		
	TTAAGAAAGGAGGTGANNNNATG	ideal RBS (SEQ ID NO:45)		
	TTAGAAA <u>GGAGG</u> ATTTAAATATG	new design A (SEQ ID NO:51)		
	TTAGAAAGGAGGTTTAATTAATG	new design B (SEQ ID NO:52)		
40	TTAGAAAGGAGGTGATTTAAATG	new design C1 (SEQ ID NO:53)		
	TTAGAAAGGAGGTGTTTAAAATG	new design C2 (SEQ ID NO:54)		
	TTAGAAAGGAGGTGANNNNATG	ideal RBS (SEQ ID NO:46)		

Additional preferred artificial RBSs (e.g., RBSs for increasing the translation of panD, for example, of B. subtilis panD) are depicted in Table 1C (e.g., SEQ ID NO:55, SEQ ID NO:56 and SEQ ID NO:57). The predicted amino acid sequence at the C-terminus of the PanC protein is shown. The start codon for PanD translation is underlined.

Table 1C: Additional Preferred panD Ribosome Binding Sites

			10		20	1					
10			1								
		A	GAA	AGG	AGG	TGA	NNN	NNN	N	ATG	ideal RBS (SEQ ID NO:47)
										ATG	native panD RBS (SEQ ID NO:43)
	Ile	Arg	Glu	Met	Glu	Arg	ITe	*		Met	SEQ ID NO:89
15		A	GAA	AGG	AGG	TGA	NNN	NNN	N	ATG	ideal RBS (SEQ ID NO:47)
	עייייע	CGA	GAA	AGG	AGG	TGA	ATA	TAA	т	ATG	NDI (SEQ ID NO:55)
20			Glu							Met	SEQ ID NO:90
20	тта	CGA	GAA	AGG	AGG	TGA	ATA	ATA	_	ATG	NDII (SEQ ID NO:56)
			Glu							Met	SEQ ID NO:90
								m = =		» III C	NDIII (SEQ ID NO:57)
									. 1	ATG	
25	Ile	Arg	Arg	Lys	Glu	Val	Asn	. *		Met	SEQ ID NO:91
			AGA	AAG	GAG	GTG	ANN	NNN	N	ATG	ideal RBS (SEQ ID NO:48)

Accordingly, in one embodiment, a vector of the present invention includes an artificial RBS as set forth in SEQ ID NO:49 or SEQ ID NO:50. In another embodiment, a vector of the present invention includes an artificial RBS as set forth in SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53 or SEQ ID NO:54. In yet another embodiment, a vector of the present invention includes an artificial RBS as set forth in SEQ ID NO:55, SEQ ID NO:56 or SEQ ID NO:57.

In another embodiment, a recombinant vector of the present invention includes sequences that enhance replication in bacteria (e.g., replication-enhancing sequences). In one embodiment, replication-enhancing sequences are derived from E. coli. In another embodiment, replication-enhancing sequences are derived from pBR322 (e.g., sequences included within the pBR322 derived portion of any of the pAN vectors as set forth in the Figures, i.e., the Not I-Not I sequences from about 5.0 kB to 9.0 kB of the vector depicted in Figure 3A).

15

20

30

35

In yet another embodiment, a recombinant vector of the present invention includes antibiotic resistance genes. The term "antibiotic resistance genes" includes sequences which promote or confer resistance to antibiotics on the host organism (e.g., Bacillus). In one embodiment, the antibiotic resistance genes are selected from the group consisting of cat (chloramphenicol resistance) genes, tet (tetracycline resistance) genes, erm (erythromycin resistance) genes, neo (neomycin resistance) genes and spec (spectinomycin resistance) genes. Recombinant vectors of the present invention can further include homologous recombination sequences (e.g., sequences designed to allow recombination of the gene of interest into the chromosome of the host organism). For example, amyE sequences can be used as homology targets for recombination into the host chromosome.

Preferred vectors of the present invention include, but are not limited to, vectors set forth in Figures 2-15, 17, 19, 22, 25 and 26. It will further be appreciated by one of skill in the art that the design of a vector can be tailored depending on such factors as the choice of microorganism to be genetically engineered, the level of expression of gene product desired and the like.

IX. Isolated Proteins

Another aspect of the present invention features isolated proteins (e.g., isolated pantothenate biosynthetic enzymes and/or valine-isoleucine biosynthetic enzymes and/or isolated CoA biosynthetic enzymes, for example isolated CoaA or CoaX). In one embodiment, proteins (e.g., isolated pantothenate biosynthetic enzymes and/or valine-isoleucine biosynthetic enzymes and/or isolated CoaA biosynthetic enzymes, for example isolated CoaA or CoaX) are produced by recombinant DNA techniques and can be isolated from microorganisms of the present invention by an appropriate purification scheme using standard protein purification techniques. In another embodiment, proteins are synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein (e.g., an isolated or purified biosynthetic enzyme) is substantially free of cellular material or other contaminating proteins from the microorganism from which the protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. In one embodiment, an isolated or purified protein has less than about 30% (by dry weight) of contaminating protein or chemicals, more preferably less than about 20% of contaminating protein or chemicals, still more preferably less than about 10% of contaminating protein or chemicals, and most preferably less than about 5% contaminating protein or chemicals.

3.7

35

In a preferred embodiment, the protein or gene product is derived from Bacillus (e.g., is Bacillus-derived). The term "derived from Bacillus" or "Bacillus-derived" includes a protein or gene product which is encoded by a Bacillus gene. Preferably, the gene product is derived from a microorganism selected from the group consisting of Bacillus subtilis, Bacillus lentimorbus, Bacillus lentus, Bacillus firmus, Bacillus pantothenticus, Bacillus amyloliquefaciens, Bacillus cereus, Bacillus circulans, Bacillus coagulans, Bacillus licheniformis, Bacillus megaterium, Bacillus pumilus, Bacillus thuringiensis, and other Group 1 Bacillus species, for example, as characterized by 16S rRNA type (Priest, supra). In another preferred embodiment, the protein or gene product is derived from Bacillus brevis or Bacillus stearothermophilus. In another 10 preferred embodiment, the protein or gene product is derived from a microorganism selected from the group consisting of Bacillus licheniformis, Bacillus amyloliquefaciens. Bacillus halodurans, Bacillus subtilis, and Bacillus pumilus. In a particularly preferred embodiment, the protein or gene product is derived from Bacillus subtilis (e.g., is Bacillus subtilis-derived). The term "derived from Bacillus subtilis" or "Bacillus 15 subtilis-derived" includes a protein or gene product which is encoded by a Bacillus subtilis gene. In yet another preferred embodiment, the protein or gene product is encoded by a Bacillus gene homologue (e.g., a gene derived from a species distinct from Bacillus but having significant homology to a Bacillus gene of the present invention, for example, a Bacillus pan gene or Bacillus ilv gene). 20

Included within the scope of the present invention are bacterial-derived proteins or gene products and/or Bacillus-derived proteins or gene products (e.g., B. subtilisderived gene products) that are encoded by naturally-occurring bacterial and/or Bacillus genes (e.g., B. subtilis genes), for example, the genes identified by the present inventors, for example, Bacillus or B. subtilis coaX genes, coaA genes, pan genes and/or ilv genes. Further included within the scope of the present invention are bacterial-derived proteins or gene products and/or Bacillus-derived proteins or gene products (e.g., B. subtilisderived gene products) that are encoded bacterial and/or Bacillus genes (e.g., B. subtilis genes) which differ from naturally-occurring bacterial and/or Bacillus genes (e.g., B. subtilis genes), for example, genes which have nucleic acids that are mutated, inserted or deleted, but which encode proteins substantially similar to the naturally-occurring gene products of the present invention. For example, it is well understood that one of skill in the art can mutate (e.g., substitute) nucleic acids which, due to the degeneracy of the genetic code, encode for an identical amino acid as that encoded by the naturallyoccurring gene. Moreover, it is well understood that one of skill in the art can mutate (e.g., substitute) nucleic acids which encode for conservative amino acid substitutions. It is further well understood that one of skill in the art can substitute, add or delete

amino acids to a certain degree without substantially affecting the function of a gene product as compared with a naturally-occurring gene product, each instance of which is intended to be included within the scope of the present invention.

In a preferred embodiment, an isolated protein of the present invention (e.g., an isolated pantothenate biosynthetic enzyme and/or an isolated isoleucine-valine biosynthetic enzyme and/or an isolated CoaA biosynthetic enzymes, for example isolated CoaA or CoaX) has an amino acid sequence shown in SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87. In other embodiments, an isolated protein of the present invention is a homologue of the at least one of the proteins set forth as SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87 (e.g., comprises an amino acid sequence at least about 30-40% identical, preferably about 40-50% identical, more preferably about 50-60% identical, and even more preferably about 60-70%, 70-80%, 80-90%, 90-95% or . 15 more identical to the amino acid sequence of SEQ ID NO:3, SEQ ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID , P., . NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87, and has an activity that is substantially similar to that of the protein encoded by the amino acid sequence of SEQ ID NO:3, SEO ID NO:9, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, SEQ ID NO:34, SEQ ID NO:36, SEQ ID NO:38 or SEQ ID NO:87, respectively.

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = # of identical positions/total # of positions x 100), preferably taking into account the number of gaps and size of said gaps necessary to produce an optimal alignment.

The comparison of sequences and determination of percent homology between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264-68, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-77. Such

an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul et al. (1990) J. Mol. Biol. 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997) Nucleic Acids Research 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See http://www.ncbi.nlm.nih.gov. Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller (1988) Comput Appl Biosci. 4:11-17. Such an algorithm is incorporated into the ALIGN program available, for example, at the GENESTREAM network server, IGH Montpellier, FRANCE (http://vega.igh.cnrs.fr) or at the ISREC server (http://www.ch.embnet.org). When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

In another preferred embodiment, the percent homology between two amino acid sequences can be determined using the GAP program in the GCG software package (available at http://www.gcg.com), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 12, 10, 8, 6, or 4 and a length weight of 2, 3, or 4. In yet another preferred embodiment, the percent homology between two nucleic acid sequences can be accomplished using the GAP program in the GCG software package (available at http://www.gcg.com), using a gap weight of 50 and a length weight of 3.

25

30

35

010177089 185

20

10

15

X. Biotransformations and Bioconversions

Another aspect of the present invention includes biotransformation processes which feature recombinant microorganisms (e.g., mutant microorganisms) and/or isolated CoA, pantothenate or isoleucine-valine biosynthetic enzymes described herein. The term "biotransformation process", also referred to herein as "bioconversion processes", includes biological processes which result in the production (e.g., transformation or conversion) of any compound (e.g., intermediate or product) which is upstream of a CoA, pantothenate or isoleucine-valine biosynthetic enzyme to a compound (e.g., substrate, intermediate or product) which is downstream of said CoA, pantothenate or isoleucine-valine biosynthetic enzyme.

10

15

20

25

30

35

In one embodiment, the invention features a biotransformation process for the production of a panto-compound comprising contacting a microorganism which overexpresses at least one pantothenate biosynthetic enzyme with at least one appropriate substrate or precursor under conditions such that said panto-compound is produced and recovering said panto-compound. In a preferred embodiment, the invention features a biotransformation process for the production of pantoate comprising contacting a microorganism which overexpresses ketopantoate reductase (the panE gene product) with an appropriate substrate (e.g., ketopantoate) under conditions such that pantoate is produced and recovering said pantoate. In another preferred embodiment, the invention features a biotransformation process for the production of pantothenate comprising contacting a microorganism which overexpresses ketopantoate reductase and pantothenate synthetase with appropriate substrates (e.g., ketopantoate and β-alanine) under conditions such that pantothenate is produced and recovering said pantothenate. In yet another preferred embodiment, the invention features a biotransformation process for the production of pantothenate comprising contacting a microorganism which overexpresses ketopantoate hydroxymethyltransferase, ketopantoate reductase and pantothenate synthetase with appropriate substrates (e.g., α-ketoisovalerate and βalanine) under conditions such that pantothenate is produced and recovering said pantothenate. Preferred recombinant microorganisms for carrying out the abovedescribed biotransformations include pantothenate kinase mutants. Conditions under which pantoate or pantothenate are produced can include any conditions which result in the desired production of pantoate or pantothenate, respectively.

In yet another embodiment, the present invention includes a method of producing β -alanine that includes culturing a microorganism which overexpresses aspartate- α -decarboxylase under conditions such that β -alanine is produced. Preferably, the aspartate- α -decarboxylase-overexpressing microorganism has a mutation in a nucleic acid sequence encoding a pantothenate biosynthetic enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase and pantothenate synthetase.

The invention further features a method of producing β -alanine that includes contacting a composition comprising aspartate with an isolated *Bacillus* aspartate- α -decarboxylase enzyme under conditions such that β -alanine is produced (e.g., an *in vitro* synthesis method).

The microorganism(s) and/or enzymes used in the biotransformation reactions are in a form allowing them to perform their intended function (e.g., producing a desired compound). The microorganisms can be whole cells, or can be only those portions of the cells necessary to obtain the desired end result. The microorganisms can be

15

20

25

30

35

suspended (e.g., in an appropriate solution such as buffered solutions or media), rinsed (e.g., rinsed free of media from culturing the microorganism), acetone-dried, immobilized (e.g., with polyacrylamide gel or k-carrageenan or on synthetic supports, for example, beads, matrices and the like), fixed, cross-linked or permeablized (e.g., have permeablized membranes and/or walls such that compounds, for example, substrates, intermediates or products can more easily pass through said membrane or wall).

Purified or unpurified CoA biosynthetic enzyme(s) (e.g., CoaA and/or CoaX), pantothenate biosynthetic enzyme(s) and/or valine-isoleucine biosynthetic enzyme(s) can also be used in biotransformation reactions. The enzyme can be in a form that allows it to perform its intended function (e.g., obtaining the desired compound). For example, the enzyme can be in free form or immobilized. Purified or unpurified CoA biosynthetic enzyme(s), pantothenate biosynthetic enzyme(s) and/or valine-isoleucine biosynthetic enzyme(s) can be contacted in one or more in vitro reactions with appropriate substrate(s) such that the desired product is produced.

With respect to at least the above-described methodologies (e.g., the production methodologies of the present invention), at least one aspect of the invention features the folowing: embodiments is which the methods do not use microorganisms of the genus Corynebacterium and/or microorganisms of the genus Escherichia; embodiments in which the methods do not use microorganisms selected from the group consiting of Escherichia coli and Corynebacterium glutamicum; embodiments in which the methods do not use gram negative microorganisms; embodiments in which the microorganisms utilized do not include, express or produce nucleic acid molecules, genes or proteins (e.g., biosynthetic emzymes) derived from microorganisms of the genus Corynebacterium and/or microorganisms of the genus Escherichia; embodiments in which the microorganisms to not include, express or produce nucleic acid molecules, genes or proteins (e.g., biosynthetic emzymes) derived from microorganisms selected from the group consisting of Escherichia coli and Corynebacterium glutamicum.

XI. Screening Assays

Because CoA is an essential factor in bacteria, proteins (e.g., enzymes) involved in the biosynthesis of CoA provide valuable tools in the search for novel anti-biotics. In particular, the CoaX protein is a valuable target for identifying bacteriocidal compounds because it bears no resemblance in primary sequence to mammalian pantothenate kinase enzymes. Accordingly, the present invention also provides a method (also referred to herein as a "screening assay") for identifying modulators, i.e., candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other drugs)

which bind to CoaX, or have a stimulatory or inhibitory effect on, for example, coaX expression or CoaX activity.

In one embodiment, the invention provides assays for screening candidate or test compounds which are capable of binding to CoaX proteins or a biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which modulate the activity of CoaX proteins or biologically active portions thereof. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) Anticancer Drug Des. 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt et al. (1993) Proc. Natl. Acad. Sci. U.S.A. 90:6909; Erb et al. (1994) Proc. Natl. Acad. Sci. USA 91:11422; Zuckermann et al. (1994). J. Med. Chem. 37:2678; Cho et al. (1993) Science 261:1303; Carrell et al. (1994) Angew. Chem. Int. Ed. Engl. 33:2059; Carell et al. (1994) Angew. Chem. Int. Ed. Engl. 33:2061; and in Gallop et al. (1994) J. Med. Chem. 37:1233. Libraries of compounds may be presented in solution (e.g., Houghten (1992) Biotechniques 13:412-421), or on beads (Lam (1991) Nature 354:82-84), chips (Fodor (1993) Nature 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull et al. (1992) Proc Natl Acad Sci USA 89:1865-1869) or on phage (Scott and Smith (1990) Science 249:386-390); (Devlin (1990) Science 249:404-406); (Cwirla et al. (1990) Proc. Natl. Acad. Sci. 87:6378-6382); (Felici (1991) J. Mol. Biol. 222:301-310); (Ladner supra.).

In one embodiment, an assay is a microorganism-based assay in which a recombinant microorganism which expresses a CoaX protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate CoaX activity is determined. Determining the ability of the test compound to modulate CoaX activity can be accomplished by monitoring, for example, intracellular phosphopanthoate or CoA concentrations or secreted pantothenate concentrations (as compounds that inhibit CoaX will result in a buildup of pantothenate in the test microorganism). CoaX substrate can be labeled with a radioisotope or enzymatic label such that modulation of CoaX activity can be determined by detecting a conversion of labeled substrate to intermediate or product. For example, CoaX substrates can be

labeled with ³²P, ¹⁴C, or ³H, either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Determining the ability of a compound to modulate CoaX activity can alternatively be determined by detecting the induction of a reporter gene (comprising a CoA-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a CoA-regulated cellular response.

In yet another embodiment, a screening assay of the present invention is a cell-free assay in which a CoaX protein or biologically active portion thereof is contacted with a test compound *in vitro* and the ability of the test compound to bind to or modulate the activity of the CoaX protein or biologically active portion thereof is determined. In a preferred embodiment, the assay includes contacting the CoaX protein or biologically active portion thereof with known substrates to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to modulate enzymatic activity of the CoaX on its substrates.

Screening assays can be accomplished in any vessel suitable for containing the microorganisms, proteins, and/or reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either CoaX protein or a recombinant microorganism expressing CoaX protein to facilitate separation of products and/or substrates, as well as to accommodate automation of the assay. For example, glutathione-S-transferase/CoaX fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates. Other techniques for immobilizing proteins on matrices (e.g., biotin-conjugation and streptavidin immobilization or antibody conjugation) can also be used in the screening assays of the invention.

In another embodiment, modulators of CoaX expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of coaX mRNA or CoaX polypeptide in the cell is determined. The level of expression in the presence of the candidate compound is compared to the level of expression in the absence of the candidate compound (or to a suitable control, for example, an appropriate buffer control or standard). The candidate compound can then be identified as a modulator of coaX mRNA or CoaX polypeptide expression based on this comparison.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an CoaX modulating agent identified as described herein (e.g., an anti-bactericidal

10

15

20

25

30

35

10

compound) can be used in an infectious animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents, patent applications (including U.S. Patent Application Serial No. 09/400,494, filed September 21, 1999 (pending), provisional U.S. Patent Application Serial No. 60/210,072, filed June 7, 2000, provisional U.S. Patent Application Serial No. 60/221,938, filed July 28, 2000 and provisional U.S. Patent Application Serial No. 60/227,860, filed August 24, 2000, to which this application relates) and published patent applications cited throughout this application are incorporated herein by reference.

15

EXAMPLES

General Methodology:

Strains. Bacillus subtilis strains of the present invention are generally derived from either of two strains. The first is variously named "168", "1A1", or "RL-1". The genotype is trpC2. This strain was derived from the wild type "Marburg" strain by mutagenesis and has been the basis of much of the molecular biology work done on B. subtilis. The second strain is PY79, a prototrophic derivative of 168 that was made Trp⁺ by transduction from the wild type strain W23.

Media. Standard minimal medium for B. subtilis is comprised of 1 x Spizizen salts and 0.5% glucose. Standard solid "rich medium" is Tryptone Blood Agar Broth (Difco), and standard liquid "rich medium" is VY, a mixture of veal infusion broth and yeast extract. For testing production of pantothenate in liquid test tube cultures, an enriched form of VY, called "Special VY" or "SVY" is used. For batch fermentations, SVYG and PFMG are used. The compositions of these media are given below.

VY, a rich liquid medium: 25 g Difco Veal Infusion Broth, 5 g Difco Yeast Extract, 1L water (autoclave).

TBAB, a rich solid medium: 33 g Difco Tryptone Blood Agar Broth, 1L water (autoclave).

MIN, a minimal medium: 100 ml 10 x Spizizen salts; 10 ml 50% glucose; 2 ml 10% arginine HCl*; 10 ml 0.8% tryptophan**; water to 1 liter. (*In some cases, arginine is omitted or replace by sodium glutamate at 0.04% final concentration. In general, B. subtilis grows faster in minimal medium when certain amino acids, such as arginine, glutamine, glutamate, or proline, are added as an auxiliary nitrogen source.) (**For strains that are tryptophan auxotrophs, tryptophan is routinely added to most minimal media.)

10 x Spizizen Salts: 174 g K₂HPO₄·3H₂O; 20 g (NH₄)₂SO₄; 60 g KH₂PO₄; 10 g Na₃Citrate·2H₂O; 2 g MgSO₄·7H₂O; water to 993 mls; then add 3.5 ml FeCl₃ solution and 3.5 ml Trace Elements solution.

FeCl₃Solution: 4 g FeCl₃·6H₂O; 197 g Na₃Citrate 2H₂O; water to 1 liter (filter 30 sterilize)

Trace Elements Solution: 0.15 g Na₂MoO₄·2H₂O; 2.5 g H₃BO₃; 0.7 g CoCl₂·6H₂O; 0.25 g CuSO₄·5H₂O; 1.6 g MnCl₂·4H₂O; 0.3 g ZnSO₄·7H₂O; water to 1 liter (filter sterilize).

SVY, Special VY, a supplemented* rich medium for testing pantothenate production in test tube cultures: 25 g Difco Veal Infusion Broth; 5 g Difco yeast extract; 5 g sodium glutamate; 2.7 g ammonium sulfate; 740 ml water (autoclave); add 200 ml 1 M potassium phosphate, pH 7.0; 60 ml 50% glucose. (*For testing pantothenate production in liquid SVY test tube cultures, Na α-ketoisovalerate and/or β-alanine can be added to a concentration of 5 g/L from filter-sterilized stocks.)

PFMG, a yeast extract based medium used in fermentors: 20 g Amberex 1003™ yeast extract; 5 g sodium glutamate, 2 g ammonium sulfate; 5 g tryptophan; 10 g KH₂PO4; 20 g K₂HPO₄·3H₂O; 1 g MgCl₂·6H₂O; 0.1 g CaCl₂·2H₂O; 1 g sodium citrate; 0.01 g FeSO₄·7H₂O; 1 ml trace elements solution; 20 g glucose; add water to 1 L. Glucose or other sugars are fed as needed. Feed solutions can contain minerals, defined or food grade nutrients.

PF, a chemically defined pantothenate free medium for testing pantothenate auxotrophy: 100 ml 10 x Spizizen Salts; 100 ml 1 x Difco Pantothenate Assay Medium; 15 10 ml 50% glucose; water to 1 liter.

For pantothenate auxotrophs, 1 mM Na pantothenate is added to both minimal and rich media, since there is generally not enough pantothenate in rich media to support *B. subtilis pan* mutants. Amino acids are at 100 mg per liter, when used.

Selection for antibiotic resistance is done with 5 mg/L chloramphenicol, 100 mg/L spectinomycin HCl, 15 mg/L tetracycline HCl, or 1 mg/L erythromycin plus 25 mg/L lincomycin.

Pantothenate Assays: Biological assay. The indicator organism, Lactobacillus plantarum, requires pantothenate for growth, and responds to low concentrations (μg/L).
Thus, using serial dilutions, a wide range of concentrations can be assayed.
Commercially available medium (e.g., Pantothenate Assay Medium (PAM), Difco), can be used. However, Difco PAM supplemented with pantothenate does not support growth to the same level as obtainable using a fresh-mixed version of Pantothenate Assay Medium (FM-PAM), made up of the individual components as specified by
Difco, which is accordingly, routinely used instead of the commercial product.

Before assaying *B. subtilis* culture supernatants, the *B. subtilis* cells must be either removed or killed. *B. subtilis* culture supernatants give approximately the same pantothenate titer when the supernatants are autoclaved as when they are sterile filtered. Accordingly, routine procedures involve autoclaving samples for 5 minutes prior to the biological assay.

35

20

20

30

Pantothenate Assays: HPLC assay. Pantothenic acid production is measured by HPLC with a detector wavelength of 197 nm and a reference at 450 nm. The procedure is a modification of one recommended by Hewlett-Packard for water soluble vitamins. Samples of culture broth are diluted into an equal volume of 60% acetronitrile (ACN), centrifuged and filtered. Typically a further 10-fold dilution before analysis brings the final dilution to 20-fold. Higher concentrations of product are diluted further. Compounds are separated on a C18 Phenomenex 5µ Aqua 250 x 4.6 mm column with 5% acetronitrile (ACN) in 50 mM Na phosphate buffer at pH 2.5. An ACN gradient from 5% to 95% washes the column between every sample. The area of the pantothenate peak is proportional to the concentration between 5 to 1000 mg/L. Other panto-compounds are also separated and quantitated by this method.

Amino Acid Analysis: HPLC assay. Amino acids present in the fermentation medium and throughout the fermentation are measured by HPLC with a detector wavelength of 338 nm and a reference at 390 nm. The procedure is a modification of 15 one recommended by Hewlett-Packard for amino acid analysis. Samples of culture broth are prepared identically as for the panto-compound analysis. Compounds are separated on a C18 Hypersil 5µ ODS 200 x 2.1 mm column. Solvent A is 20 mM Na acetate buffer at pH 7.2. Solvent B contains 40% ACN and 40% methanol. A gradient from 100% Solvent A to 100% Solvent B separates amino acids and washes the column between every sample.

Batch Fermentations. Pantothenate producing strains are grown in stirred tank fermentors, for example, in CF3000 Chemap 14 liter vessels with 10 liter working volumes. Computer control and data collection is by commercial software, for example, B. Braun Biotech MFCS software. Fermentations can be batch processes but are preferably sugar-limited, fed batch processes. Some media components (e.g. of SVYG and PMFG) are added to the fermentor and sterilized in place. Portions of the media are sterilized separately and added to the fermentors aseptically. This procedure is well known to those familiar with the art. Additional nitrogen sources in feeds are sterilized separately and added to the carbon source after cooling.

The initial sugar in the medium is consumed in approximately 6 hours. Afterwards, glucose or other sugars are fed with the possible addition of minerals, and defined or food grade nutrients. Alternatively, feeds are scheduled based on a consensus profile of nutritional requirements from samples taken from earlier fermentations.

15

After inoculation, agitation is set at a relatively low speed, e.g. 200 rpm. When the dissolved oxygen (pO2) falls to 30%, computer control automatically adjusts the agitation to maintain a dissolved oxygen concentration between 25 and 30% pO2.

5 EXAMPLE I: Enhanced Production of a Panto-Compound Using Bacteria Overexpressing panBCD Gene Products.

This Example describes the cloning of the *B. subtilis panBCD* operon and the generation of microorganisms overexpressing the *panBCD* gene products.

To clone the *B. subtilis panBCD* operon, a plasmid library of *B. subtilis* GP275 (a derivative of 168) genomic DNA was transformed in *E. coli* BM4062 (birA^{ts}), and temperature resistant clones were selected at 42°C. By comparison of restriction maps to the genome sequence, one particular clone was deduced to contain the *B. subtilis birA* gene and the adjacent panBCD genes. This plasmid was named pAN201.

To overexpress the panBCD operon and produce pantothenate, the native promoter of the panBCD operon was replaced by either of two strong, constitutive promoters derived from the B. subtilis bacteriophage SP01. These two promoters are named P_{26} and P_{15} . In addition, either of two artificial ribosome binding sites (RBSs) were used to replace the native panB RBS. These two artificial RBSs (set forth as SEQ ID NO:49 and SEQ ID NO:50) were predicted to increase translation of panBCD; their sequences are shown in Table 1A. Three such engineered panBCD expression cassettes were built into circular plasmids capable of replicating in E. coli. Other features of the plasmids include a strong rho-independent transcription terminator from the E. coli ribosomal RNA transcription unit, called T₁T₂, a Gram-positive chloramphenicol resistance gene (cat), derived from pC194, and a pair of NotI restriction sites at the junctions between the E. coli replicon and the segment intended for integration into B. subtilis. Three plasmids of this series, pAN004, pAN005, and pAN006 were constructed. pAN004 contains the P26 promoter, RBS1, and a low copy E. coli replicon. pAN005 contains the P_{15} promoter, which in our-experience is not as strong as P_{26} , RBS1, and the low copy replicon. pAN006 contains the P_{26} promoter, RBS2, and a medium copy replicon.

The three panBCD expression cassettes contained in the above-mentioned three plasmids were all ligated to a DNA fragment consisting of sequences that naturally occur immediately upstream from the native panB gene and integrated in single copy by homologous recombination into the panBCD locus of B. subtilis strains RL-1 and PY79, replacing the wild-type operon. This was accomplished in two steps. First a deletion-substitution that replaced about two thirds of the panB coding region with a Gram-

20

positive spectinomycin resistance gene (*spec*) was integrated at *panB* to yield Spec^{*}, pantothenate auxotrophs. These intermediate strains were than transformed with the *panBCD* expression cassettes of pAN004, pAN005, and pAN006 after ligating them to a DNA fragment containing chromosomal sequences just upstream of *panB*. Selection of the incoming cassette was for pantothenate prototrophy. The resulting strains were named PA221, PA222 and PA223 (from RL-1), and PA235, PA232 and PA233 (from PY79), respectively. An example of a plasmid that contains the joined upstream sequence that is in the integrated strain in PA221 is pAN240 (see Figure 2). The nucleotide sequence of pAN240 is set forth as SEQ ID NO:76.

Polymerase chain reaction using appropriate primers was used to verify the correct chromosomal structures of these engineered strains. When extracts of strain PA221 were examined by SDS-PAGE, two proteins were found to be overexpressed. One protein had an apparent molecular weight of 29,000 and the other protein appeared to be 39,000 daltons. The 29,000 dalton bands is presumably PanB (predicted molecular weight of 29,761). The larger protein band presumably represents PanC (predicted size 31,960 daltons).

The ability of these strains to produce pantothenate in test tube cultures was assessed as follows. Each strain was grown in SVY medium supplemented with 5 g/L α-ketoisovalerate (α-KIV) and 5 g/L β-alanine, to ensure that these precursors were not limiting. Culture supernatants were autoclaved and assayed using the bioassay. Relative to the parent strains, RL-1 and PY79, the engineered strains produced about 8-to 30-fold more pantothenate, attaining 1 g/L pantothenate in some cases.

Table 2. Production of pantothenate by engineered B. subtilis strains in liquid test tube cultures grown in SVY medium with 5 g/L α -KIV and 5 g/L β -alanine.

Expt.	Strain	Promoter	RBS at panB	[pantothenate] mg/L
1	RL-1	Native	Native	30
	PA221	P ₂₆ .	RBS1	990 . 790
	PA222	P_{ls}	RBS1	250 250
	PA223	P ₂₆	RBS2	790 790
2	PY79	Native	Native	40
,	PA235	P ₂₆	RBS1	930 860
	PA221	P ₂₆	RBS1	1100 1030

The P_{26} promoter was about 3- to 4-fold more effective than the P_{15} promoter, 5 while RBS1 and RBS2 were roughly equivalent. Plasmids such as pAN004, pAN005, pAN006 can also be recombined as circles into the B. subtilis wild type panBCD locus by Campbell-type (single crossover) integration, selecting for chloramphenicol resistance at 5 mg/L. Strains obtained in this fashion produce about the same amount of pantothenate as strains PA221, PA222, and PA223, respectively. pAN004 containing the P_{26} promoter, RBS1 and a low copy E. coli replicon, is depicted schematically in Figure 3A. The nucleotide sequence of plasmid pAN004 is set forth as SEQ ID NO:93. pAN006 containing the P_{26} promoter, RBS2 and a medium copy E. coli replicon, is depicted schematically in Figure 3B. The nucleotide sequence of plasmid pAN006 is set forth as SEQ ID NO:94. The nucleotide sequence of panBCD is set forth as SEQ ID NO:59 and the predicted amino acid sequences of PanB, PanC and PanD are set forth as SEQ ID NO:24, SEQ ID NO:26 and SEQ ID NO:28, respectively. Methods for manipulating Bacilli are described, for example, in Harwood, C.R. and Cutting, S.M. (editors), Molecular Biological Methods for Bacillus (1990) John Wiley & Sons, Ltd., Chichester, England, the content of which is incorporated herein by reference.

10

20

EXAMPLE II: Enhanced Production of a Panto-Compound Using Bacteria Overexpressing the panE1 Gene Product - Ketopantoate Reductase.

This Example describes the cloning of the B. subtilis panEl gene and the generation of microorganisms overexpressing the panE1 gene product.

Pan B. subtilis strains (e.g., B. subtilis mutants blocked in the synthesis of pantothenic acid) had previously been isolated, one of which was reported to be affected in ketopantoate reductase activity (Baigori et al. (1991) J. Bacteriol. 173:4240-4242). However, the mutations in these strains were incorrectly mapped to the purE-tre interval of the B. subtilis genetic map which does not contain the panE or panBCD genes. Furthermore as shown below, a panE mutant does not have a Pan phenotype as the ilvC gene product can substitute for the panE gene product in B. subtilis as in other bacterial strains such as E. coli. More recently, the S. typhimuruim panE gene has been located and determined to be allelic to apbA, a gene required for anaerobic purine biosynthesis (Frodyma et al. (1998) J. Biol. Chem. 273:5572-5576). E. coli carries a highly

homologous gene at the same map location. Identification of the panE genes in E. coli and S. typhimurium was complicated by the fact that the ilvC gene product, acetohydroxy acid isomeroreductase, is also capable of carrying out the ketopantoate reductase reaction. As a result, pantothenate auxotrophy is not obtained unless both panE and ilvC are mutated.

To identify the B. subtilis panEl gene, the B. subtilis genome was searched using the protein sequence of E. coli or S. typhimurium ApbA (PanE), and two open reading frames were identified having homology to ApbA, named ylbQ and ykpB. These genes were renamed panE1 and panE2, due to their proposed function in pantothenate biosynthesis. Both panE1 and panE2 were cloned as PCR products generated from RL-1 genomic DNA as a template. Both genes were disrupted by either a 25 spectinomycin resistance gene (spec) or a chloramphenicol resistance gene (cat). The interrupted genes were each integrated by double crossover into PY79 to give PA240 (ΔpanEl::spec) and PA241 (ΔpanE2::cat). Neither of these strains were pantothenate auxotrophs when tested on pantothenate-free (PF) plates, although PA240 containing ΔpanEl::spec grew slightly more slowly on TBAB without added pantothenate than 30 with a 1 mM pantothenate supplement. By comparison, a ΔpanB::spec strain does not produce single colonies on TBAB, presumably because B. subtilis has no active uptake system for pantothenate.

It was hypothesized that the B. subtilis gene, ilvC, could function for panE as had been shown for E. coli. Accordingly, the panEl and panE2 disruptions were introduced 35 into a strain, CU550, which is reported to be trpC2 ilvC4 leuC124. Both the single

10

panE1 and the double panE1, panE2 disruptants were pantothenate auxotrophs on PF medium.

Table 3. Phenotypes of various panE1 and panE2 mutants on rich and defined media.

Strain	Medium	Growth*:	+ pan
PY79	TBAB	+++	+++
	PF	++	++
PA240	TBAB spec	+	+++
	PF	++	++
PA241	TBAB cam	+++	+++ .
	PF	++	++
CU550	TBAB	+++	+++
	PF	++	++
PA256	TBAB spec	-	+++
	PF	-	++
PA258	TBAB spec, cam	_	+++
	PF	- ,	++

^{*}Each "+" represents about 1 mm of colony diameter after overnight at 37°C.

Thus, mutating both panE1 and ilvC results in pantothenate auxotrophy, while mutating only panE1 does not, similar to what has been reported for E.coli and S. typhimurium.

Next, the quantitative effect of panE1 and panE2 knockouts in a pantothenate overproducing strain (PA235 described herein) was examined. The panE1 and panE2 disruptions were introduced into PA235, either singly or together to produce PA245 (ΔpanE1::spec), PA248 (ΔpanE2::cat) and PA244 (ΔpanE1::cat, ΔpanE2::spec). The effect of each mutation on pantothenate production was then tested in liquid test tube cultures.

20

25

0101770A2 IA

Table 4. Pantothenate production by PA235 derivatives containing panE1 and panE2 disruptions.

[pan] mg/L	% of PA235
990	(100)
940	95
59	6
82	8
1060	106
1030	104
25	3
50	5
	990 940 59 82 1060 1030 25

Thus, deletion analysis indicated that the panE1 gene contributes to over 90% of the pantothenate production, while deletion of panE2 did not have a significant effect on pantothenate production. It is therefore concluded that panE1 accounts for most, but not necessarily all, of the ketopantoate reductase activity in B. subtilis. The rest of the ketopantoate reductase activity is predicted to be supplied by ilvC.

Having identified panE1 as an important gene for pantothenate production, increased panE1 expression was tested to determine whether it could enhance pantothenate production in strains such as PA221 or PA235. The panE1 coding sequence was installed downstream of the P26 promoter and RBS2 in a vector, pOTP61, designed to integrate and amplify at either the bpr locus (a non-essential protease gene) or at the locus of the cloned insert. The resulting plasmid, pAN236 (Figure 4) was transformed into PA221, selecting for resistance to tetracycline at 15 mg/L. The nucleotide sequence of pAN236 is set forth as SEQ ID NO:77. One transformant, named PA236 was chosen for further study.

PA236 was shown to overexpress a protein of about 31,000 daltons, which is close to the expected molecular weight of 33,290 daltons for *panE1* protein. Briefly, whole cell extracts were prepared from PY79, RL-1, PA221, PA221/pOTP61and PA236 (2 samples). Cell extracts were separated by gel electrophoresis and the gels were coomassie stained to visualize proteins. In cells engineered to overexpress *panE* (PA236-1 and PA236-2), a band was visible having an approximate molecular weight of ~31,000 daltons (as compared to molecular weight markers). Moreover, PA221 and PA236 expressed increased levels of a ~29,000 dalton band, corresponding to the *panB*

15

gene product, and a ~39,000 dalton band, presumably corresponding the panC gene product. Furthermore, E. coli transformed with pAN006 (Figure 3B)-expressed bands correlating to the panB and panC gene products and E. coli transfected with PAN236 expressed a ~31,000 dalton band corresponding to the panE gene product.

Next, PA236 was compared to PA221 carrying the empty vector pOTP61 for pantothenate production in liquid test tube cultures supplemented with 5 g/L β -alanine and 5 g/L α -KIV.

Table 5. Effect of overexpression of panE1 and panE2 on pantothenate production by engineered strains in liquid test tube cultures.

Strain	Additional Plasmid	Gene Overexpressed	[Pantothenate] mg/L
PA221	pOTP61	none	1,000
			940
PA236	pAN236	panE1	2,030
		•	2,050
PA238	pAN238	panE2	530
1	-		680

Overexpression of panE1 caused a two-fold increase in pantothenate production when compared to the parent strain (e.g., to slightly over 2 g/L) whereas overexpression of panE2 resulted in a strain that produced about 35% less pantothenate than the parent strain. The panE1 nucleotide sequence and predicted amino acid sequence are set forth as SEQ ID NO:29 and SEQ ID NO:30.

EXAMPLE III: Enhanced Production of a Panto-Compound by Culturing Bacteria Overexpressing panE1 or panBCD in the Presence of Valine.

The ability of valine to function as a media supplement (e.g., as a substitute for α-KIV) in strains engineered to overexpress the panBCD operon and panE1 was evaluated. Valine is closely related to α-KIV by transamination, is less expensive than α-KIV, and is commercially available in kilogram quantities. Valine was substituted for α-KIV in the standard liquid test tube cultures in SVY medium. The concentration of valine was varied from 5 to 50 g/L. Although valine at 5 g/L was slightly less effective

than α -KIV in promoting pantothenate production, valine at 10 or 20 g/L equaled or surpassed 5 g/L α -KIV in promoting pantothenate production.

EXAMPLES IV-X Generation of Microorganisms Capable of Producing Pantothenate in a Precursor-Independent Manner

B. subtilis strains such as PA221 and PA235 (engineered to overexpress panBCD) and PA236 (engineered to overexpress panBCD and panE1) need to be fed α-ketoisovalerate (α-KIV) (or valine) and aspartate (or β-alanine) to achieve maximal pantothenate production, as both these precursors are limiting for pantothenate
synthesis. Accordingly, manipulated microorganisms were designed to eliminate the need to feed limiting precursors of pantothenate biosynthesis in the production of pantothenate. These strains are also useful in the production of various pantothenate biosynthetic pathway intermediates.

15 EXAMPLE IV: Generation of Microorganisms Capable of Producing Pantothenate in an Aspartate- (or β-Alanine) Independent Manner

The panD gene was cloned into B. subtilis expression vector pOTP61 to construct pAN423 (Figure 5). The nucleotide sequence of pAN423 is set forth as SEQ ID NO:78. The NotI restriction fragment containing panD was isolated from pAN423, self ligated and used to transform PA221. Transformants resistant to Tet¹⁵, Tet³⁰, and Tet⁶⁰ were isolated and saved for further analysis.

Six of the pAN423 transformants plus two control transformants were grown in SVY containing 5 g/l α -KIV with and without 10 g/l aspartate and then assayed for pantothenate production (Table 6).

Table 6. Effect of overproducing PanD on pantothenate production with and without added aspartate.

Culture* (PA221 transformants)	Asp (10 g/L)	TetR** (μg/ml)	OD550	[pan] (mg/L)
pOTP61-1	-	60	8.0	76
pOTP61-2	-	60	7.7	91
423#1-1	-	· 15	8.5	180
423#1-2	-	15	8.0	150
423#1-3	-	30	8.3	220
423#1-4	-	30	8.5	280
423#1-5	-	60	8.9	580
423#1-6	-	60	8.8	280

20

25

		1		
pOTP61-1	+	60	7.5	380
pOTP61-2	+	60	6.9	560
423#1-1	+	15	8.5	1200
423#1-2	+	15]	8.6	1000
423#1-3	+	30	8.8	1200
423#1-4	+	30	9.0	1200
423#1-5	+	60	9.0	1200
423#1-6	.+	60	9.0	1200

^{*}Test tubes cultures were grown in SVY + α -KIV (5 g/L) with Asp (10 g/L) where indicated.

The pAN423 transformants produced at least twice the amount of pantothenate as the controls (i.e., to a level at or near that which was obtained in earlier experiments by the addition of β-alanine to the culture medium). The data also show that in the absence of added aspartate, transformants containing additional copies of the panD gene expression cassette produce more pantothenate than the control transformants. One of the transformants, 423#1-5, produced about five times as much pantothenate as the controls. These results indicated that increased levels of PanD protein "pull" the conversion of available aspartate towards β-alanine, and that increasing panD gene expression can result in enhancement of pantothenate production both in the presence and absence of added aspartate.

Transformant 423#1-5 was re-named strain PA401 and studied further in shake

flask fermentations. The shake flask medium was SVY with maltose instead of SVY

with glucose. Results of shake flask experiments agreed well with test tube experiments

during the first 24 hours. In shake flask experiments without the addition of β-alanine,

PA401 produced approximately 1.5 g/l of pantothenate in 24 hours. Addition of β
alanine to the culture medium did not further improve pantothenate titers (Table 7),

indicating that with this strain and these fermentation conditions, β-alanine is not

limiting pantothenate production. In fact, when no β-alanine is fed, one can observe that

PA401 is secreting β-alanine in significant amounts into the medium.

^{**}TetR = Approximate Tet-resistance of transformant

Table 7. Shake flask cultures with strain PA401 (panD) with and without β -alanine.

	Amino ac	ids (g/l)		24 hours	
Initial β-ala Added	β-ala	Val	pН	OD600	Pantothenate (g/l)
0	0.7	1.5	7.5	13.7	1.5
5 g/l	7.1	1.4	7.6	12.4	1.5

Each value represents the average of duplicate 250 ml baffled flasks containing 50 ml of medium, incubated at 37°C with shaking (200 rpm).

Base Medium: SVY with 10 g/l α-KIV, 30 g/l maltose

2% Inoculum: SVY with Tet 15 grown 24 hours.

EXAMPLE V: Engineering the *panD* gene for Further Increased Synthesis of Aspartate Decarboxylase and Enhanced Production of Pantothenate

This Example describes the generation of improved ribosome binding sites (RBSs) in the panD gene to increase the translation of panD mRNA.

Increasing the translation of the panD gene mRNA by generation of synthetic panD RBSs

The RBS (SEQ ID NO:88) used to express panD in pAN423 is a synthetic RBS and has been used to successfully produce other proteins in B. subtilis at a high level. However, it contains six mismatches when aligned to the "ideal" B. subtilis RBS (SEQ ID NO:45) (e.g., an RBS having a sequence which is complementary to the 16S RNA sequence within the B. subtilis ribosome). (See e.g., Table 1B, mismatches in bold).

Two new RBSs were designed to more closely mimic the ideal RBS. These synthetic

RBSs, named new design A (NDA) and new design B (NDB) (also referred to herein as RBS3 and RBS4), are set forth as SEQ ID NO:51 and SEQ ID NO:52 and are aligned with the ideal RBS in Table 1B.

Oligonucleotides corresponding to the top and bottom strands of each new RBS were synthesized, annealed, then used to replace the RBS in pAN420, generating plasmids pAN426 and pAN427. These constructions are illustrated in Figure 6. The presence of the NDA and NDB RBS in pAN426 and pAN427 was confirmed by DNA sequence analysis. Next, the *panD* genes from pAN426 and pAN427 were transferred to *B. subtilis* expression vector pOTP61 as shown in Figure 7, creating pAN428 and pAN429. The nucleotide sequence of pAN429 is set forth as SEQ ID NO:79.

Not I restriction fragments lacking the E. coli vector sequences were isolated from pAN428 and pAN429, self-ligated, and used to transform strain PA221 to resistance to Tet¹⁵. Four isolates resistant to Tet⁶⁰ were picked from each transformation and assayed for pantothenate and β -alanine production along with PA221 transformed with the empty vector (pOTP61) and PA221 transformed with pAN423 (strain PA401) (see Table 8).

Table 8. Panthothenate production by test tube cultures of PA221 transformed with pAN428 and pAN429

	1	1 1	
Plasmid	Medium	OD550	Pan

Plasmid		Medium Supplements	OD ₅₅₀	Pan g/l	β-Ala g/l
pOTP61		α-KIV ⁵	10 10	UND 0.4	0:04
pAN423	ļ	α-KIV ⁵	10	0.4	0.04
pAN428-1	*	α -KIV ⁵	12	0.6	0.04
pAN428-2		α -KIV ³	11	0.5	0.03
pAN428-3		α-KIV	11	0.3	0.03
pAN428-4		α-KIV ⁵	10	0.1	UND
pAN429-1		α -KIV ⁵	12	0.6	0.04
pAN429-2		α -KIV ⁵	11	0.5	0.04
pAN429-3		α -KIV ²	11	0.6	0.05
pAN429-4	#	α-KIV ⁵	12	0.8	0.10
pOTP61		α -KIV ⁵ + Asp ¹⁰	11	0.5	0.08
pAN423		α -KIV ^S + Asp ¹⁰	12	0.9	1.32
pAN428-1	*	α -KIV ⁵ + Asp ¹⁰	12	0.8	1.97
pAN428-2		α -KIV ⁵ + Asp ¹⁰	12	0.8	1.51
pAN428-3		$\begin{array}{c} \alpha \text{-KIV}^5 + \text{Asp}^{10} \\ \alpha \text{-KIV}^5 + \text{Asp}^{10} \end{array}$	12	0.9	1.02
pAN428-4		α -KIV ⁵ + Asp ¹⁰	11	0.8	0.30
pAN429-1		α -KIV ⁵ + Asp ¹⁰	12	0.8	1.78
pAN429-2		α -KIV + Asp α -KIV ⁵ + Asp 10	12	0.8	1.66
pAN429-3			12	0.8	1.78
pAN429-4	#	α -KIV ⁵ + Asp ¹⁰	13	0.8	2.28

UND: Below the limits of detection. * Renamed PA402 # Renamed PA403

When grown in medium supplemented with a-KIV at 5 g/l (a-KIV⁵), the pAN428-1 transformant and all four of the pAN429 transformants produced more

pantothenate than did PA401, suggesting that these transformants contain higher levels of aspartate decarboxylase activity. When grown in medium supplemented with α -KIV⁵ and Asp¹⁰ none of the pAN428 or pAN429 transformants produced more pantothenate than PA401. However, the pAN428-1 transformant and all four of the pAN429 transformants produced significantly more β -alanine than did PA401. It is possible that the excess β -alanine produced from added aspartate causes inhibition of pantothenate production. Alternatively, β -alanine may accumulate because pantoate is limiting in these strains.

The strains that produced the highest level of β-alanine, the pAN428-1 and pAN429-4 transformants, were renamed PA402 and PA403, respectively. These two strains were grown in SVY medium supplemented with various intermediates and reassayed for pantothenate and β-alanine production. PA221 and PA401 were included as controls. The results of the assays are presented in Table 9.

15 Table 9. Pantothenate production of PA402 and PA403 in test tube cultures.

Strain	Medium Supplements	OD550	Pan g/l	β-Ala g/l	Val g/l
PA221	α-KIV ⁵	7.9	UND	UND	0.9
PA401	α-KIV ⁵	8.7	0.3	0.04	0.9
PA402	α-KIV ⁵	8.5	0.5	0.04	0.9
PA403	α-KIV ⁵	9.4	0.7	0.07	0.9
PA221	$\alpha\text{-KIV}^{5} + \text{Asp}^{10}$	9.8	0.4	0.11	0.8
PA401		9.1	0.8	1.15	0.8
PA402		9.4	0.8	2.02	0.8
PA403		9.7	0.7	2.40	0.8
PA221	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	8.9	UND	UND	0.2
PA401		8.7	0.3	0.02	0.2
PA402		10.6	0.5	0.02	0.2
PA403		10.5	0.7	0.02	0.2
PA221	Pantoate ⁵ + Asp ¹⁰	9.5	0.4	0.06	0.2
PA401		9.2	2.2	0.62	0.2
PA402		9.1	2.8	1.17	0.2
PA403		10.2	2.9	1.58	0.2

UND: Below the limits of detection.

When grown in medium supplemented with either α-KIV⁵ or Pantoate⁵, PA402 and PA403 produced significantly more pantothenate than did PA401. As before, even though PA402 and PA403 produced significantly more β-alanine than PA401 when grown in medium supplemented with α-KIV⁵ and Asp¹⁰, they did not produce a proportional increase in pantothenate. However, when grown in medium supplemented with Pantoate⁵ plus Asp¹⁰, both PA402 and PA403 produced significantly more pantothenate than PA401, about a 30% increase.

It can be concluded from these experiments that the improved NDA and NDB panD ribosome binding sites, engineered into pAN428 and pAN429, respectively, lead to increased levels of aspartate decarboxylase activity.

Increasing the translation of the panD gene mRNA by generation of synthetic panD RBSs within the panBCD operon

15

20

25

30

The native B. subtilis panD gene ribosome binding site (RBS) (SEQ ID NO:43), which is found in the $P_{26}panBCD$ operon cassette present in PA221 (and in other engineered pantothenate production strains described herein), is shown in Table 1C aligned with the ideal ribosome binding site (SEQ ID NO:47). The alignment shows mismatches between the native B. subtilis panD-gene RBS, which is located within the coding sequence for PanC, and the the ideal RBS. Three new RBSs (within the P26 panBCD operon cassette) were generated to increase translation of the panD gene mRNA and to yield increased synthesis of aspartate decarboxylase. These synthetic RBSs (termed NDI, NDII, and NDIII, also referred to herein as RBS5, RBS6 and RBS7. respectively) are set forth as SEQ ID NO:55, SEQ ID NO:56 and SEQ ID NO:57, respectively) and are included in Table 1C. It should be noted that although changes in the panD RBS within the panBCD operon also changes the C-terminal amino acid sequence of the PanC protein encoded by that operon, an alignment of known and suspected PanC protein amino acid sequences showed that the sequence of the last nine amino acids of the B. subtilis PanC protein could be altered without affecting any conserved amino acid residues indicating that such changes should not reduce pantothenate synthetase activity or expression. The new RBSs were synthesized and incorporated into the P_{26} panBCD operon expression cassette as follows.

First, PCR primers were designed to contain the following elements: (1) a

nucleic acid sequence encoding the first five amino acids of PanD up to and including a
unique BsiWI restriction site that had been previously introduced into panD by PCR; (2)

10

15

20

25

30

35

a stop codon for panC, (3) at least one synthetic RBS; and (4) 30-39 bp of nucleic acid sequence having 100% identity with panC upstream of the panD RBS. The primers were named TP102, TP103, and TP104 and contain the NDI, NDII, and NDIII ribosome binding sites, respectively. These three primers were used in conjunction with the 5' primer TP101, which hybridizes near the start codon of panC, in three independent PCR reactions to generate the NDI, NDII, and NDIII PCR products. The PCR products were purified, digested with XbaI, then cloned into plasmid vector pASK-1BA3 which had been digested with XbaI and SmaI. The resulting plasmids were named pAN431, pAN432, and pAN433. The construction of pAN431 is illustrated in Figure 8 and is representative of all three plasmid constructions. The presence of the desired synthetic panD gene RBS in each new plasmid was confirmed by DNA sequencing.

Next, the modified panC genes containing the new panD RBSs were joined with the panD gene utilizing the unique BsiWI restriction site. This was accomplished by isolating the appropriate NsiI-BsiWI restriction fragments from pAN431, pAN432, and pAN433 and ligating them with a 2395 bp NsiI-BsiWI restriction fragment from pAN420, which supplied the BsiWI-modified panD gene. These constructions resulted in plasmids pAN441, pAN442, and pAN443, respectively. A representative construction (pAN441) is illustrated in Figure 9. The nucleotide sequence of pAN443 is set forth as SEQ ID NO:80.

The new panD gene RBSs were then substituted into the P₂₆panBCD operon expression cassette as follows. First, a deletion-insertion mutation which removes the region of panC containing the panD RBS was created. This was constructed by digesting pAN430 with a mixture of BspE1 and BglII and recovering the 4235 bp fragment which is now missing the 3' end of panC and the 5' end of panD. This fragment was ligated with an AvaI-BamHI restriction fragment from plasmid pECC4, which contains the chloramphenical acetyl transferase (cat) gene. The 5' extension produced by AvaI digestion is compatible with that produced by BspEI while the BglII and BamHI extensions are also compatible. The resulting plasmid was named pAN440, and its construction is illustrated in Figure 10.

The resulting deletion-insertion mutation was crossed into the P_{26} panBCD operon via homologous recombination by transforming PA221 with linearized pAN440 and selecting for resistance to chloramphenicol on Cam⁵ plates containing 1 mM pantothenate. Several transformants were tested, and were all found to require 1 mM pantothenate for growth, as expected. Two of these transformants were remaned PA408A and PA408B and were assayed for pantothenate production. Neither strain synthesized measurable quantities of pantothenate, even when grown in medium

containing pantoate and β -alanine at 5 g/l, indicating that the strains are deficient in pantothenate synthetase activity. Next, the new panD RBSs were crossed into the P_{26} panBCD operon by transforming PA408 with linearized pAN441, pAN442, and pAN443 plasmid DNA and selecting for growth on TBAB plates without pantothenate supplementation. A transformation with linearized pAN430 (including the native panD RBS) was included as a control and was expected to give rise to transformants identical to PA221 described herein. Four isolates from each transformation were assayed for pantothenate and β -alanine production in SVY medium supplemented with various intermediates (Tables 10 and 11).

10
Table 10. Pantothenate production of PA410 - PA413 in test tube cultures.

Strain	RBS	Medium Supplements	OD ₅₅₀	Pan .g/l	β-Ala g/l
PA221	native	Pantoate ⁵	11	UND	UND
PA410-1 PA410-2 PA410-3 PA410-4	native	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 12 12 12	UND UND UND UND	UND UND UND UND
PA411-1 PA411-2 PA411-3 PA411-4	NDI	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 12 12 12	0.23 0.20 0.19 UND	UND UND UND UND
PA412-1 PA412-2 PA412-3 PA412-4	NDII	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 11 13 12	UND UND 0.18 0.18	UND UND UND UND
PA413-1 PA413-2 PA413-3 PA413-4	NDIII	Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5 Pantoate 5	12 12 12 12	0.18 0.17 0.16 0.17	UND UND UND UND

UND: Below the limits of detection.

Pantothenate production of PA410 - PA413 in test tube cultures. Table 11.

Strain	RBS	Medium Supplements	OD550	Pan g/l	β -Ala g/l
PA221	native	Pantoate ⁵ + Asp ¹⁰	11	0.3	UND
PA410-1 PA410-2 PA410-3 PA410-4	native	Pantoate ⁵ + Asp ¹⁰	12 12 12 12	0.4 0.4 0.4 0.4	UND UND UND UND
PA411-1 PA411-2 PA411-3 PA411-4	NDI	Pantoate 5 + Asp 10	13 13 13 13	1.7 1.7 1.8 0.4	0.4 0.4 0.3 UND
PA412-1 PA412-2 PA412-3 PA412-4	NDII	Pantoate 5 + Asp 10 Pantoate 5 + Asp 10 Pantoate 5 + Asp 10 Pantoate 5 + Asp 10 Pantoate 5 + Asp	13 12 12 12	0.4 0.4 1.6 1.5	UND UND 0.3 0.2
PA413-1 PA413-2 PA413-3 PA413-4	NDIII	Pantoate ⁵ + Asp ¹⁰	13 13 13 13	1.6 1.6 1.7 1.7	0.3 0.4 0.4 0.4

UND: Below the limits of detection.

As expected from previous experiments using PA221, none of the transformants that contained the native panD RBS produced measurable quantities of pantothenate when grown in medium supplemented with pantoate. However, nine of the twelve transformants expected to contain modified panD RBSs produced significant quantities of pantothenate (160-230 mg/l) under these conditions, indicating that they possess elevated levels of aspartate decarboxylase activity. When grown in medium supplemented with both pantoate and aspartate, these same nine transformants produced approximately four times more pantothenate than those with the native panD RBS. In addition, these nine transformants accumulated measurable quantities of β -alanine (230-410 mg/l). All transformants produced roughly equivalent quantities of pantothenate when grown in medium containing pantoate and β -alanine, demonstrating that each 15 contains a functional pantothenate synthetase.

5

10

These data demonstrate that the synthetic panD RBSs are about four times more effective than the native panD RBS in directing translation of the panD gene mRNA and evidence the utility of such synthetic RBSs in enhancing pantothenate production. Additional approaches to increasing pantothenate production can include, for example, increasing the half-life of the panD gene mRNA, increasing the strength of the promoter for panD transcription and/or increasing the stability of the PanD protein.

EXAMPLE VI: Construction of Strains Containing an Integrated P_{26} panE1 Cassette without an Antibiotic Resistance Gene.

10

20

Example II describes the identification of the B. subtilis panE1 gene that encodes the enzyme responsible for the majority of the ketopantoate reductase activity in B. subtilis. PA236 (containing the pAN236 plasmid) produced about twice as much pantothenate (2 g/l) as its parent strain, PA221 (1 g/l) in 24 hour SVY test tube cultures. PA236 was presumed to contain an amplified (\sim 3 copies) integrated pAN236 plasmid based on selection for tetracycline resistance (the tetR gene product being encoded on the pAN236 plasmid in addition to the P_{26} panE1 cassette). Also useful in the methodologies of the present invention are strains that contain a single integrated unamplifiable copy of P_{26} panE1 at the panE1 locus, for example, without an antibiotic resistance gene in the strain. Such a strain was generated as follows.

A plasmid named pAN251 was derived from pAN236 by inserting additional chromosomal sequences just upstream and just downstream from the P_{26} panE1 cassette. These additional sequences, which provide homology to allow integration of the P_{26} panE1 cassette at panE1 by double crossover, were obtained by PCR from chromosomal DNA as a template. pAN251 is shown in Figure 11. The nucleotide sequence of pAN251 is set forth as SEQ ID NO:81.

Next, a strain was constructed which allowed selection for the incoming P_{26} panE1 cassette. The strain included the following three components: (1) P_{26} panBCD; (2) $\Delta panE1$; and (3) ilvC, since both panE1 and ilvC must be mutated to have a Panphenotype. The starting strain was CU550 (trpC2, ilvC4, leuC124). The P_{26} panBCD cassette from PA221 chromosomal DNA was introduced in two steps to create strain PA290. Next, $\Delta panE1$::spec was transformed into PA290, using chromosomal DNA from strain PA240, to give strain PA294 (trpC2, ilvC4, leuC124, P_{26} panBCD, $\Delta panE1$::spec), which is a strict pantothenate auxotroph. Finally, PA294 was transformed with plasmid pAN251, selecting for pantothenate prototrophy, to give strain PA303. This strain was expected to have the genotype trpC2, ilvC4, leuC124, P_{26}

20

25

panBCD, P_{26} panE1. PA303 was checked for the correct chromosomal structure at the panE1 locus by PCR using primers that flank the P_{26} insertion just upstream of panE1. The PCR product from PA303 was of the expected size, with a concomitant loss of the PCR product from the wild type panE1 gene, consistent with having obtained the desired double crossover event. Furthermore, PA303 was tetracycline sensitive, which is also consistent with the desired double crossover event, as opposed to a Campbell-type single crossover of the plasmids into the chromosome. The trp, ilv, and leu auxotrophies from the parent strain were all maintained in PA303.

In 24 hour liquid SVY test tube cultures, PA303 produced almost the same level of pantothenate as positive control PA236, and about twice as much as PA221, which does not contain engineered *panE1* as indicated in Table 12.

Table 12. Pantothenate production by 24 hr. test tube cultures of PA303 and controls grown in SVY plus 5 g/l α -KIV and 5 g/l β -alanine.

Strain	OD600	[pan] g/l
PA221-1	10.9	0.85
PA221-2	10.5	0.85
PA236-1	9.5	1.74
PA236-2	9.3	1.70
PA303-1	10.8	1.66
PA303-2	10.7	1.61

EXAMPLE VII: Generation of Microorganisms Capable of Producing Pantothenate in an α-KIV (or Valine) Independent Manner

 α -ketoisovalerate (α -KIV) is a rate limiting intermediate for pantothenate production in certain strains deregulated for pantothenate synthesis. Addition of either α -KIV or valine at 5 g/l increases pantothenate production about 5-fold in test tube cultures with strains such as PA221. In order to alleviate the need to feed either α -KIV or valine, strains were engineered that have an increased capacity to synthesize α -KIV.

 α -KIV is produced in B. subtilis from pyruvate by the sequential action of three enzymes encoded by four genes, ilvB and ilvN, ilvC, and ilvD. In a wild type B. subtilis, three of the genes (ilvB, ilvN, and ilvC) are the first three genes of the large ilv-leu operon. The fourth gene necessary for α -KIV synthesis, ilvD, is located by itself elsewhere on the chromosome. The B. subtilis ilv-leu operon is thought to be regulated

10

15

20

only by leucine levels. Feeding of exogenous leucine reduces transcription of the *ilv-leu* operon by about 13-fold, probably by an attenuation mechanism (Grandoni *et al.* (1992) *J. Bacteriol.* 174: 3212-3219). The only known feedback regulation in the *ilv-leu* pathway is the inhibition of the *leuA* gene product by leucine.

As a first step to deregulate the synthesis of α -KIV, a copy of the *ilvBNC* region from the wild type *B. subtilis ilv-leu* operon was isolated by PCR, and installed adjacent to the P_{26} promoter and RBS2 on a vector, pOLL8, that was designed to integrate a single P_{26} expression cassette by double recombination at the *amyE* locus. The *amyE* gene encodes a nonessential α -amylase, and is a useful locus for installing expression cassettes. The resulting plasmid, pAN267, is illustrated in Figure 12. The nucleotide sequence of pAN267 is set forth as SEQ ID NO:82. pAN267 readily gave stable transformants by double crossover at the *amyE* locus of *B. subtilis* strains, as described in detail below.

Construction of pantothenate overproducing strains that are leucine prototrophs

Initially, a *B. subtilis* strain containing ilvC4 and $\Delta panE1$ was used to introduce a single copy of P_{26} panE1 into the chromosome without using an antibiotic resistance gene. The double mutant was required to select for the incoming P_{26} panE1 cassette because a $\Delta panE1$ mutation alone does not result in pantothenate auxotrophy. A strain named CU550 was obtained containing ilvC4 to be used as a basis for this type of strain construction. However, CU550 also contains a closely linked leuC124 mutation, so all strains derived from CU550 required leucine. Having shown that the combination of P_{26} panBCD and P_{26} panE1 was favorable for pantothenate production, the next step was to reassemble this combination of two cassettes in a leucine prototroph.

Accordingly, the two cassettes were combined in two different strain backgrounds, RL-1 and PY79. To introduce chromosomal P_{26} panE1 into the PY79 and RL-1 strain backgrounds without using an antibiotic resistance gene, a strategy was used that did not rely on ilvC4. (The strategy took advantage of the observation that the $\Delta panE1$ mutation causes a pantothenate bradytrophy, manifested by relatively small colonies on TBAB (rich) plates). First, $\Delta panB$::cat and $\Delta panE$::spec were introduced into both strain backgrounds. Next, the resulting strains were transformed simultaneously with DNA from two strains, PA221 (P_{26} panBCD) and PA303.(P_{26} panE1), selecting for Pan⁺ on TBAB plates. Colonies of two distinct sizes grew on the selective plates, with the larger size comprising about 2% of the colonies. The larger colonies were presumed to represented co-transformants that received both P_{26} panBCD and P_{26} panBCD. Consistent

with this prediction, the larger colonies had lost both Cam^r and Spec^r, while the smaller colonies had lost only the *cat* gene, and retained the *spec* gene. Furthermore, a representative derivative of PY79 named PA327, and a representative derivative of RL-1, named PA328, both produced the elevated levels of pantothenate in test tube cultures which was about 1.6 to 1.7 g/l (Table 13).

Table 13. Pantothenate production of PA327, PA328, and controls from 24 hr test tube cultures grown in SVY plus 5 g/l α-KIV and β-alanine.

Strain	Background	P26 panE1 copy number	[pan] g/l	
PA221-1	RL-1	0	0.92	
PA221-2	RL-1	0	0.95	
PA236-1	RL-1	amplified (~3)	1.60	
PA236-2	RL-1	amplified (~3)	1.73	
		• • •		
PA327-1	PY79	1	1.66	
PA327-2	PY79	1	1.65	•
				•
PA328-1	RL-1	1	1.61	
PA328-2	RL-1	1 '	1.91	•
1115252	1 1	-		

10

20

Thus, PA327 and PA328 were concluded to contain both P_{26} panBCD and P_{26} panE1, and were used for further constructions as described below. PCR analysis confirmed the presence of the two cassettes.

15 <u>Installation of a stable P26 ilvBNC cassette into two lineages of pantothenate overproducing strains</u>

Having constructed PA327 and PA328, derivatives of PY79 and RL-1 that contain P26 panBCD and P26 panE1, and that are Leu⁺, the next step was to introduce stable copies of P26 ilvBNC. This was accomplished by transforming PA327 and PA328 with plasmid pAN267, selecting for Spec^r. Screening by PCR showed that about 85% of the obtained transformants contain P26 ilvBNC integrated at amyE by double crossover. One transformant of PA327, named PA340, and one transformant of PA328, named PA342, were chosen for further study.

In test tube cultures grown in SVY medium plus 5 g/l β-alanine but without added α-KIV, both PA340 and PA342 gave the expected increase in pantothenate production over that of PA327 and PA328, to about 1.3 to 2 g/l (Table 14).

15

Table 14. Pantothenate and valine production by PA340 and PA342, both containing P_{26} ilvBNC in 24 hr test tube cultures grown in SVY with 5-g/l β -alanine and with or without 5 g/l α -KIV

Strain	Back- ground	OD600 - α-KIV	+ α-KIV	{pan] g/l - α-KIV	+ α-KIV	[val] g/l - α-KIV	+ α- K IV
PA340-1 PA340-2	PY79 PY79	11.8 10.3	7.1 7.5	2.02	2.10 2.03	0.38	0.90 0.91
PA342-1 PA342-2	RL-1 RL-1	10.2	8.0 9.2	1.29	1.89 2.04	0.27	0.78 0.79

The two new strains also gave a slight increase in valine secretion, indicating that the *ilvBNC* genes had been deregulated. However, when the same strains were grown with 5 g/l α-KIV added, a further increase in pantothenate production occurred from PA342, suggesting that α-KIV was still rate limiting in this strain background. Similar results, only with more growth and hence higher pantothenate levels, were seen in shake flask cultures (Table 15).

Table 15. Pantothenate and valine production by PA340 and PA342, both containing P_{26} ilvBNC in 24 hour shake flask cultures grown in SVY with 5 g/l β -alanine and with or without 5 g/l α -KIV.

Strain	Back- ground	OD600 - α-KIV	+ α-KIV .	[pan] g/l - α-KIV	+ a-KIV	{val] g/l - α-KIV	+α-KIV
PA327	PY79	21	22	0.6	3.0	0.5	1.3
PA340-1	PY79	20	20	3.5	4.1	1.0	1.9
PA340-2	PY79	22	19	3.0	2.1	0.8	1.4
PA328	RL-1	20	16	1.4	2.7	0.6	1.3
PA342-1	RL-1	17	16	3.3	3.6	0.9	1.6
PA342-2	RL-1	18	18	3.1	4.2	0.8	1.4

EXAMPLE VIII: Increasing panD Copy Number in Strains Engineered to Overproduce panE1 and the ilvBNC Gene Products Enhances Pantothenate **Production**

Experiments where β-alanine was fed to cultures of engineered B. subtilis strains consistently showed that \beta-alanine was a rate limiting intermediate in pantothenate synthesis. The effect of adding additional copies of panD on pantothenate production in PA340 and PA342 was examined. Strains PA340 and PA342 were transformed with chromosomal DNA isolated from PA401 with selection on plates containing 15 µg/ml of tetracycline (Tet15 plates). Transformants derived from each parent were patched onto Tet⁶⁰ plates to identify those which were likely to contain multiple copies of the expression cassette. Twelve transformants from each transformation which grew on Tet⁶⁰ were streaked for single colonies on this medium and then assayed in SVY medium test tube cultures for pantothenate production. One transformant from each group was found to produce greater than 300 mg/l pantothenate in 24 hours. These two 15 transformants were saved and named PA404 (PA340 strain background) and PA405 (PA342 strain background). Both strains were resistant to spectinomycin, indicating that the P_{26} ilvBNC expression cassette was still present at amyE. PCR analysis of chromosomal DNA isolated from each strain confirmed that the deregulated panEl gene had also been retained.

20 Next, PA404 and PA405 were evaluated in shake flask cultures which were grown in SVY medium containing maltose as the carbon source and supplemented with various intermediates. The cultures were grown for 24 and 48 hours and then assayed for pantothenate, β-alanine, and valine production. The results of this experiment are presented in Table 16. Analogous shake flask culture data for the parent strains (PA340 25 and PA342) are included in the tables for comparison.

Table 16. Pantothenate production by PA404 and PA405 in shake flask cultures after 24 hours

Strain	Medium Supplements	ОД600	Pan g/i	β-Ala g/l	Val g/l
PA340	none	20	0.4	<0.1	1.0
PA404	none	22	1.8	<0.1	0.7
PA342	none	19	0.3	0.2	0.7
PA405	none	19	1.4	0.4	0.5

PA340	β-alanine ⁵	18	3.6	3.2	0.6	
PA404	β-alanine ⁵	18	2.8	5.1	0.7	
PA342*	β-alanine ⁵	17	3.3	3.3	0.5	
PA405*	β-alanine ⁵	19	1.3	6.5	0.6	

Values are the average of duplicate flasks except where indicated by *.

In the absence of any medium supplementation, PA404 and PA405 made four to five times more pantothenate in 24 hours compared to their isogenic parent strains (Table 16). The supply of β -alanine was clearly limiting in the parent strains PA340 and PA342. Addition of amplified *P26 panD* greatly increased the supply of β -alanine.

EXAMPLE IX: Deregulation of the *B. subtilis ilvD* Gene Enhances Pantothenate Production

10 To deregulate expression of the *ilvD gene*, standard procedures (described above) were used to integrate the constitutive P_{26} promoter and an artificial ribosome binding site, RBS2, just upstream of the *ilvD* coding region. The *ilvD* gene maps by itself, unlinked to the *ilvBNC* operon. First, a 2.4 kb region of the RL-1 chromosome that contains the *ilvD* coding region and 730 bp of upstream sequence was cloned by PCR into a low copy (about 15 per *E. coli* cell) vector called pOK12, to give plasmid pAN257, shown in Figure 13.

Taking advantage of a natural EcoRI site just upstream of the native ilvD gene promoter, and a natural NcoI site at the ilvD start codon, an artificial sequence containing P_{26} and RBS2 was inserted into pAN257 to give pAN263 (Figure 14). The nucleotide sequence of pAN263 is set forth as SEQ ID NO:83. In parallel with this construction, the cat gene was also inserted into pAN257, between the same upstream EcoRI site and a BgIII site in the middle of the ilvD coding region, to give pAN261, which is deleted for a large portion of the ilvD gene (Figure 15). Using pAN261 and pAN263, the P_{26} ilvD cassette could then be installed in the B subtilis chromosome in two steps. In the first step, pAN261 is introduced by transformation, selecting for chloramphenicol resistance, and then confirming an Ilv phenotype. In the second step, pAN263 is introduced, selecting for Ilv⁺, checking for chloramphenicol sensitivity, and confirming correct local structure by PCR.

pAN261 was first transformed into strain RL-1 (highly competent) to give strain PA343 (\(\Delta\ildelt

20

in transforming *B. subtilis*. Similarly, pAN263 DNA was transformed into PA343 (moderately competent) to give strain PA345 (P_{26} ilvD), and then PA345 chromosomal DNA was used to transform PA348 and PA349 to Ilv⁺ prototrophy, yielding strains PA374 and PA354, respectively.

As predicted, PA374 and PA354 gave further increases in pantothenate production, to about 2.5 to 2.9 g/l, in test tube cultures grown in SVY plus 5 g/l β-alanine (Table 17).

Table 17. Pantothenate and valine production by PA374 and PA354, containing P_{26} ilvD, and controls, in 24 hr test tube cultures grown in SVY with 5 g/l β -alanine and with or without 5 g/l α -KIV.

		ı	OD ₆₀₀		[pan] g/l		[val] g/l	
Strain	Back- ground	ilvD status	α-KIV -	+	α-KIV -	+	α-KIV -	+
PA340	PY79	w.t.	9.2	9.0	2.14	2.23	0.38	0.90
PA348	PY79	ilvD::cat	11.7	10.0	0.19	2.23	0.19	0.91
PA374-1 PA374-2	PY79 PY79	P ₂₆ ilvD P ₂₆ ilvD	9.1 8.2	7.3 7.7	2.93 2.99	2.40 2.36	0.58 0.60	0.87 0.95
PA342	RL-1	w.t.	10.2	8.0	1.29	1.89	0.27	0.78
PA349	RL-1	ilvD::cat	8.1	7.7	0.17	1.87	0.22	0.88
PA354-1 PA354-2	RL-1 RL-1	P26 ilvD P26 ilvD	9.6 7.5	9.6 8.2	2.57 2.48	2.03 2.24	0.65 0.64	1.23 0.97

In the absence of added β -alanine, strains PA374 and PA354 produced only about 0.2 g/l pantothenate in test tube cultures, indicating that PanD activity is significantly rate limiting.

To alleviate this limitation, the amplifiable P_{26} panD cassette from strain PA401 was installed. PA401 chromosomal DNA was transformed into PA374 and PA354, selecting for Tet^r at 15 mg/l, to yield strains PA377 and PA365, respectively. After transformants were obtained, the strains were streaked on plates containing 30 and 60 mg/l tetracycline to reamplify the copy number of the P_{26} panD cassette integrated at the bpr locus. In test tube cultures grown in SVY without α -KIV or β -alanine, a substantial improvement in pantothenate titers over those of PA374 and PA354 was obtained (Tables 18 and 19).

20

Table 18. Pantothenate production by PA365, containing amplified P_{26} panD, and controls, in 24 and 36 hr test tube-cultures grown in SVY-glucose without β -alanine or α -KIV.

		OD ₆₀₀		[pan] g/l		
Strain	Relevant genotype	24 hrs.	36 hrs	24 hrs.	36 hrs.	
PA342-1-1	w.t. ilvD	11.7	8.8	b.d.	0.27	
PA342-1-2	w.t. <i>ilvD</i>	12.8	8.8	b.d.	0.26	
PA354-1-1	P ₂₆ ilvD	n.d.	11.0	n.d.	0.19	
PA354-1-2	P ₂₆ ilvD	n.d.	8.4	n.d.	0.20	
PA365-1	P ₂₆ ilvD, P ₂₆ panD	9.8	10.0	1.01	2.07	
PA365-2	P ₂₆ ilvD, P ₂₆ panD	9.9	10.4	0.96	2.09	

n.d. = not determined; b.d. = below detection

Table 19. Pantothenate production by PA377, containing amplified P_{26} panD, and controls, in 27 hr test tube cultures grown in SVY-glucose or SVY-maltose, without α -KIV, and with or without β -alanine.

		OD_{600}			
	Relevant genotype	- β-ala	+β-ala	- β-ala	+β-ala
Strain		Glucose	Glucose	Maltose	Maltose
PA374-1	P26 ilvD	9.4	9.8	7.0	6.4
PA374-2	P ₂₆ ilvD	9.2	9.6	6.6	6.3
		1			
PA377-1	P26 ilvD, P26 panD	10.0	7.6	7.2	6.1
PA377-2	P26 ilvD, P26 panD	10.5	7.8	9.4	5.4
	•] [pan] g/l			
	Relevant genotype	[pan] g/l - β-ala	+ β-ala	- β-ala	+ B-ala
Strain	Relevant genotype	1	+ β-ala Glucose	- β-ala Maltose	+β-ala Maltose
Strain	Relevant genotype	- β-ala	•	•	•
Strain PA374-1	Relevant genotype P26 ilvD	- β-ala	•	•	•
		- β-ala Glucose	Glucose	Maltose	Maltose
PA374-1	P ₂₆ ilvD	- β-ala Glucose	Glucose 2.76	Maltose 0.14	Maltose 1.31
PA374-1	P ₂₆ ilvD	- β-ala Glucose	Glucose 2.76	Maltose 0.14	Maltose 1.31
PA374-1 PA374-2	P ₂₆ ilvD P ₂₆ ilvD	- β-ala Glucose 0.04 0.10	Glucose 2.76 2.65	Maltose 0.14 0.15	1.31 1.33

In SVY with glucose, an increase in pantothenate production can still be
achieved by feeding 5 g/l β-alanine suggesting that increasing panD expression further
might increase pantothenate production. In SVY with maltose, no further increase in
pantothenate was obtained by feeding β-alanine suggesting that β-alanine and/or

aspartate synthesis is suppressed by glucose. Strains PA377 and PA365 have been evaluated in 10 liter fermentors, where they typically produce above 20 g/l pantothenate in 48 hours without supplemental β -alanine and α -KIV or valine, described in detail below.

5

10

EXAMPLE X: 10 liter Fermentations of Pantothenate-Producing Microbes

Engineering of the P_{26} ilvBNC and P_{26} ilvD cassettes to give strains PA342 and PA354 allowed the production of 22 and 26 g/l of pantothenate, respectively, without the addition of valine or α -KIV to the fermentation medium (Table 20). At 48 hours, both strains had secreted about 0.5 g/l of valine into the medium.

Table 20. 10-liter fermentations of five pantothenate overproducing strains.

,	Strain	Medium	Feed 40% Glucose plus	OD 600 48 hr	Valine 48 hours g/l	β-ala 48 hr g/l	Pa 36 hr	ntothena g/L 48 hr	te 72 hr
	PA 236	svyg	50 g/l β-ala 25 g/l α-KIV	108	added	· added	16	19	21
	PA 342	SVYG	50 g/l β-ala	92	0.5	added	17	22	
	PA 354	SVYG	50 g/l β-ala	90	0.5	added	19	26	
	PA 365	SVYG	25g/l YE	77	0.85	0.4	18	21	27
	PA 377	SVYG	25g/l YE	85	1.5	0.5	18	22	31
	PA 377	PFMG	25g/l YE	96	0.8	0.4	19	25	29
	PA377	PFMG	-	71	0.7	0.1	16	21	-

15 Pantothenate synthesis in fermentors

With the addition of the P_{26} panD cassette to strains PA354 and PA374 to create strains PA365 and PA377, neither β -alanine nor α -KIV needed to be added to the fermentors. Strain PA365 produced 21 g/l pantothenate in 48 hours and 27 g/l in 72 hours with no precursors added to the medium (Table 20). PA377 was somewhat better, producing 18 g/l of pantothenate in 36 hours, 22 g/l in 48 hours, and 31 g/l in 72 hours). Valine was measured at 0.85 and 1.5 g/l for strains PA365 and PA377, respectively, at

20

48 hours in SVYG medium. Strain PA377 maintained valine between 1-1.5 g/l throughout most of the fermentation and β -alanine between 0.2 and 0.5 g/l.

Strain PA377 was further evaluated in 10-liter fermentors in yeast extract based PFMG medium. Pantothenate yields in PFMG and SVYG medium were similar. In PFMG, PA377 produced 19 g/l of pantothenate in 36 hours, 25 g/l in 48 hours, and 29 g/l in 72 hours. In SVYG, PA377 produced 18 g/L pantothenate in 36 hours, 22 g/L in 48 hours and 31 g/L in 72 hours (Table 20).

EXAMPLE XI: Converting Strain PA377 to a Tryptophan Prototroph

10,

PA377 (Trp) was transformed to Trp⁺ using chromosomal DNA from PY79 to give strain PA824. After re-amplification of the $P_{26}panD$ casette, PA824 was compared to PA377 for pantothenate production in test tube cultures grown in SVY glucose with or without 5 g/L β -alanine (Table 21).

15

Table 21: Trp^{+} derivatives of PA377: Pantothenate production in 48 hour test tube cultures grown in SVY glucose, $\pm \beta$ -alanine

		OD600		[pan].g/L	
Strain	trpC donor	- β-alanine	+ β-alanine	- β-alanine	+ β-alanine
PA377-1	RL-1	8	8	1.5	3.4
PA377-2	RL-1	8	9	1.6	3:6
PA824-1	PY79	12	10	0.7	3.7
PA824-2	PY79	11	11	1.9	4.9
				1	

20

The Trp+ strains grew to slightly higher densities than PA377. In the absence of exogenous β -alanine, all of the strains produced similar levels of pantothenate, while with the addition of β -alanine, the Trp+ derivatives produced somewhat more pantothenate.

25 Fermentor studies with PA824

PA824 was evaluated in CF3000 Chemap 14 liter vessels with 10 liter working volumes. Formulations for two of the media used in the fermentors are given in Tables 22 and 23.

Table 22: Formulation for PFMG-5 medium

BATCH

	2.1	
	MATERIAL	g/L (final [])
1	Amberex 1003	10
2	Na Glutamate	5 '
3	(NH ₄) ₂ SO ₄	8
4	MAZU DF 37C	2.5
	Added After Sterilization and Co	ol Down
		<u> </u>
1	KH₂PO₄	10

20 2 $K_2HPO_4 \cdot 3H_2O$ Glucose 20 1 1 $MgCl_2{\cdot}6H_2O$ 2 3 CaCl₂·2H₂O 0.1 1 Sodium Citrate 1 0.01 FeSO₄·7H₂O 2 SM-1000X $1.0 \; ml$ 3 qs to 6000 ml H_2O

5

FEED

	MATERIAL	g/L
1	Glucose	600
2	CaCl ₂ ·2H ₂ O	0.6
	H ₂ O	qs to 3000 ml

Table 23: Formulation for SVY-4 medium

BATCH

	MATERIAL	g/L (final []) 25		
1	Veal Infusion			
2	Yeast Extract	5		
3	Na Glutamate	5		
4	(NH ₄) ₂ SO ₄	. 4		
5	MAZU DF 37C	2.5		
·				

Added After Sterilization and Cool Down

1	KH₂PO₄	10 20		
2	K₂HPO₄·3H₂O			
· 1	Glucose	20		
2	MgCl ₂ ·6H ₂ O	1		
3	CaCl ₂ ·2H ₂ O	0.1		
1	Sodium Citrate 1			
2	FeSO ₄ ·7H ₂ O	0.01		
3	SM-1000X	1.0 ml		
	H ₂ O	qs to 6000 ml		

5

FEED

	MATERIAL	g/L	
1	Glucose	600	
2	CaCl₂·2H₂O	0.6	
	H ₂ O	qs to 3000 ml	

All fermentations were glucose limited fed batch processes. Immediately after inoculation, agitation was set at 200 rpm. The initial batched 2% glucose was consumed during exponential growth. Afterwards, glucose concentrations were maintained

10

15

20

25

30

between 0.2 and 1.0 g/L by continuous feeding of a 60% glucose solution. The variable rate feed pump was computer controlled and linked to the dissolved oxygen concentration [pO₂] in the tank by an algorithm. When the [pO₂]fell to 30%, computer control began to automatically adjust the agitation rate to maintain a dissolved oxygen concentration between 25 and 30% [pO₂]. Computer control and data recording were by Braun MFCS software.

In one study, PA284 was grown in fermentors at two temperatures (40°C and 43°C) in the medium described in Table 22. Results of two experiments demonstrated that the highest pantothenate titers at early time points were produced at 43°C. The cell mass approached 150 optical density units at OD₆₀₀ and 56 hours at 43°C, and the pantothenate titers were 21 g/L, 28 g/L and 36 g/L at 36, 48 and 72 hours respectively. In the parallel fermentation at 40°C, the cell mass approached 120 optical density units at OD₆₀₀ and 56 hours, and the pantothenate titers were 18 g/L, 26 g/L and 37 g/L at 36, 48 and 72 hours, respectively.

In another study, PA824 was grown in a fermentor at 43°C in the medium described in Table 23. The cell mass exceeded 160 optical density units at OD₆₀₀ and 36 hours, and the pantothenate titers were 23 g/L, 34 g/L, 37 g/L and 40 g/L at 24, 36, 48 and 60 hours, respectively. In other fermentations, increasing the amount of trace elements in the glucose feed (e.g., increasing the concentration of SM from 1X to 2X) resulted in even higher titers of pantothenate.

EXAMPLE XII: Identification and characterization of the B. subtilis coaA gene product

The annotated version of the *B. subtilis* genome sequence available on the "Subtilist" web site contains no gene labeled as *coaA*. However a homology search using the protein sequence of *E. coli* pantothenate kinase as a query sequence gave a good match with *B. subtilis* gene *yqjS*, which is annotated as "unknown; similar to pantothenate kinase." This gene appears to be the penultimate gene in an operon containing five open reading frames (Figure 18). Two of the open reading frames encode proteins which are similar to D-serine dehydratase and to "ketoacyl reductase"; the other two have no known homologies. For the open reading frame corresponding to *coaA*, there are three possible start codons; each having a possible ribosome-binding site (RBS) associated with it. The three potential *coaA* ORFs were named *coaA1*, *coaA2*, and *coaA3*, from longest to shortest.

All three potential coal open reading frames were cloned along with their respective RBSs by PCR followed by ligation into expression plasmid pAN229. pAN229 is a low copy vector in *E. coli* that provides expression from the SP01 phage P₁₅ promoter and can integrate by single crossover at bpr with tetracycline selection. A representative resulting plasmid, pAN281, is shown in Figure 19.

To determine if the cloned putative coal ORFs actually encode a pantothenate kinase activity, several isolates of all three plasmids were transformed into the E. coli strain YH1, that contains the coal 15(Ts) allele. Transformants were streaked to plates incubated at 30° and 43°C to test for complementation of the temperature sensitive allele. All isolates of all three coal variants, except for one isolate of pan282, complemented well at 43°C, indicating that all three plasmid constructs encode an active pantothenate kinase. Accordingly, it can be concluded that the B. subtilis yajS open reading frame codes for an active pantothenate kinase.

15 EXAMPLE XIII: Deletion of the coaA gene from the B. subtilis genome

The coaA gene of B. subtilis (yqjS) was deleted from the chromosome of a B. subtilis strain by conventional means. The majority of the coaA coding sequence was deleted from a plasmid clone and replaced by a chloramphenical resistance gene (cat), while leaving approximately 1 kb of upstream and downstream sequence to allow homologous recombination within the chromosome, to give plasmid pAN296 (see Figure 17). pAN296 was then used to transform a B. subtilis strain (PY79), selecting for chloramphenical resistance. The majority of transformants result from a double crossover event that effectively substitutes the cat gene for the coaA gene. The transformed strain containing the coaA deletion—cat insertion grew normally due the presence of a second B. subtilis pantothenate kinase encoding gene described herein.

EXAMPLE XIV: Identification and characterization of a second *B. subtilis* gene encoding pantothenate kinase activity

As described in detail in the instant specification, in order to maximize

pantothenate production, it is necessary to restrict the flow of pantothenate toward

Coenzyme A (CoA), for example, by reducing the activity of pantothenate kinase, the

first enzyme in the pathway from pantothenate to CoA. After finding that deletion of
the coaA gene from the chromosome of B. subtilis is not a lethal event (see Example

XIII), it was concluded that B. subtilis must contain a second gene that encodes an active
pantothenate kinase, since pantothenate kinase is an essential enzyme activity.

A second pantothenate kinase-encoding gene was identified by complementing the *E. coli* strain YH1 (coaA15(Ts)) with a *B. subtilis* gene bank and selecting for transformants that were able to grow at 43°C. Found among the transformants were two families of plasmids that had overlapping restriction maps within each family, but not between the families. As expected, the restriction map of one family was identical to that predicted from the *B. subtilis* genome sequence for the homologue of the *E. coli coaA* gene (which we named coaA also, see above) and surrounding sequences. The other family had a restriction map that was completely non-overlapping with the first.

DNA sequencing of the ends of the cloned inserts from the second family showed that the clones came from a region of the *B. subtilis* chromosome that includes the 3' end of the *fts*H gene, the 5' end of the *sul* gene, and all of the *yacB*, *yacC*, *yacD*, *cysK*, *pabB*, *pabA* and *pabC* genes. None of the open reading frames of these cloned inserts showed homology to any known pantothenate kinase sequences, either prokaryotic or eukaryotic.

Several deletions were created through the B. subtilis genomic sequences in the cloned inserts. Each deletion was tested for complementation of the E. coli temperature sensitive pantothenate kinase. In particular, a deletion that removed all DNA between a Stu I site in the cloning vector and a Swa I site in the yacC gene, leaves yacB as the only intact open reading frame in the cloned insert (see Figure 21). This deleted plasmid still complemented the E. coli pantothenate kinase mutant. However, another deletion that removed DNA from the Swa I site in yacC through a Bst1107I site in the (already truncated) ftsH gene, could not complement the E. coli pantothenate kinase mutant. From these results, it was concluded that the yacB open reading frame was responsible for the complementation activity. To confirm that yacB is a pantothenate kinase gene, the yacB ORF plus 112 base pairs of downstream flanking sequence was amplified by PCR in two independent reactions and cloned downstream of a constitutive promote to give plasmids pAN341 and pAN342 (Figure 22). Both pAN341 and pAN342 complemented the defect in YH1 at 44°C, while a control plasmid, which has the same backbone, but expresses panBCD instead of yacB did not. This confirmed that the yacB open reading frame was responsible for the complementation of YH1.

As such, a novel gene that encodes pantothenate kinase activity in *B. subtilis* has been discovered that is not related by homology to any previously known pantothenate kinase gene. This gene has been renamed *coaX*, as a second, alternative gene that encodes an enzyme that catalyzes the first step in the pathway from pantothenate to CoaA. Deletion of *coaX* by methods described above for deleting *coaA*, in conjunction

10

15

20

30

with reduction in the activity of the CoaA enzyme, provides a means to reduce pantothenate kinase activity to the desired level.

Several homologues of the *B. subtilis coaX* gene were identified by homology searching of various publically available databases using the published yacB (coaX) open reading frame sequence and predicted amino acid sequence (as set forth in SEQ ID NOs:84 and 85 respectively). In two cases (Mycobacterium tuberculosis and Streptomyces coelicolor) the homologous coaX genes are adjacent to, or almost adjacent to, pantothenate biosynthetic genes, consistent with these homologs having a role in pantothenate metabolism. The CoaX proteins show no homology to the CoaA family of pantothenate kinases, nor to the eukaryotic family of pantothenate kinases exemplified by PanK of Saccharomyces cerevisiae.

Alignment of the amino acid sequences of several bacterial CoaX homologs with the amino acid sequence predicted from translating the *B. subtilis yacB* ORF described in the published *B. subtilis* genome sequence revealed that the CoaX proteins from other bacteria contained additional amino acid residues at their carboxy-terminal ends. Moreover, these extensions beyond the end of the predicted amino acid for the *B. subtilis* gene product contained two relatively well-conserved segments of sequence.

Translation of nucleotide sequences just downstream from the stop codon of the B. subtilis yacB ORF in a different reading frame revealed the existence of amino acid sequences very similar to the carboxy-terminal extensions of the other bacterial CoaX proteins. It is thus believed that an error exists in the published DNA sequence of the B. subtilis yacB ORF sequence that causes a frame shift leading to an artifactual downstream amino acid sequence and premature termination.

The PCR-generated sequences of *B. subtilis CoaX* in pAN341 and pAN342 (described above) contain enough downstream flanking sequence to encode the putative carboxy-terminal extension described above, which is consistent with the result that the clones were functional in the complementation assay. However when the 3' PCR primer was positioned to include only the shorter *yacB* ORF predicted from the published sequence, but not to include the putative carboxy-terminal extension, then the resulting plasmids, pAN329 and pAN330 (similar in structure to pAN341 and pAN342; *see* Figure 22), did not complement the defect in YH1. This result supports the notion that the published *yacB* coding sequence contains a frame-shift error, and that the carboxy-terminal end of CoaX is necessary for pantothenate kinase activity. The predicted correct nucleotide sequence for *B. subtilis coaX* is set forth as SEQ ID NO:19 and the translated amino acid sequence is set forth as SEQ ID NO:9. A multiple

20

sequence alignment of the CoaX amino acid sequences of B. subtilis and 11 homologues thereof is set forth in Figure 23.

EXAMPLE XV: Generation of mutant coaA genes encoding pantothenate kinase having reduced or temperature sensitive activities

This Example describes strategies for modifying the *coaA* gene (i.e., by introducing point mutations) to reduce the activity of pantothenate kinase after *coaX* is deleted from the genome.

10 Cloning and sequencing of the temperature sensitive allele of the E. coli coaA gene.

Two E. coli strains, each exhibiting a different mutant CoaA phenotype, were obtained from the E. coli Genetic Stock Center. Strain DV62 contains the coaA15(Ts) allele, and DV79 contains the coaA16(Fr) mutation. DV62 is temperature sensitive at 43°C and produces a pantothenate kinase that is temperature sensitive. DV79 was obtained by reversion of DV62 to temperature resistance, and it produces a temperature stable, feedback resistant pantothenate kinase activity. Since the DNA sequences of these alleles are not available in the literature, the coaA genes from the two mutant strains were cloned by PCR and sequenced, in addition to a coaA gene from a strain that is wild type at the coaA locus, MM294. The PCR primer at the 5' end was designed to include the start codon plus four bases upstream, and added an arbitrarily chosen ribosome binding site (RBS). The three PCR generated fragments were each ligated between the XbaI and BamHI sites of pAN229 to give pAN284 (from coaA15(Ts)), pAN285 (from wild type coaA), and pAN286 (from coaA16(Fr)). pAN229 is a low copy E. coli vector that provides expression from the P15 promoter and that can integrate by single crossover at bpr in B. subtilis with tetracycline selection.

All three plasmids were transformed into the *E. coli* strain YH1 for complementation testing. All three plasmids complemented the temperature sensitive *coaA* mutation in *E. coli* YH1. It is presumed that the *coaA15(Ts)* gene in pAN284 is probably significantly overexpressed relative to the normal chromosomal gene, such that the overproduction compensates for the temperature sensitive defect. Complementation of a defect by overproduction is a well-documented phenomenon in *E. coli*.

The coal coding regions from pAN284, 285, and 286 were subcloned into pGEM7 to give pAN306, 307, and 308, respectively, for DNA sequencing. As expected, the DNA sequence of the insert in pAN307 (from wild type coal) matched the coal sequence from the E. coli genome database (GenBankTM). The sequence from pAN306 contains a single base change that causes a S176L substitution (i.e., a Ser \rightarrow

10

15

Leu substitution in the amino acid sequence set forth as SEQ ID NO:2). Interestingly, the DNA sequence of the pAN308 insert, derived from the feedback resistant strain, was identical to that derived from its temperature sensitive parent (represented in pAN306). This is in accord with the genetic data that indicates that the reversion of the temperature sensitive mutation occurred at a second site unlinked to the *coaA* gene.

The S176L mutation, predicted to cause the temperature sensitive defect in *E. coli* pantothenate kinase, changed a serine residue that is conserved in all known or suspected bacterial *coaA* encoded pantothenate kinases, including that of *B. subtilis* (see SEQ ID NO:3 and refer to alignment). Based on this, a serine to leucine change at the homologous residue in the *B. subtilis* pantothenate kinase is predicted to result in either a temperature sensitive enzyme or one which is less active. Accordingly, to produce a mutant *B. subtilis coaA* gene, this specific change was introduced into the *B. subtilis coaA* gene. The mutant version is installed in the chromosome of a *B. subtilis* strain deleted for *coaX*, for example, and the recombinant microorganism is checked for temperature sensitivity (*e.g.*, reduced growth at 43°C). The mutation is then installed into a pantothenate overproducing strain, preferably a strain deleted for the above mentioned *coaX* gene by standard methods to give strains favorable for pantothenate production in *B. subtilis*, *i.e.*, a strain that has reduced pantothenate kinase activity under typical fermentation conditions.

20

Additional coaA point mutations resulting in reduced pantothenate kinase activity

Of course it is expected that many other point mutations or combinations of more than one point mutation in B. subtilis coaA will also lead to reduced activity.

Appropriate mutations can be generated by mutagenic polymerase chain reaction and in vitro recombination, and identified by screening for alleles that poorly complement the E. coli coaA15(Ts) mutant. An example of such a mutation of this type is a tyrosine to histidine substitution at amino acid 181 of B. subtilis coaA, generated by mutagenic polymerase chain reaction (see SEQ ID NO:3 and first line of the alignment of Figure 24).

Isolate pAN282A was derived from the middle-sized B. subtilis coaA open reading frame described in Example XII. pAN282A complemented the E. coli coaA15(Ts) mutant very poorly, but nonetheless at a level that was detectable above background. As was done for the E. coli coaA clones, the open reading frame from pAN282A was subcloned into pGEM7 to give pAN303. The DNA sequence of the insert in pAN303 showed a single base change that led to a tyrosine to histidine amino

30

20

30

acid change at the tyrosine corresponding to Y181 of SEQ ID NO:3. This tyrosine residue is conserved in all bacterial coal genes/homologues present in GenBank (Figure 24). This tyrosine residue and the serine that is altered in the E. coli temperature sensitive pantothenate kinase described above are separated by only three amino acid residues in a region which is highly conserved in bacterial pantothenate kinases whereas the DNA sequence of a second isolate of the middle-sized open reading frame, from pAN282B, was identical to the wild type sequence from the B. subtilis genome sequencing project. The single base change found in pAN303 probably occurred during PCR amplification of the coal gene. If this variant of coal has sufficient residual biological activity in B. subtilis, it may be useful in the future for providing reduced pantothenate kinase activity.

A preferred plasmid that can serve as a basis for mutagenizing the *coaA* open reading frame is pAN294 (see *e.g.*, Figure 25 and Example XII). Briefly, mutagenic PCR is performed using pAN294 as a template and variants of *coaA* having reduced pantothenate kinase activity are screened as described above. Alternatively, mutations such as the one isolated in pAN282A can be installed into pAN294. The desired mutation is then introduced into the chromosome of a *B. subtilis* strain by transformation with the appropriate pAN294 derivative and selected for chloramphenicol resistance at 5 mg/L. Among the resulting transformants will be isolates that contain the desired mutation.

In a similar fashion, mutations that reduce the activity of the CoaX enzyme can be generated and identified, and such mutations used for optimizing pantothenate production by reducing CoA production as described above.

25 EXAMPLE XVI: Deleting the second pantothenate kinase gene, coaX gene from B. subtilis

With the knowledge gained above concerning the existence and nature of coaX, one can create a deletion of the coaX open reading frame from the B. subtilis chromosome that will remove the encoded activity, and that will not adversely affect the expression of the genes downstream from coaX. In such a deleted strain, the coaA gene will be the only gene that encodes pantothenate kinase.

To delete the coaX gene from B. subtilis, plasmid pAN336 (SEQ ID NO:92), which contains upstream and downstream homology for double crossover, was constructed with a kanamycin resistance gene replacing most of the coaX ORF (Figure 26). Strain PY79 was transformed to kanamycin resistance by pAN336, and an isolate confirmed to have resulted from a double crossover by PCR was named PA876. As predicted, deletion of coaX by itself is not lethal for B. subtilis. Furthermore, chromosomal DNA from PA876 would not transform competent PA861 (PY79 AcoaA::cat) to kanamycin resistance. These results indicate that it is the combination of AcoaA::cat and AcoaX:: kan that is lethal for B. subtilis, confirming that B. subtilis

10 contains two unlinked genes that encode pantothenate kanase, coaA and coaX, and that either gene alone is capable of supplying sufficient pantothenate kinase for a normal rate of growth.

EXAMPLE XVII: Construction of a plasmid designed to allow directed mutagenesis of the *B. subtilis coaA* gene

In order to easily introduce mutated coaA genes into the B. subtilis chromosome, it was necessary to install an antibiotic resistance gene adjacent to the coaA gene. This was accomplished by joining together in the vector pGEM5 three DNA fragments: (1) a 3.4 kb DNA sequence containing 2.5 kb of genomic sequence upstream from coal and the coal open reading frame(s); (2) a 1.1 kb DNA sequence containing a 20 chloramphenicol resistance gene (cat); and (3) a 1.4 kb DNA sequence comprising a region downstream from the operon that contains coaA. The resulting plasmid, named pAN294, effectively replaces the open reading frame yqjT (the open reading frame just downstream from coaA) with the cat gene, with enough homology flanking both sides of the cat gene to allow double recombination into the B. subtilis chromosome (Figure 25). 25 pAN294 was transformed into B. subtilis strain PY79, selecting for chloramphenicol resistance at 5 mg/l to give strains PA836 and PA837, which are presumably identical. PA836 and 837 were checked by diagnostic PCR to show that the cat gene had integrated by double crossover, as opposed to single crossover. PA836 and PA837-grow normally, leading to the conclusion that the open reading frame yqjT is not essential 30 (i.e., the yajT open reading frame could be deleted from strains PA836 and PA837 with no significant effect on growth or pantothenate production). Thus, variant alleles (i.e., mutations) of the coaA gane can be introduced into pAN294 and the resulting plasmids can be used to introduce the variant alleles into the chromosome of, for example, a B. 35 subtilis strain.

EXAMPLE XVIII: Generation of mutant coaX genes encoding pantothenate kinase having reduced or temperature sensitive activities

Mutant coaX genes are generated by introducing point mutations into the gene and testing the resulting mutants for the ability to complement the E. coli YH1 strain as described in Example XII. Preferred mutations in the coaX gene sequences are those that encode a substitution of a residue conserved among CoaX sequences from a variety of bacterial sources (e.g., a conserved residue set forth in Figure 23). Alternatively, random mutations in the coaX gene sequence are generated by mutagenic PCR and in vitro recombination and identified by screening for alleles that poorly complement the E. coli coaA15(Ts) mutant.

Mutants so generated (i.e., mutants having reduced coaX activity) can be further engineered such that the endogenous coaA gene is deleted (as described in Example XIII). CoaX reduced-activity mutants can also be further engineered to contain reduced-activity CoaA gene products as described in Example XV.

15

20

25

35

10

EXAMPLE XIX: Enhanced Production of Panto-Compounds Using Bacteria Having Deletions in One or More Pantothenate Biosynthetic Enzymes

If the desired panto-compound is not pantothenate, then an appropriate deletion of one or more of the pantothenate biosynthetic genes from a pantothenate overproducing strain will provide a strain that produces said desired panto-compound. In this example, the desired panto-compound is pantoate. Starting with, for example, strain PA236, PA313 or PA824 either one or both of the panC and panD genes is deleted. In another example, ketopantoate is the desired panto-compound. Starting with, for example strain PA244, PA245 or PA824 one, two or all of the ilvC, panE1, panC and panD genes are deleted from the starting strain. If β-alanine is the desired panto-compound, then panB and panC can be deleted, preferably in a fashion that leaves an in frame fusion of a small portion of the 5' end of panB with a small portion of the 3' end of panC, from the strain PA221, PA235, PA245, or PA313. In all of the abovementioned examples, the panto-compound producing strain will be a pantothenate auxotroph. Accordingly, the growth medium requires sufficient pantothenate for adequate growth. Vectors designed to overexpress panD as described above are then transformed into the above strains to further enhance β-alanine production.

The above-mentioned deletions are accomplished by methods well-known to those skilled in the art, for example, by insertion of an antibiotic resistance gene and removing sufficient sequence from the target gene(s) to inactivate said target gene(s).

Alternatively, removal of targeted sequences is accomplished without simultaneous introduction of an antibiotic resistance gene in said target gene and then introduced by congression (co-transformation with any other appropriate selectable DNA sequence) followed by screening for the loss of function of said target gene by replica plating.

Table 24: Strains (and corresponding phenotypes) for panto-compound production

Name	Pheno type	Drug resist.	locus	panE locus	<i>ilvD</i> locus	amyE locus	<i>bpr</i> locus	Parent
PA221	Trp-		P26panBCD			•		DT 1
PA222			P ₁₅ panBCD					RL-1
PA235			P26panBCD					
PA236			P ₂₆ panBCD	P ₂₆ panEl			•	PA221
PA327	Trp-		P26panBCD	P26panE1				PA221
PA328	Trp-		P26panBCD	P26panE1				PA235
PA340	Trp-	Spc	P26panBCD	P26panE1		P26ilvBNC		PA327
PA342	Тгр-	Spc	P26panBCD	P26panE1		P26ilvBNC		PA328
PA354	Trp-	Spc	P26panBCD	P26panE1	P26ilvD	P26ilvBNC		PA342
PA365		Spc, Tet	P26panBCD	P26panE1	P26ilvD	P26ilvBNC	P26panD423	PA354.
PA374	Trp-	Spc	P26panBCD	P26panE1	P26ilvD	P26ilvBNC		PA340
PA377	Trp-	Spc, Tet	P26panBCD	P26panE1	P26ilvD	P26ilvBNC	P26panD423	PA374
PA40	Тгр-		P26panBCD			,	P26panD423	PA221
PA402	2 Trp-		P26panBCD				P26panD428	PA221
PA40:	3 Trp-		P26panBCD				P26panD429	PA221
PA40	4 Trp-	Spc, Tet	P26panBCD	P26panE1		P26ilvBNC	P26panD423	PA340
PA40	5 Trp-	Spc, Tet	P26panBCD	P26panE1		P26ilvBNC	P26panD423	
PA65	l Trp-	Spc	P26panBC*D	P26panE1				PA374
PA28	4	Spc, Tet	P26'panBCD	P26panE1	P26ilvD	P26ilvBNC	P26panD423	PA377

Equivalents Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed:

- 1. A method of producing a panto-compound comprising culturing a microorganism which overexpresses at least one *Bacillus* pantothenate biosynthetic enzyme under conditions such that the panto-compound is produced.
 - 2. The method of claim 1, wherein the microorganism overexpresses at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.
- 10 3. The method of claim 1 or 2, wherein the pantothenate biosynthetic enzyme is selected from the group consisting of ketopantoate hydroxymethyltransferase, pantothenate synthetase, aspartate-α-decarboxylase and ketopantoate reductase.
- 4. The method of any one of claims 1 to 3, wherein the microorganism overexpresses at least two pantothenate biosynthetic enzymes.
 - 5. The method of any one of claims 1 to 3, wherein the microorganism overexpresses at least three pantothenate biosynthetic enzymes.
- 20 6. The method of any one of claims 1 to 5, wherein the panto-compound is selected from the group consisting of pantothenate, pantoate, ketopantoate and β-alanine.
- 7. A method of producing a panto-compound comprising culturing a ketopantoate reductase-overexpressing (KPAR-O) microorganism under conditions such that the panto-compound is produced.
 - 8. The method of claim 7, wherein the panto-compound is pantothenate or pantoate.
 - 9. The method of claim 7 or 8, wherein the ketopantoate reductase is bacterial-derived.
- 10. The method of claim 7 or 8, wherein the ketopantoate reductase is derived from *Bacillus*.

25

30

- 11. The method of claim 7 or 8, wherein the ketopantoate reductase is derived from *Bacillus subtilis*.
- The method of any one of claims 7 to 11, wherein the KPAR-O
 microorganism further overexpresses at least one pantothenate biosynthetic enzyme in addition to overexpressing ketopantoate reductase.
- The method of claim 12, wherein the KPAR-O microorganism further overexpresses at least one of ketopantoate hydroxymethyltransferase, pantothenate
 synthetase and aspartate-α-decarboxylase.
 - 14. A method of producing pantothenate in a manner independent of precursor feed comprising culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism having a deregulated isoleucine-valine (*ilv*) pathway under conditions such that pantothenate is produced.
- 15. A method of producing at least 2 g/L pantothenate in a manner independent of aspartate or β-alanine feed comprising culturing an aspartate-α-decarboxylase-overexpressing (AαD-O) microorganism under conditions such that
 20 pantothenate is produced.
 - 16. A method of producing at least 2 g/L pantothenate in a manner independent of valine or α-ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (ilv) biosynthetic pathway under conditions such that pantothenate is produced.
 - 17. A method of producing at least 30 g/L pantothenate in a manner independent of aspartate or β -alanine feed comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that pantothenate is produced.
 - 18. A method of producing at least 30 g/L pantothenate in a manner independent of valine or α-ketoisovalerate feed comprising culturing a microorganism having a deregulated isoleucine-valine (*ilv*) biosynthetic pathway under conditions such that pantothenate is produced.

WO 01/021772 PCT/US00/25993

- 19. A β -alanine independent high yield production method for producing pantothenate comprising culturing a manipulated microorganism under conditions such that pantothenate is produced at a significantly high yield.
- 5 20. The method of any one of claims 14 to 19, wherein the microorganism overexpresses acetohydroxyacid synthetase or is transformed with a vector comprising an *ilvBN* nucleic acid sequence or an *alsS* sequence.
- 21. The method of any one of claims 14 to 19, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase or is transformed with a vector comprising an *ilvC* nucleic acid sequence.
- 22. The method of any one of claims 14 to 19, wherein the microorganism overexpresses dihydroxyacid dehydratase or is transformed with a vector comprising an 15 ilvD nucleic acid sequence.
 - 23. The method of any one of claims 19 to 22, wherein the microorganism overexpresses aspartate-α-decarboxylase or is transformed with a vector comprising a panD nucleic acid sequence.
 - 24. The method of any one of claims 14 to 23, wherein the microorganism further has a deregulated pantothenate biosynthetic pathway.
- 25. The method of any one of claims 14 to 24, wherein the microorganism further has at least one mutant gene selected from the group consisting of a mutant avtA gene, a mutant ilvE gene, a mutant ansB gene and a mutant alsD gene.
- The method of claim 24, wherein the microorganism overexpresses any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate
 synthetase and aspartate-α-decarboxylase.
 - 27. The method of claim 24 or 26, wherein the microorganism is transformed with a vector comprising a *panBCD* nucleic acid sequence or a vector comprising a *panE1* nucleic acid sequence.

PCT/US00/25993

- 28. The method of any one of claims 14 to 16 and 19 to 27, wherein pantothenate is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 5 29. The method of claim 20, wherein the microorganism overexpresses acetohydroxyacid synthetase derived from *Bacillus* or is transformed with a vector comprising an *ilvBN* nucleic acid sequence or an *alsS* nucleic acid sequence derived from *Bacillus*.
- 10 30. The method of claim 21, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase derived from Bacillus or is transformed with a vector comprising an ilvC nucleic acid sequence derived from Bacillus.
- 31. The method of claim 22, wherein the microorganism overexpresses dihydroxyacid dehydratase derived from *Bacillus* or is transformed with a vector comprising av *ilvD* nucleic acid sequence derived from *Bacillus*.
- 32. The method of claim 23, wherein the microorganism overexpresses aspartate-α-decarboxylase derived from Bacillus or is transformed with a vector
 comprising a panD nucleic acid sequence derived from Bacillus.
 - 33. The method of claim 24 or 26, wherein the microorganism overexpresses any of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate-α-decarboxylase derived from *Bacillus*.
 - 34. The method of claim 27, wherein the vector comprises a panBCD nucleic acid sequence or a panE1 nucleic acid sequence derived from Bacillus.
- 35. A method of producing a panto-compound comprising contacting a composition comprising at least one pantothenate biosynthesis pathway precursor or isoleucine-valine biosynthesis pathway precursor with at least one isolated *Bacillus* enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase, pantothenate synthetase and aspartate-α-decarboxylase, under conditions such that the panto-compound is produced.

WO 01/021772 PCT/US00/25993

- 105 -

- 36. A method of producing β -alanine comprising culturing an aspartate- α -decarboxylase-overexpressing (A α D-O) microorganism under conditions such that β -alanine is produced.
- 5 37. The method of claim 36, wherein the AαD-O microorganism has a mutation in a nucleic acid sequence encoding a pantothenate biosynthetic enzyme selected from the group consisting of ketopantoate hydroxymethyltransferase, ketopantoate reductase and pantothenate synthetase.
- 10 38. A method of producing β-alanine comprising contacting a composition comprising aspartate with an isolated *Bacillus* aspartate-α-decarboxylase enzyme under conditions such that β-alanine is produced.
- 39. A method for enhancing production of a panto-compound comprising culturing a mutant microorganism having a mutant coaX gene under conditions such that the panto-compound production is enhanced.
 - 40. The method of claim 39, wherein said recombinant microorganism has a mutant *coaA* gene.
 - 41. A method of producing a panto-compound comprising a pantothenate kinase mutant microorganism under conditions such that the panto-compound is produced at a significantly high yield.
- 25 42. The method of claim 41, wherein said mutant microorganism has a mutant coaA gene.
 - 43. The method of claim 41, wherein said mutant microorganism has a mutant *coaX* gene.
 - 44. The method of claim 41, where said mutant microorganism has a mutant coal and coal gene.
- 45. The method of any one of claims 39 to 44, wherein said panto-compound is selected from the group consisting of ketopantoate, pantoate or pantothenate.

20

- 46. The method of any one of claims 39 to 44, wherein said panto-compound is pantothenate.
- 47. The method of any one of claims 39 to 44, wherein said panto-compound is produced at a level selected from the group consisting of a level greater than 10g/L, a level greater than 20g/L and a level greater than 40g/L.
- 48. The method of any one of claims 39 to 44, wherein said recombinant microorganism further has a deregulated pantothenate biosynthetic pathway or further has a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.
 - 49. The method of claim any one of claims 39 to 44, wherein said recombinant microorganism further overexpresses panD and panE.
- 15 50. The method of any one of claims 39 to 44, wherein said recombinant microorganism further has at least one mutant gene selected from the group-consisting of a mutant avtA gene, a mutant ilvE gene, a mutant ansB gene and a mutant alsD gene.
- 51. A method for enhancing production of a panto-compound comprising culturing a microorganism that has a deregulated pantothenate biosynthetic pathway and that also has a mutation that results in reduced pantothenate kinase activity under conditions such that the panto-compound production is enhanced.
- 52. A method for identifying compounds which modulate pantothenate
 25 kinase activity comprising contacting a recombinant cell expressing pantothenate kinase
 encoded by the coaX gene with a test compound and determining the ability of the test
 compound to modulate pantothenate kinase activity in said cell.
- 53. The method of claim 52, wherein said cell further comprises a mutant coal gene encoding a pantothenate kinase having reduced activity.
 - 54. The method of any one of claims 1 to 51, wherein the microorganism is Gram positive.
- 35 55. The method of any one of claims 1 to 51, wherein the microorganism is Gram negative.

25

- 56. The method of any one of claims 1 to 51, wherein the microorganism is a microorganism belonging to a genus selected from the group consisting of *Bacillus*, *Cornyebacterium*, *Lactobacillus*, *Lactococci* and *Streptomyces*.
- 5 57. The method of any one of claims 1 to 51 and 54 to 56, wherein the microorganism is of the genus *Bacillus*.
 - 58. The method of any one of claims 1 to 51 and 54 to 57, wherein the microorganism is *Bacillus subtilis*.
 - 59. The method of any one of claims 1 to 13, 35, 39 to 51 and 54 to 58, further comprising recovering the panto-compound.
- 60. The method of any one of claims 14 to 34 and 54 to 58, further comprising recovering the pantothenate.
 - 61. The method of any one of claims 1 to 14, 35, 39 to 46, 48 to 51 and 54 to 59, wherein the panto-compound is produced at a level greater than 2 g/L.
- 20 62. A recombinant microorganism which overexpresses at least one *Bacillus* pantothenate biosynthetic enzyme.
 - 63. The recombinant microorganism of claim 62, which overexpresses at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.
 - 64. The recombinant microorganism of claim 62 or 63, wherein the pantothenate biosynthetic enzyme is selected from the group consisting of ketopantoate hydroxymethyltransferase, pantothenate synthetase, aspartate-α-decarboxylase and ketopantoate reductase.
 - 65. The recombinant microorganism of any one of claims 62 to 64, wherein the pantothenate biosynthetic enzyme is ketopantoate reductase.
- 66. A recombinant microorganism which overexpresses aspartate-α 35 decarboxylase and has a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.

PCT/US00/25993

- 67. A recombinant microorganism having a mutant coaX gene, said mutant coaX gene encoding reduced pantothenate kinase activity in said microorganism.
- 68. The recombinant microorganism of claim 67 further having a mutant
 5 coaA gene, said mutant coaA gene encoding reduced pantothenate kinase activity in said microorganism.
- 69. A recombinant microorganism having a mutant coaX gene and optionally having a mutant coaA gene, said mutant microorganism having reduced pantothenate

 10 kinase activity as compared to a microorganism having wild-type coaA and coaX genes.
 - 70. A recombinant microorganism comprising a vector comprising an isolated *coaX* gene.
- 15 71. A recombinant microorganism that overproduces a panto-compound, the microorganism having a deregulated pantothenate biosynthetic pathway and having at least one mutation that results in a decrease in the capacity of the microorganism to synthesize Coenzyme A (CoA).
- 72. The recombinant microorganism of claim 71, having at least one mutation that results in a reduced level of pantothenate kinase activity.
- 73. The recombinant microorganism of claim 72, having a mutation in a coaA gene, or homologue thereof, that results in a reduced level of CoaA enzyme 25 activity.
 - 74. The recombinant microorganism of claim 72, having a mutation in a coaX gene, or homologue thereof, that results in a reduced level of CoaX enzyme activity.
 - 75. The recombinant microorganism of claim 72, having a mutation in a coal gene, or homologue thereof, and having a mutation in a coal gene, or homologue thereof, the mutations resulting in reduced levels of Coal enzyme activity and reduced Coal enzyme activity.
 - 76. The recombinant microorganism of any one of claims 66 to 70 which further has a deregulated pantothenate biosynthetic pathway.

35

- 77. The recombinant microorganism of any one of claims 62 to 65 and 67 to 75, further having a deregulated isoleucine-valine (*ilv*) biosynthetic pathway.
- 5 78. The recombinant microorganism of any one of claims 62 to 77, which is Gram positive.
- 79. The recombinant microorganism of claim 78 belonging to a genus selected from the group consisting of Bacillus, Cornyebacterium, Lactobacillus, 10 Lactococci and Streptomyces.
 - 80. The recombinant microorganism of claim 79 belonging to the genus *Bacillus*.
 - 81. The recombinant microorganism of claim 80 which is Bacillus subtilis.
- 82. A recombinant microorganism selected from the group consisting of PA221, PA235, PA236, PA313, PA410, PA402, PA403, PA411, PA412, PA413, PA303, PA327, PA328, PA401, PA340, PA342, PA404, PA405, PA374, PA354, PA365, PA377, PA651 and PA824.
 - 83. A recombinant vector for use in the production of panto-compounds comprising a nucleic acid sequence which encodes at least one *Bacillus* pantothenate biosynthetic enzyme operably linked to regulatory sequences.
 - 84. The vector of claim 83, comprising a nucleic acid sequence which encodes at least one *Bacillus subtilis* pantothenate biosynthetic enzyme.
- 85. The vector of claim 84, wherein the nucleic acid sequence encodes at least one of ketopantoate hydroxymethyltransferase, pantothenate synthetase, aspartate-α-decarboxylase and ketopantoate reductase.
 - 86. A recombinant vector comprising at least one nucleic acid sequence selected from the group consisting of SEQ ID NO:23, SEQ ID NO:25, SEQ ID NO:27,
 5 SEQ ID NO:29 and SEQ ID NO:59.

- 87. The vector of claim 84, wherein the nucleic acid sequence encodes ketopantoate reductase.
- 88. A vector comprising a mutant *coaX* gene, said mutant encoding a pantothenate kinase enzyme having reduced activity.
 - 89. A vector comprising an isolated coaX gene.
 - 90. A vector comprising an isolated *Bacillus coaX* gene.

- 91. A vector comprising an isolated Bacillus subtilis coaX gene.
- 92. The vector of any one of claims 86 and 89 to 91, which further comprises regulatory sequences.

15

- 93. The vector of any one of claims 83 to 85, 87 and 92, wherein the regulatory sequences comprise a constitutively active promoter.
- 94. The vector of claim 93, wherein the constitutively active promoter comprises P_{veg} (SEQ ID NO:41), P_{15} (SEQ ID NO:39) or P_{26} (SEQ ID NO:40) sequences.
 - 95. The vector of claim 83, wherein the regulatory sequences comprise at least one artificial ribosome binding site (RBS).

25

96. The vector of claim 95, wherein the artificial RBS comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:55 and SEQ ID NO:57.

30

97. A vector selected from the group consisting of pAN004, pAN005, pAN006, pAN236, pAN423, pAN428, pAN429, pAN441, pAN442, pAN443, pAN251, pAN267, pAN256, pAN257, pAN263, pAN240, pAN294, pAN296, pAN336, pAN341 and pAN342.

35

98. A recombinant microorganism comprising the vector of claim 86 or 93.

PCT/US00/25993

- 99. An isolated nucleic acid molecule which encodes at least one *Bacillus* pantothenate biosynthetic gene.
- 100. The isolated nucleic acid molecule of claim 99 which encodes at least one *Bacillus subtilis* pantothenate biosynthetic gene.
 - 101. The isolated nucleic acid molecule of claim 99 or 100 which encodes ketopantoate reductase.
- 10 102. An isolated *Bacillus* pantothenate biosynthetic enzyme polypeptide.
 - 103. An isolated *Bacillus subtilis* pantothenate biosynthetic enzyme polypeptide.
- 15 104. An isolated Bacillus ketopantoate reductase polypeptide.
 - 105. An isolated Bacillus subtilis ketopantoate reductase polypeptide.
 - 106. An isolated *Bacillus* aspartate-α-decarboxylase polypeptide.
 - 107. An isolated Bacillus subtilis aspartate-α-decarboxylase polypeptide.
 - 108. An isolated nucleic acid molecule comprising a mutant coaX gene.
- 25 109. An isolated nucleic acid molecule comprising a coaX gene.
 - 110. An isolated pantothenate kinase protein encoded by a coaX gene.




FIG. 2





FIG. 3A



SUBSTITUTE SHEET (RULE 26)

FIG. 3B







FIG. 4






PCT/US00/25993



Swa I CTAGATTAGAAAGGAGGATTTAAAT ATG TAT CGT AC TAATCTTTCCTCCTAAATTTA TAC ATA GCA TG



SUBSTITUTE SHEET (RULE 26)







FIG. 8









SUBSTITUTE SHEET (RULE 26)

FIG. 11

PCT/US00/25993




FIG. 12




FIG. 13




FIG. 14




FIG. 15






FIG. 16




SUBSTITUTE SHEET (RULE 26)

FIG. 17




19/31



SUBSTITUTE SHEET (RULE 26)

FIG. 19



21/31

FIG. 20A CLUSTAL W (1.7) Multiple Sequence Alignments

ance type explicitly set to Protein ance 1: sp Q9X795 M.leprae ance 2: sp O86779 S.coelicolor ance 3: sp O53440 M.tuberculosis ance 4: sp P54556 B.subtilis ance 5: sp P44793 H.influenzae 316 aa 316 aa	3X795 M.leprae	GLGEQIDLLEVEEVYLPLARLIHLQVAARQRLFAATAEFLGEPQONPGRP 53440 M. tuberculosis GLGEQIDLLEVEEVYLPLARLIHLQVAARQRLFAATAEFLGEPQONPDRP 64779 S. coelicolor GEGDVIDLDEVRDIYLPLSRLINLYVGATDGLRGALNTFLGEQGSQSG 64779 H. influenzae GENEDLSLDEVSTIYLPLTRLINYYIDENLHRQTVLHRFLGRNNAK 61NEDLSLEEVAEIYLPLSRLINFYISSNLRRQAVLEQFLGTNGQR 61NDYLSVEEVETIYIPLVRLHHVKSAAERNKHVNVFLKHPHSAK 7	9X795 M.leprae VPFIIGVAGSVAVGKSTTARVLQALLARWDHHTRVDLVTTDGFLYPNAEL S3440 M.tuberculosis VPFIIGVAGSVAVGKSTTARVLQALLARWDHHPRVDLVTTDGFLYPNAEL B6779 S.coelicolor TPFVIGVAGSVAVGKSTVARLLQALLSRWPEHPRVELVTTDGFLPTREL TPYIISIAGSVAVGKSTSARILQSLLSHWPTERKVDLITTDGFLYPLNKL IPYIISIAGSVAVGKSTTARVLQALLSRWPEHRRVELITTDGFLHPNQVL IPYIISIAGSVAVGKSTTARVLQALLSRWPEHRRVELITTDGFLHPNQVL IPYIIGIAGSVAVGKSTTARVLQALLSRWPEHRRVELITTDGFLHPNQVL IPYIIGIAGSVAVGKSTTARVLQALLSRWPEHRRVELITTDGFLHPNQVL IPYIIGIAGSVAVGKSTTARVLQALLSRWPEHRRVSLITTDGFLHPNAEL FASTER
Sequence Sequence Sequence Sequence Sequence Sequence	sp 09X79 sp 05344 sp 05344 sp 08677 sp P4479 sp P1504	sp Q9X79 sp O5344 sp O6677 sp P4479 sp P1504	sp Q9X79 sp O5344 sp O8677 sp P4479 sp P1504

FIG. 20B

eargimsrkgfpesydrraltrfvadikagkaevtapvyshliydivpdo Kerglakkrgfpesydmhrlvkfvsdlksgvpnvtapvyshliydvipdg <u>Orrnimhrkgfpesynrralmrfvtsvksgsdyacapvyshlhydiipga</u> KODNILIOKKGFPVSYDTPKLIRFLADVKSGKSNVTAPIYSHLTYDIIPDK kkknmmsrkgfpesydvkallefindiksgkdsvkapvyshliydreegv grrnimhrkgfpesynrraimrfvtsvksgadyacapvyshlrydtipga ** **** * * * * * *** **** ***

RLVVRRPDILIVEGINVLQPALPGKDGRT-RVGLADYFDFSVYVDARTED FEVVEQADIVIIEGINVLOSPTLEDDRENPRIFVSDFFDFSIYVDAEESR KHVVRHPDILILEGLNVLQTGP-----TLMVSDLFDFSLYVDARIQD eqvvrhpdilileglnvlqtg------tlmvsdlfdfslxvdaried fdvvdkpdilileglnvlqtgnnk---td-Qtfvsdfvdfsiyvdaeekl dktvvopdilileginvlosæmyphdph-hvfvsdfvdfsiyvdapedl ************

sp|053440|M.tuberculosis

sp|Q9X795|M.leprae

spl086779|S.coelicolor sp|P44793|H.influenzae

sp|P54556|B.subtilis

SP| P15044|E.coli

ieowyvsrelamrgtafadpeshfhhysaltdskaiiaareiwrsinrpn ieowyvsrelamritafadpeshfhhyaafsdsoavvaareiwriinrpn ierwylnrfrklratafonpssyfrkytovseeealdyaritwrtinkpn <u>Lkewyikrfikfresafndpnsyfkhyaslskeeaiataskiwdeingln</u> <u>lotwy inrflkfregaftdpdsyfhnyakltkeeaiktamtlwkeinwl</u>n iftwylerfillretafonpdsyfhkfydlsdoeademaasiwesvnrpn ****** ** ** ** ** ** ** ** **

Lyenilptkfrolilrkgdghkveevlyrrv **LVENILPTRPRATLVLRKDADHSINRLRLRKL** LVENILPTRPRATLVLRKDADHSINRLRLRKL LVENVAPTRGRATLVLRKGPDHKVORLSLRKL Lnonilptreranlilkkghnhoveliklrklkonilptrerasliltksanhaveevrlrk-*** ** ** ** ** ** ** ** ** **

sp|053440|M.tuberculosis sp|P44793|H.influenzae sp|086779|S.coelicolox splP54556|B.subtilis sp|Q9X795|M.leprae sp| P15044|E.coli

sp|053440|M.tuberculosis

sp|Q9X795|M.leprae

sp|P44793|H.influenzae sp|086779|S.coelicolor

sp|P54556|B.subtilis

sp|P15044|E.coli

SUBSTITUTE SHEET (RULE 26)


sp|053440|M.tuberculosis

sp|Q9X795|M.leprae

spi086779|S.coelicolor sp|P44793|H.influenzae

sp|P54556|B.subtilis

sp|P15044|E.coli






FIG. 22




FIG. 23A

CLUSTAL W (1.7) Multiple Sequence Alignments

258 aa Seq.8: sp 051477 B.burgdorferi 262 212 aa Seq.9: sp P74045 Synechocystis 257 246 aa Seq.10: sp 025533 H.pylori 223 265 aa Seq.11: sp 067753 A.aeolicus 229 267 aa Seq.12: sp Q9RX54 D.radiodurans 262 272 aa Seq.13: WIT RCA03301 C.acetobutylicum 250 273 aa Seq.14: WIT RRC02473 R.capsulatus 258	NKRAAFMLLETRSVLKVILVIDVGNTNIVLGIYNDTKLTAEWRLS
Sequence type explicitly set to Protein Sequence format is Pearson Seq. 1: B.subtilis Coax SEQNO 9 258 aa Seq.2: dbj BAA21476.1 D.vulgaris 212 aa Seq.3: gb AAD35964.1 T.maritima 246 aa Seq.4: pir T36391 S.coelicolor 265 aa Seq.5: sp Q45338 B.pertussis 267 aa Seq.6: sp Q66282 M.tuberculosis 272 aa Seq.7: sp Q83446 T.pallidum 273 aa	B. subtilis Coak SEQIDNO_9 WIT RCA03301 C. acetobutylicum pir T36391 S. coelicolor sp 006282 M. tuberculosis WIT RRC02473 R. capsulatus dbj BAA21476.1 D. vulgaris sp Q9RK54 D. radiodurans gb AAD35964.1 T. maritima sp 083446 T. pallidum sp 051477 B. burgdorferi sp 057753 A. aeolicus sp 07753 A. aeolicus sp 025533 B. pylori sp 025533 B. pylori sp Q45338 B. pertussis

B.subtilis|Coax|SEQIDNO_9
WIT|RCA03301|C.acetobutylicum
pir|T36391|S.coelicolor
sp|O06282|M.tuberculosis
WIT|RRC02473|R.capsulatus
dbj|BAA21476.1|D.vulgaris
sp|Q9RX54|D.radiodurans
gb|AAD35964.1|T.maritima
sp|O83446|T.pallidum
sp|O51477|B.burgdorferi
sp|O67753|A.aeolicus
sp|O67753|A.aeolicus
sp|O25533|H.pylori
sp|Q45338|B.pertussis

B.subtilis | Coax | SEQIDNO_9
WIT | RCA03301 | C.acetobutylicum
pir | T36391 | S.coelicolor
sp | 006282 | M.tuberculosis
WIT | RRC02473 | R.capsulatus
db | BAA21476.1 | D.vulgaris
sp | Q9RX54 | D.radiodurans
gb | AAD35964.1 | T.maritima
sp | 083446 | T.pallidum
sp | 067753 | A.aeolicus
sp | 067753 | A.aeolicus
sp | 25533 | H.pylori
sp | 025533 | B.pertussis

FIG. 23B

TSRHKTEDEFGMILRSLFDHS----GLMFEQIDGIIISSVVPPIMFALER
TDVLRSADEYGIQVMNLFQQD----KLDPTLVEGVIISSVVPPIMFALEH
TDSRRTADELAVLLQGLMGMHPLLGDELGDGIDGIAICATVPSVLHELRE
TESEVTADELALTIDGLIG-----EDSERLTGTAALSTVPSVLHEVRI
TDHRRTADEYFVWLNTLMQLK----GLQGRISEAIISSTAPRVVFNLRV
TDPGQTTDSIGLRLLEVLRHAG----GRADVGACVASSVVPGVNPLIRR
TGVFQTEDELFSHLHPLLG-----GAP-IPRAAVLSSVAPPVGENYAL
TGVFQTEDELFSHLHPLLG-----DAMREIKGIGVASVVPTQNTVIER
PDARKTQDEYSLLIHALCERAG----VGRASLRDAFISSVVPVLTKTIAD
TNIMLRYDEVYSFFEENFDFN-----VN---K-VFISSVVPILNETFKN
DFLKLSHEEFIKEEFPKLK-------ALGISVKQSFSEKVRG
SGNAPLQTWVTDYNPKSAQLP--------LGIQKEIFYISVNEE
IHFAQNYQLFSSAKEDLKR-------LGIQKEIFYISVNEE

MCTKYFHIEPQIVG-PG-MKTGLNIKYDNPKEVGADRIVNAVAAIHLYGMIRKYFKINPLVVG-PG-IKTGINIKYDNPKEVGADRIVNAVAAHEIYKVTRRYYGDVPAVLVEPG-VKTGVPILTDHPKEVGADRIINAVAAVELYGMLDQYWPSVPHVLIEPG-VRTGIPLLVDNPKEVGADRIINAVAAVELYGLCNRYFDCRPYVVGKPG-CELPVAPRVDPGTTVGFDRLVNTVAGYDRERACERYL--YRKLLFAPGDIAIPLDNRYERPAEVGADRLVNTVAGYDREGALKRHFMIDAFAVSAEN--LPDVTVELDTPGSVGADRLCNLFGAEKYLGFSQKYFHISPIWVKAKN---GCVKWNVKNPSEVGADRLVANAVAFVHFRAVAQISGVQPVVFGPWAYEHLPVRIPEPVRAEIGTDLVANAVAAYHFRVIFSFFKIKPLFIGFDLNYDLTFNPYKSDKFLLGSDVFANLVAAYHFRVIFSFFKIKPLFIGFDLNYDLTFNPYKSDKFLLGSDVFANLVAAYHFRQTEVWRVYQPKILTLKN----FPIQVDYKTPFTLGTDRQMALAYSAKKFYGNEKALLNCYPNAKNIAG--FFHLETDYVG---LGIDRQMACLA--VN-ATLRAGGCDIRWLRAQP-LAMGLRNGYRNPDQLGADRWACMVGVLARQPS

B.subtilis|Coax|SEQIDNO_9
WIT|RCA03301|C.acetobutylicum
pir|T36391|S.coelicolor
sp|O06282|M.tuberculosis
WIT|RRC02473|R.capsulatus
dbj|BAA21476.1|D.vulgaris
sp|Q9RX54|D.radiodurans
sp|Q9RX54|D.radiodurans
sp|Q9RX54|T.pallidum
sp|O83446|T.pallidum
sp|O67753|A.aeolicus
sp|O67753|A.aeolicus
sp|Q5533|H.pylori
sp|Q45338|B.pertussis

B. subtilis|Coax|SEQIDNO_9
WIT|RCA03301|C.acetobutylicum
pir|T36391|S.coelicolor
sp|O06282|M.tuberculosis
WIT|RRC02473|R.capsulatus
dbj|BAA21476.1|D.vulgaris
sp|Q9RX54|D.radiodurans
gb|AAD35964.1|T.maritima
sp|O83446|T.pallidum
sp|O67753|A.aeolicus
sp|P74045|Synechocystis
sp|Q5533|H.pylori
sp|Q45338|B.pertussis

FIG. 23C

NP--LIVVDFGTATTYCYIDENKQYMGGAIAPGITISTEALYSRAAKLPR
RS--LIIIDFGTATTFCAVRENGDYLGGAICPGIKVSSEALFEKAAKLPR
GP--AIVVDFGTATTFDAVSARGEYIGGVIAPGIEISVEALGYKGAQLRK
KA--AIVVDFGTATTFDAVSARGEFLGGAIAPGYQVSSDAAARSAALRR
GD--LIVVDFGTATTFDVVAPDGAYIGGVIAPGVQVSSDAAARSAALRR
GD--LIVVDFGTATTFDVVAPDGAYIGGVIAPGVVSSDAAARSAALPR
GLDYAVVVDFGTATTFDVVGG-GAYLGGLICPGVUSSAGALSSRTAKLPR
KN--GIIIDMGTATTVDLVVN-GSYEGGAILPGFFMMVHSLFRGTAKLPL
SA--CVVVDCGTALTFTAVDGTGLIQGVAIAPGLRTAVQSLHTGTAQLPL
FEN-VLVVDLGTACTIFAVSRQDGILGGIINSGPLINFNSLLDNAYLIKK
KN--VVVISAGTALVIDLVLE-GKFKGGFITLGLGKKLKILSDLAEGIPE
FP--CLVVDAGSAITTDLIKE-GKHLGGCILPGLATLGDRLAALPK
NG---VVVDAGSAITTDLIKE-GKHLGGCILPGLAYKKSAKILEQ
VHPPLLVASFGTATTLDTIGFDNVFPGGLILPGPAMMRGALAYGTAHLPL


TEITRPDN---IIGKNTVSAMQSGILEGYVGQVEGIVKRÆKKAÇAKÇDLKVELIKPAY---AICKNTISSIQSGIVYRYLRQVKYLFEKLKENLPDGRRT
IEVARPRS---VIGKNTVEAMQSGIVYGFAGQVDGVVNRMARELADD--P
VELARPRS---VIGKNTVECMQAGAVFGFAGLVDGLVGRIREDVSGFSVD
VDVTKPQG---VIGTNTVACIQSGVYWGYIGLVEGIVRQIRWERDRP--ISLEVEEDS-PVIGRSTTTSINHGFIFGFAAMTEGVLAA-----ITLQAPET---AIGKNTVHALQSGLVFGYAEMYDGLLRRIRRELPGE--VEVKPADF---VVGKDTEENIRLGVVNGSVYALEGIIGRIKEVYGDLP-VPLALPDS---VVGKDTTHAVQAGVVRGTLFVIRAMIAQCQKELGCR--FPISTPNN---LLERTTSGSVNSGLFYQYKYLIEGVYRDIKQMYKK--FFPEEVEI---FLGRSTRECVLGGAYRESTEFIKSTLKLWRKVFRRK--LEMDQLTELPDRWALDTPSAIFSGVYYGVLGALQSYLQDWQKLFPGA--PFKALDSL--EVLPRSTRDAVNYGMVLSVIACIQHLAK--NQK-----ADGLVADY-----PIDTHQAIASGIAAAQAGAIVRQWLAGRQRYGQAP---

--DCIDIVDPFLTLKGLELI

-LAPLIANES--


---VIATGG----

NUM RISLVLATGG	YERNRYGSV FERNLEYORGRIKTAR FDYNKGLGA WASRSEVR CFGD ARLVPTSLLPPATVSGSS GNSIDFKFVN LYLYHRI LXLYHRI LXLXHRI
B. subtilis Coax SEQIDNO_9 WIT RCA03301 C. acetobutylicum pir T36391 S. coelicolor sp O06282 M. tuberculosis WIT RRC02473 R. capsulatus db BAA21476.1 D. vulgaris sp Q9RX54 D. radiodurans gb AAD35964.1 T. maritima sp O83446 T. pallidum sp O51477 B. burgdorferi sp O57753 A. aeolicus sp P74045 Synechocystis sp Q25533 H. pylori sp Q25533 B. pertussis	B. subtilis Coax SEQIDNO_9 WIT RCA03301 C. acetobutylicum pir T36391 S. coelicolor sp 006282 M. tuberculosis WIT RRC02473 R. capsulatus db BAA21476.1 D. vulgaris sp Q9RX54 D. radiodurans sp Q9RX54 D. radiodurans sp Q83446 T. pallidum sp O51477 B. burgdorferi sp O67753 A. aeolicus sp O67753 R. pylori sp Q25533 R. pylori sp Q25533 R. pylori



SUBSTITUTE SHEET (RULE 26)


FIG. 25



PCT/US00/25993

WO 01/021772

FIG. 26



PCT/US00/25993

-1-

SEQUENCE LISTING

<110> OMNIGENE BIOPRODUCTS <120> METHODS AND MICROORGANISMS FOR PRODUCTION OF PANTO-COMPOUNDS <130> BGI-141CPPC <140> <141> <150> USSN 09/400,494 <151> 1999-09-21 <150> USSN 60/210,072 <151> 2000-06-07 <150> USSN 60/221,836 <151> 2000-07-28 <150> USSN 60/221,836 <151> 2000-08-24 <160> 94 <170> PatentIn Ver. 2.0 <210> 1 <211> 311 <212> PRT <213> Haemophilus influenzae <400> 1 Met Glu Phe Ser Thr Gln Gln Thr Pro Phe Leu Ser Phe Asn Arg Glu 15 5 10 Gln Trp Ala Glu Leu Arg Lys Ser Val Pro Leu Lys Leu Thr Glu Gln 25 Asp Leu Lys Pro Leu Cly Phe Asn Glu Asp Leu Ser Leu Asp Glu 35 40 Val Ser Thr Ile Tyr Leu Pro Leu Thr Arg Leu Ile Asn Tyr Tyr Ile Asp Glu Asn Leu His Arg Gln Thr Val Leu His Arg Phe Leu Gly Arg 70 75 65 Asn Asn Ala Lys Thr Pro Tyr Ile Ile Ser Ile Ala Gly Ser Val Ala Val Gly Lys Ser Thr Ser Ala Arg Ile Leu Gln Ser Leu Leu Ser His 100 105 110 Trp Pro Thr Glu Arg Lys Val Asp Leu Ile Thr Thr Asp Gly Phe Leu 120 125

Tyr Pro Leu Asn Lys Leu Lys Gln Asp Asn Leu Leu Gln Lys Lys Gly Phe Pro Val Ser Tyr Asp Thr Pro Lys Leu Ile Arg Phe Leu Ala Asp 150 155 Val Lys Ser Gly Lys Ser Asn Val Thr Ala Pro Ile Tyr Ser His Leu Thr Tyr Asp Ile Ile Pro Asp Lys Phe Asp Val Val Asp Lys Pro Asp Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Thr Gly Asn Asn Lys Thr Asp Gln Thr Phe Val Ser Asp Phe Val Asp Phe Ser Ile Tyr Val Asp Ala Glu Glu Lys Leu Leu Lys Glu Trp Tyr Ile Lys Arg Phe Leu Lys Phe Arg Glu Ser Ala Phe Asn Asp Pro Asn Ser Tyr Phe Lys His Tyr Ala Ser Leu Ser Lys Glu Glu Ala Ile Ala Thr Ala Ser Lys Ile Trp Asp Glu Ile Asn Gly Leu Asn Leu Asn Gln Asn Ile Leu Pro Thr 280 Arg Glu Arg Ala Asn Leu Ile Leu Lys Lys Gly His Asn His Gln Val Glu Leu Ile Lys Leu Arg Lys 310 <210> 2 <211> 316 <212> PRT <213> Escherichia coli <400> 2 Met Ser Ile Lys Glu Gln Thr Leu Met Thr Pro Tyr Leu Gln Phe Asp Arg Asn Gln Trp Ala Ala Leu Arg Asp Ser Val Pro Met Thr Leu Ser Glu Asp Glu Ile Ala Arg Leu Lys Gly Ile Asn Glu Asp Leu Ser Leu Glu Glu Val Ala Glu Ile Tyr Leu Pro Leu Ser Arg Leu Leu Asn Phe 50 Tyr Ile Ser Ser Asn Leu Arg Arg Gln Ala Val Leu Glu Gln Phe Leu 75

Gly Thr Asn Gly Gln Arg Ile Pro Tyr Ile Ile Ser Ile Ala Gly Ser 85 90 95

Val Ala Val Gly Lys Ser Thr Thr Ala Arg Val Leu Gln Ala Leu Leu 100 , 105 110

Ser Arg Trp Pro Glu His Arg Arg Val Glu Leu Ile Thr Thr Asp Gly 115 120 125

Phe Leu His Pro Asn Gln Val Leu Lys Glu Arg Gly Leu Met Lys Lys 130 135 140

Lys Gly Phe Pro Glu Ser Tyr Asp Met His Arg Leu Val Lys Phe Val 145 150 155 160

Ser Asp Leu Lys Ser Gly Val Pro Asn Val Thr Ala Pro Val Tyr Ser 165 170 175

His Leu Ile Tyr Asp Val Ile Pro Asp Gly Asp Lys Thr Val Val Gln 180 185 190

Pro Asp Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Ser Gly Met 195 200 205

Asp Tyr Pro His Asp Pro His His Val Phe Val Ser Asp Phe Val Asp 210 215 220

Phe Ser Ile Tyr Val Asp Ala Pro Glu Asp Leu Leu Gln Thr Trp Tyr 225 230 235 240

Ile Asn Arg Phe Leu Lys Phe Arg Glu Gly Ala Phe Thr Asp Pro Asp 245 250 255

Ser Tyr Phe His Asn Tyr Ala Lys Leu Thr Lys Glu Glu Ala Ile Lys 260 270

Thr Ala Met Thr Leu Trp Lys Glu Ile Asn Trp Leu Asn Leu Lys Gln 275 280 285

Asn Ile Leu Pro Thr Arg Glu Arg Ala Ser Leu Ile Leu Thr Lys Ser 290 295 300

Ala Asn His Ala Val Glu Glu Val Arg Leu Arg Lys 305 310 315

<210> 3

<211> 319

<212> PRT

<213> Bacillus subtilis

<400> 3

Met Lys Asn Lys Glu Leu Asn Leu His Thr Leu Tyr Thr Gln His Asn 1 5 10 15

Arg Glu Ser Trp Ser Gly Phe Gly Gly His Leu Ser Ile Ala Val Ser 20 25 30

Glu Glu Glu Ala Lys Ala Vai Glu Gly Leu Asn Asp Tyr Leu Ser Val Glu Glu Val Glu Thr Ile Tyr Ile Pro Leu Val Arg Leu Leu His Leu His Val Lys Ser Ala Ala Glu Arg Asn Lys His Val Asn Val Phe Leu Lys His Pro His Ser Ala Lys Ile Pro Phe Ile Ile Gly Ile Ala Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala Arg Ile Leu Gln Lys Leu Leu Ser Arg Leu Pro Asp Arg Pro Lys Val Ser Leu Ile Thr Thr Asp 120 Gly Phe Leu Phe Pro Thr Ala Glu Leu Lys Lys Lys Asn Met Met Ser 135 Arg Lys Gly Phe Pro Glu Ser Tyr Asp Val Lys Ala Leu Leu Glu Phe 150 Leu Asn Asp Leu Lys Ser Gly Lys Asp Ser Val Lys Ala Pro Val Tyr 165 170 Ser His Leu Thr Tyr Asp Arg Glu Glu Gly Val Phe Glu Val Val Glu Gln Ala Asp Ile Val Ile Ile Glu Gly Ile Asn Val Leu Gln Ser Pro Thr Leu Glu Asp Asp Arg Glu Asn Pro Arg Ile Phe Val Ser Asp Phe 215 Phe Asp Phe Ser Ile Tyr Val Asp Ala Glu Glu Ser Arg Ile Phe Thr 230 Trp Tyr Leu Glu Arg Phe Arg Leu Leu Arg Glu Thr Ala Phe Gln Asn 250 Pro Asp Ser Tyr Phe His Lys Phe Lys Asp Leu Ser Asp Gln Glu Ala 260 Asp Glu Met Ala Ala Ser Ile Trp Glu Ser Val Asn Arg Pro Asn Leu 280 Tyr Glu Asn Ile Leu Pro Thr Lys Phe Arg Ser Asp Leu Ile Leu Arg 295 Lys Gly Asp Gly His Lys Val Glu Glu Val Leu Val Arg Arg Val

<210> 4 <211> 312 <212> PRT <213> Mycobacterium leprae

Met Pro Arg Leu Ser Glu Pro Ser Pro Tyr Val Glu Phe Asp Arg Lys 10 Gln Trp Arg Ala Leu Arg Met Ser Thr Pro Leu Ala Leu Thr Glu Glu 25 Glu Leu Ile Gly Leu Arg Gly Leu Gly Glu Gln Ile Asp Leu Leu Glu 40 Val Glu Glu Val Tyr Leu Pro Leu Ala Arg Leu Ile His Leu Gln Val Ala Ala Arg Gln Arg Leu Phe Ala Ala Thr Ala Glu Phe Leu Gly Glu Pro Gln Gln Asn Pro Gly Arg Pro Val Pro Phe Ile Ile Gly Val Ala Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala Arg Val Leu Gln Ala Leu Leu Ala Arg Trp Asp His His Thr Arg Val Asp Leu Val Thr Thr Asp Gly Phe Leu Tyr Pro Asn Ala Glu Leu Gly Arg Arg Asn Leu Met His Arg Lys Gly Phe Pro Glu Ser Tyr Asn Arg Arg Ala Leu Met Arg Phe Val Thr Ser Val Lys Ser Gly Ala Asp Tyr Ala Cys Ala Pro Val Tyr Ser His Leu Arg Tyr Asp Thr Ile Pro Gly Ala Lys His Val Val Arg His Pro Asp Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Thr 200 Gly Pro Thr Leu Met Val Ser Asp Leu Phe Asp Phe Ser Leu Tyr Val 215 Asp Ala Arg Ile Gln Asp Ile Glu Gln Trp Tyr Val Ser Arg Phe Leu Ala Met Arg Gly Thr Ala Phe Ala Asp Pro Glu Ser His Phe His His 245 Tyr Ser Ala Leu Thr Asp Ser Lys Ala Ile Ile Ala Ala Arg Glu Ile

Trp Arg Ser Ile Asn Arg Pro Asn Leu Val Glu Asn Ile Leu Pro Thr 280 285

Arg Pro Arg Ala Thr Leu Val Leu Arg Lys Asp Ala Asp His Ser Ile

0101770A2 IA-

290 .295 .300

Asn Arg Leu Arg Leu Arg Lys Leu 305 . 310

<210> 5

<211> 312

<212> PRT

<213> Mycobacterium tuberculosis

<400> 5

Met Ser Arg Leu Ser Glu Pro Ser Pro Tyr Val Glu Phe Asp Arg Arg 1 5 10 15

Gln Trp Arg Ala Leu Arg Met Ser Thr Pro Leu Ala Leu Thr Glu Glu 20 25 30

Glu Leu Val Gly Leu Arg Gly Leu Gly Glu Gln Ile Asp Leu Leu Glu 35 40 45

Val Glu Glu Val Tyr Leu Pro Leu Ala Arg Leu Ile His Leu Gln Val 50 55 60

Ala Ala Arg Gln Arg Leu Phe Ala Ala Thr Ala Glu Phe Leu Gly Glu
65 70 75 80

Pro Gln Gln Asn Pro Asp Arg Pro Val Pro Phe Ile Ile Gly Val Ala 85 90 95

Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala Arg Val Leu Gln Ala 100 105 110

Leu Leu Ala Arg Trp Asp His His Pro Arg Val Asp Leu Val Thr Thr 115 120 125

Asp Gly Phe Leu Tyr Pro Asn Ala Glu Leu Gln Arg Arg Asn Leu Met 130 140

His Arg Lys Gly Phe Pro Glu Ser Tyr Asn Arg Arg Ala Leu Met Arg 145 150 155 160

Phe Val Thr Ser Val Lys Ser Gly Ser Asp Tyr Ala Cys Ala Pro Val 165 170 175

Tyr Ser His Leu His Tyr Asp Ile Ile Pro Gly Ala Glu Gln Val Val 180 185 190

Arg His Pro Asp Ile Leu Ile Leu Glu Gly Leu Asn Val Leu Gln Thr 195 200 205

Gly Pro Thr Leu Met Val Ser Asp Leu Phe Asp Phe Ser Leu Tyr Val

Asp Ala Arg Ile Glu Asp Ile Glu Gln Trp Tyr Val Ser Arg Phe Leu 225 230 235 240

Ala Met Arg Thr Thr Ala Phe Ala Asp Pro Glu Ser His Phe His His

245 250 255

Tyr Ala Ala Phe Ser Asp Ser Gln Ala Val Val Ala Ala Arg Glu Ile 260 265 270

Trp Arg Thr Ile Asn Arg Pro Asn Leu Val Glu Asn Ile Leu Pro Thr 27.5 280 285

Arg Pro Arg Ala Thr Leu Val Leu Arg Lys Asp Ala Asp His Ser Ile 290 295 300

Asn Arg Leu Arg Leu Arg Lys Leu 305 310

<210> 6

<211> 329

<212> PRT

<213> Streptomyces coelicolor

<400> 6

Met Ile Ser Pro Val Pro Ser Ile Pro Arg Ser Ala His Arg Gln Arg
1 5 10 15

Pro Glu Ala Thr Pro Tyr Val Asp Leu Thr Arg Pro Glu Trp Ser Ala 20 25 30

Leu Arg Asp Lys Thr Pro Leu Pro Leu Thr Ala Glu Glu Val Glu Lys
35 40 45

Leu Arg Gly Leu Gly Asp Val Ile Asp Leu Asp Glu Val Arg Asp Ile 50 55 60

Tyr Leu Pro Leu Ser Arg Leu Leu Asn Leu Tyr Val Gly Ala Thr Asp 65 70 75 80

Gly Leu Arg Gly Ala Leu Asn Thr Phe Leu Gly Glu Gln Gly Ser Gln 85 90 95

Ser Gly Thr Pro Phe Val Ile Gly Val Ala Gly Ser Val Ala Val Gly
100 105 110

Lys Ser Thr Val Ala Arg Leu Leu Gln Ala Leu Leu Ser Arg Trp Pro 115 120 125

Glu His Pro Arg Val Glu Leu Val Thr Thr Asp Gly Phe Leu Leu Pro 130 135 140

Thr Arg Glu Leu Glu Ala Arg Gly Leu Met Ser Arg Lys Gly Phe Pro 145 150 155 160

Glu Ser Tyr Asp Arg Arg Ala Leu Thr Arg Phe Val Ala Asp Ile Lys
165 170 175

Ala Gly Lys Ala Glu Val Thr Ala Pro Val Tyr Ser His Leu Ile Tyr 180 185 190

Asp Ile Val Pro Asp Gln Arg Leu Val Val Arg Arg Pro Asp Ile Leu

MICROCID. AND 010177080 IA-

		195					200					205			
Ile	Val 210	Glu	Gly	Leu '		Val 215	Leu '	Gln	Pro	Ala.	Leu 220	Pro	Gly	Lys	Asp
Gly 225	Arg	Thr	Arg	Val	Gly 230	Leu	Ala	Asp	Tyr	Phe 235	Asp	Phe	Ser	Val	Tyr 240
Val	Asp	Ala	Arg	Thr 245	Glu	Asp	Ile	Glu	Arg 250	Trp	Tyr	Leu	Asn	Arg 255	Phe
Arg	Lys	Leu	Arg 260	Ala	Thr	Ala	Phe	Gln 265	Asn	Pro	Ser	Ser	Tyr 270	Phe	Arg
Lys	Tyr	Thr 275	Gln	Val	Ser	Glu	Glu 280	Glu	Ala	Leu	Asp	Tyr 285	Ala	Arg	Thr
Thr	Trp 290	Arg	Thr	Ile	Asn	Lys 295	Pro	Asn	Leu	Val	Glu 300	Asn	Val	Ala	Pro
Thr 305	Arg	Gly	Arg	Àla	Thr 310	Leu	Val	Leu	Arg	Lys 315	Gly	Pro	Asp	His	Lys 320
Val	Gln	Arg	Leu	Ser 325	Leu	Arg	Lys	Leu							
<21 <21 <21		65 RT trep	tomy	ces	coel.	icol	or	•							
			Thr	Ile 5		Val	Gly	Asn	Thr 10		Thr	'Val	Leu	Gly 15	Leu
Phe	Asp	Gly	/ Glu	Asp	Tle	Val	_								***
			20		, 110	V G I	Glu	His 25	Trp	Arg	Ile	Ser	Thr 30) Ser
Arç	, Arg	, Thi 35	20 Ala					25 Val			*		30 Leu)	Gly
		35 Pro	20 Ala	Asp	Glu	Leu	Ala 40	25 Val	Leu	. Leu	Gln	Gly 45	30 Leu	Met	
Met	His 50	35 Pro	20 Ala Leu	Asp Leu	Glu Gly	Asp 55	Ala 40 Glu	25 Val	Leu Gly	Leu Asp	Gln Gly 60	Gly 45	30 Leu	Met	Gly
Met Ala	His 50 a Ile	35 Pro) • Cy:	20 Alas	Asp Leu	Glu Gly Val 70 Gly	Asp 55	Ala 40 Glu Ser	25 Val Leu Val	Leu Gly	Leu Asp His 75	Gln Gly 60 Glu	Gly 45 Ile	30 Leu Asr	Met Gly Glu	Gly Ile Val 80 Gly
Met Ala 65 Thi	His 50 a Ile 5	35 Pro) ≥ Cy:	20 Alasi	Asp Leu Thi Tyi 85	Glu Gly Val 70 Gly	Asp 55 Pro	Ala 40 Glu Ser	Val Lev Val	Lev Gly Lev Ala 90	Asp His 75	Gly 60 Glu Glu Leu	Gly 45 Ile Leu Val	Jeu Leu Asp Arc	Gly Glu Pro	Gly Ile Val 80 Gly

Pro Ala Ile Val Val Asp Phe Gly Thr Ala Thr Thr Phe Asp Ala Val

130 135 140 Ser Ala Arg Gly Glu Tyr Ile Gly Gly Val Ile Ala Pro Gly Ile Glu 150 , Ile Ser Val Glu Ala Leu Gly Val Lys Gly Ala Gln Leu Arg Lys Ile Glu Val Ala Arg Pro Arg Ser Val Ile Gly Lys Asn Thr Val Glu Ala Met Gln Ser Gly Ile Val Tyr Gly Phe Ala Gly Gln Val Asp Gly Val Val Asn Arg Met Ala Arg Glu Leu Ala Asp Asp Pro Asp Asp Val Thr Val Ile Ala Thr Gly Gly Leu Ala Pro Met Val Leu Gly Glu Ser Ser Val Ile Asp Glu His Glu Pro Trp Leu Thr Leu Met Gly Leu Arg Leu 250 Val Tyr Glu Arg Asn Val Ser Arg Met 260 <210> 8 <211> 272 <212> PRT <213> Mycobacterium tuberculosis <400> 8 Met Leu Leu Ala Ile Asp Val Arg Asn Thr His Thr Val Val Gly Leu Leu Ser Gly Met Lys Glu His Ala Lys Val Val Gln Gln Trp Arg Ile 25 30 Arg Thr Glu Ser Glu Val Thr Ala Asp Glu Leu Ala Leu Thr Ile Asp Gly Leu Ile Gly Glu Asp Ser Glu Arg Leu Thr Gly Thr Ala Ala Leu Ser Thr Val Pro Ser Val Leu His Glu Val Arg Ile Met Leu Asp Gln Tyr Trp Pro Ser Val Pro His Val Leu Ile Glu Pro Gly Val Arg Thr 85 Gly Ile Pro Leu Leu Val Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val Asn Cys Leu Ala Ala Tyr Asp Arg Phe Arg Lys Ala Ala Ile

120

Val Val Asp Phe Gly Ser Ser Ile Cys Val Asp Val Val Ser Ala Lys

125

115

	130					135					140				
31y 145	Glu	Phe	Leu	GĻ	Gly 150	Ala	Ile	Ala	Pro	Gly 155	'Val	Gln	Val	Šer	Ser 160
Asp	Ala	Ala	Ala	Ala 165	Arg	Ser	Ala	Ala	Leu 170	Arg	Arg	Val	Glu	Leu 175	Ala
Arg	Pro	Arg	Ser 180	Val	Val	Gly	Lys	Asn 185	Thr	Val	Glu	Суѕ	Met 190	Gln	Ala
Gly	Ala	Val 195	Phe	Gly.	Phe	Ala	Gly 200	Leu	Val	Asp	Gly	Leu 205	Val	Gly	Arg
Ile	Arg 210		Asp	Val	Ser	Gly 215	Phe	Ser	Val	Asp	His 220	Asp	Val	Ala	Ile
Val 225		Thr	Gly	His	Thr 230	Ala	Pro	Leu	Leu	Leu 235	Pro	Glu	Leu	His	Thr 240
Val	Asp	His	Tyr	Asp 245		His	Leu	Thr	Leu 250	-Gln	Gly	Leu	Arg	Leu 255	Val
Phe	Glu	Arg	Asn 260		Glu	Val	Gln	Arg 265	Gly	Arg	Leu	Lys	Thr 270	Ala	Arg
								•							

<210> 9 <211> 258 <212> PRT <213> Bacillus subtilis

Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg 20 25 30

His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp 35 40 45

His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser 50 55 60

Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr
65 70 75 80

Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu $85 \hspace{1cm} 90 \hspace{1cm} 95$

Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val 100 105 110

Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 115 120 125 Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 135 140

Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala 145 150 155 160

Leu Tyr Ser Arg Ala Ala Lys Leu Pro Arg Ile Glu Ile Thr Arg Pro 165 170 , 175

Asp Asn Ile Ile Gly Lys Asn Thr Val Ser Ala Met Gln Ser Gly Ile 180 185 190

Leu Phe Gly Tyr Val Gly Gln Val Glu Gly Ile Val Lys Arg Met Lys 195 200 205

Trp Gln Ala Lys Gln Asp Leu Lys Val Ile Ala Thr Gly Gly Leu Ala 210 215 220

Pro Leu Ile Ala Asn Glu Ser Asp Cys Ile Asp Ile Val Asp Pro Phe 225 230 235 240

Leu Thr Leu Lys Gly Leu Glu Leu Ile Tyr Glu Arg Asn Arg Val Gly 245 250 255

Ser Val

<210> 10

<211> 262

<212> PRT

<213> Deinococcus radiopugnans

<400> 10

0101770A0 IA.

Met Pro Ala Phe Pro Leu Leu Ala Val Asp Ile Gly Asn Thr Thr 1 5 10 15

Val Leu Gly Leu Ala Asp Ala Ser Gly Ala Leu Thr His Thr Trp Arg
20 25 30

Ile Arg Thr Asn Arg Glu Met Leu Pro Asp Asp Leu Ala Leu Gln Leu 35 40 45

His Gly Leu Phe Thr Leu Ala Gly Ala Pro Ile Pro Arg Ala Ala Val 50 55 60

Leu Ser Ser Val Ala Pro Pro Val Gly Glu Asn Tyr Ala Leu Ala Leu 65 70 75 80

Lys Arg His Phe Met Ile Asp Ala Phe Ala Val Ser Ala Glu Asn Leu 85 90 95

Pro Asp Val Thr Val Glu Leu Asp Thr Pro Gly Ser Val Gly Ala Asp 100 105 110

Arg Leu Cys Asn Leu Phe Gly Ala Glu Lys Tyr Leu Gly Gly Leu Asp

- 12 -

120 115 Tyr Ala Val Val Asp Phe Gly Thr Ser Thr Asn Phe Asp Val Val 135 Gly Arg Gly Arg Arg Phe Leu Gly Gly Ile Leu Ala Thr Gly Ala Gln 155 150 Val Ser Ala Asp Ala Leu Phe Ala Arg Ala Ala Lys Leu Pro Arg Ile 170 Thr Leu Gln Ala Pro Glu Thr Ala Ile Gly Lys Asn Thr Val His Ala 185 Leu Gln Ser Gly Leu Val Phe Gly Tyr Ala Glu Met Val Asp Gly Leu 200 Leu Arg Arg Ile Arg Ala Glu Leu Pro Gly Glu Ala Val Ala Val Ala Thr Gly Gly Phe Ser Arg Thr Val Gln Gly Ile Cys Gln Glu Ile Asp 235 Tyr Tyr Asp Glu Thr Leu Thr Leu Arg Gly Leu Val Glu Leu Trp Ala 245 250 Ser Arg Ser Glu Val Arg 260 <210> 11 <211> 212 <212> PRT <213> Desulfovibrio vulgaris <400> 11 Met Thr Gln His Phe Leu Leu Phe Asp Ile Gly Asn Thr Asn Val Lys Ile Gly Ile Ala Val Glu Thr Ala Val Leu Thr Ser Tyr Val Leu Pro Thr Asp Pro Gly Gln Thr Thr Asp Ser Ile Gly Leu Arg Leu Leu Glu Val Leu Arg His Ala Gly Leu Gly Pro Ala Asp Val Gly Ala Cys Val Ala Ser Ser Val Val Pro Gly Val Asn Pro Leu Ile Arg Arg Ala Cys Glu Arg Tyr Leu Tyr Arg Lys Leu Leu Phe Ala Pro Gly Asp Ile Ala Ile Pro Leu Asp Asn Arg Tyr Glu Arg Pro Ala Glu Val Gly Ala Asp 105 Arg Leu Val Ala Ala Tyr Ala Ala Arg Arg Leu Tyr Pro Gly Pro Arg

- 13 -

115 120 125 Ser Leu Val Ser Val Asp Phe Gly Thr Ala Thr Thr Phe Asp Cys Val 135 Glu Gly Gly Ala Tyr Leu Gly Gly Leu Ile Cys Pro Gly Val Leu Ser Ser Ala Gly Ala Leu Ser Ser Arg Thr Ala Lys Leu Pro Arg Ile Ser 165 170 Leu Glu Val Glu Glu Asp Ser Pro Val Ile Gly Arg Ser Thr Thr 180 Ser Leu Asn His Gly Phe Ile Phe Gly Phe Ala Ala Met Thr Glu Gly 200 Val Leu Ala Ala 210 <210> 12 <211> 246 <212> PRT <213> Thermotoga maritima <400> 12 Met Tyr Leu Leu Val Asp Val Gly Asn Thr His Ser Val Phe Ser Ile Thr Glu Asp Gly Lys Thr Phe Arg Arg Trp Arg Leu Ser Thr Gly Val Phe Gln Thr Glu Asp Glu Leu Phe Ser His Leu His Pro Leu Leu Gly Asp Ala Met Arg Glu Ile Lys Gly Ile Gly Val Ala Ser Val Val Pro Thr Gln Asn Thr Val Ile Glu Arg Phe Ser Gln Lys Tyr Phe His Ile Ser Pro Ile Trp Val Lys Ala Lys Asn Gly Cys Val Lys Trp Asn Val 85 Lys Asn Pro Ser Glu Val Gly Ala Asp Arg Val Ala Asn Val Val Ala Phe Val Lys Glu Tyr Gly Lys Asn Gly Ile Ile Ile Asp Met Gly Thr 115 120 Ala Thr Thr Val Asp Leu Val Val Asn Gly Ser Tyr Glu Gly Gly Ala 135 Ile Leu Pro Gly Phe Phe Met Met Val His Ser Leu Phe Arg Gly Thr 150 155 Ala Lys Leu Pro Leu Val Glu Val Lys Pro Ala Asp Phe Val Val Gly

- 14 -

165 170 Lys Asp Thr Glu Glu Asn Ile Arg Leu Gly Val Val Asn Gly Ser Val 185 Tyr Ala Leu Glu Gly Ile Ile Gly Arg Ile Lys Glu Val Tyr Gly Asp Leu Pro Val Val Leu Thr Gly Gly Gln Ser Lys Ile Val Lys Asp Met 215 Ile Lys His Glu Ile Phe Asp Glu Asp Leu Thr Ile Lys Gly Val Tyr 230 His Phe Cys Phe Gly Asp 245 <210> 13 <211> 273 <212> PRT <213> Treponema pallidum <400> 13 Met Leu Leu Ile Asp Val Gly Asn Ser His Val Val Phe Gly Ile Gln Gly Glu Asn Gly Gly Arg Val Cys Val Arg Glu Leu Phe Arg Leu Ala Pro Asp Ala Arg Lys Thr Gln Asp Glu Tyr Ser Leu Leu Ile His Ala Leu Cys Glu Arg Ala Gly Val Gly Arg Ala Ser Leu Arg Asp Ala Phe Ile Ser Ser Val Val Pro Val Leu Thr Lys Thr Ile Ala Asp Ala Val Ala Gln Ile Ser Gly Val Gln Pro Val Val Phe Gly Pro Trp Ala Tyr 85 Glu His Leu Pro Val Arg Ile Pro Glu Pro Val Arg Ala Glu Ile Gly 105 Thr Asp Leu Val Ala Asn Ala Val Ala Ala Tyr Val His Phe Arg Ser 115 Ala Cys Val Val Val Asp Cys Gly Thr Ala Leu Thr Phe Thr Ala Val 135 Asp Gly Thr Gly Leu Ile Gln Gly Val Ala Ile Ala Pro Gly Leu Arg Thr Ala Val Gln Ser Leu His Thr Gly Thr Ala Gln Leu Pro Leu Val

Pro Leu Ala Leu Pro Asp Ser Val Leu Gly Lys Asp Thr Thr His Ala

- 15 -

180 185 190 Val Gln Ala Gly Val Val Arg Gly Thr Leu Phe Val Ile Arg Ala Met 200 Ile Ala Gln Cys Gln Lys Glu Leu Gly Cys Arg Cys Ala Ala Val Ile 215 Thr Gly Gly Leu Ser Arg Leu Phe Ser Ser Glu Val Asp Phe Pro Pro 230 Ile Asp Ala Gln Leu Thr Leu Ser Gly Leu Ala His Ile Ala Arg Leu 245 250 Val Pro Thr Ser Leu Leu Pro Pro Ala Thr Val Ser Gly Ser Ser Gly 265 Asn <210> 14 <211> 262 <212> PRT <213> Borrelia burgdorferi <400> 14 Met Asn Lys Pro Leu Leu Ser Glu Leu Ile Ile Asp Ile Gly Asn Thr Ser Ile Ala Phe Ala Leu Phe Lys Asp Asn Gln Val Asn Leu Phe Ile Lys Met Lys Thr Asn Leu Met Leu Arg Tyr Asp Glu Val Tyr Ser Phe Phe Glu Glu Asn Phe Asp Phe Asn Val Asn Lys Val Phe Ile Ser Ser Val Val Pro Ile Leu Asn Glu Thr Phe Lys Asn Val Ile Phe Ser Phe Phe Lys Ile Lys Pro Leu Phe Ile Gly Phe Asp Leu Asn Tyr Asp Leu 85 90 Thr Phe Asn Pro Tyr Lys Ser Asp Lys Phe Leu Leu Gly Ser Asp Val 105 Phe Ala Asn Leu Val Ala Ala Ile Glu Asn Tyr Ser Phe Glu Asn Val 115 120 Leu Val Val Asp Leu Gly Thr Ala Cys Thr Ile Phe Ala Val Ser Arg Gln Asp Gly Ile Leu Gly Gly Ile Ile Asn Ser Gly Pro Leu Ile Asn 150 155

Phe Asn Ser Leu Leu Asp Asn Ala Tyr Leu Ile Lys Lys Phe Pro Ile

- 16 -

175 170 165 Ser Thr Pro Asn Asn Leu Leu Glu Arg Thr Thr Ser Gly Ser Val Asn Ser Gly Leu Phe Tyr Gln Tyr Lys Tyr Leu Ile Glu Gly Val Tyr Arg Asp Ile Lys Gln Met Tyr Lys Lys Phe Asn Leu Ile Ile Thr Gly 215 Gly Asn Ala Asp Leu Ile Leu Ser Leu Ile Glu Ile Glu Phe Ile Phe Asn Ile His Leu Thr Val Glu Gly Val Arg Ile Leu Gly Asn Ser Ile Asp Phe Lys Phe Val Asn 260 <210> 15 <211> 229 <212> PRT <213> Aquifex aeolicus <400> 15 Met Arg Phe Leu Thr Val Asp Val Gly Asn Ser Ser Val Asp Ile Ala Leu Trp Glu Gly Lys Lys Val Lys Asp Phe Leu Lys Leu Ser His Glu Glu Phe Leu Lys Glu Glu Phe Pro Lys Leu Lys Ala Leu Gly Ile Ser. Val Lys Gln Ser Phe Ser Glu Lys Val Arg Gly Lys Ile Pro Lys Ile Lys Phe Leu Lys Lys Glu Asn Phe Pro Ile Gln Val Asp Tyr Lys Thr Pro Glu Thr Leu Gly Thr Asp Arg Val Ala Leu Ala Tyr Ser Ala Lys Lys Phe Tyr Gly Lys Asn Val Val Val Ile Ser Ala Gly Thr Ala Leu 105 Val Ile Asp Leu Val Leu Glu Gly Lys Phe Lys Gly Gly Phe Ile Thr 120 Leu Gly Leu Gly Lys Lys Leu Lys Ile Leu Ser Asp Leu Ala Glu Gly 130 Ile Pro Glu Phe Phe Pro Glu Glu Val Glu Ile Phe Leu Gly Arg Ser 150 Thr Arg Glu Cys Val Leu Gly Gly Ala Tyr Arg Glu Ser Thr Glu Phe

- 17 -

165 170 175 Ile Lys Ser Thr Leu Lys Leu Trp Arg Lys Val Phe Lys Arg Lys Phe 185 Lys Val Val Ile Thr Gly Gly Glu Gly Lys Tyr Phe Ser Lys Phe Gly 200 Ile Tyr Asp Pro Leu Leu Val His Arg Gly Met Arg Asn Leu Leu Tyr 215 Leu Tyr His Arg Ile <210> 16 <211> 257 <212> PRT <213> Synechocystis sp. <400> 16 Met Glu Thr Ser Lys Pro Gly Cys Gly Leu Ala Leu Asp Asn Asp Lys 10 Gln Lys Pro Trp Leu Gly Leu Met Ile Gly Asn Ser Arg Leu His Trp Ala Tyr Cys Ser Gly Asn' Ala Pro Leu Gln Thr Trp Val Thr Asp Tyr Asn Pro Lys Ser Ala Gln Leu Pro Val Leu Leu Gly Lys Val Pro Leu Met Leu Ala Ser Val Val Pro Glu Gln Thr Glu Val Trp Arg Val Tyr Gln Pro Lys Ile Leu Thr Leu Lys Asn Leu Pro Leu Val Asn Leu Tyr 85 Pro Ser Phe Gly Ile Asp Arg Ala Leu Ala Gly Leu Gly Thr Gly Leu Thr Tyr Gly Phe Pro Cys Leu Val Val Asp Gly Gly Thr Ala Leu Thr 120 Ile Thr Gly Phe Asp Gln Asp Lys Lys Leu Val Gly Gly Ala Ile Leu 135 Pro Gly Leu Gly Leu Gln Leu Ala Thr Leu Gly Asp Arg Leu Ala Ala 155 Leu Pro Lys Leu Glu Met Asp Gln Leu Thr Glu Leu Pro Asp Arg Trp 170 Ala Leu Asp Thr Pro Ser Ala Ile Phe Ser Gly Val Val Tyr Gly Val 180

185

Leu Gly Ala Leu Gln Ser Tyr Leu Gln Asp Trp Gln Lys Leu Phe Pro

190

200 195 Gly Ala Ala Met Val Ile Thr Gly Gly Asp Gly Lys Ile Leu His Gly 215 Phe Leu Lys Glu His Ser Pro Asn Leu Ser Val Ala Trp Asp Asn 235 Leu Ile Phe Leu Gly Met Ala Ala Ile His His Gly Asp Arg Pro Ile 250 Cys <210> 17 <211> 223 <212> PRT <213> Helicobacter pylori <400> 17 Met Pro Ala Arg Gln Ser Phe Thr Asp Leu Lys Asn Leu Val Leu Cys Asp Ile Gly Asn Thr Arg Ile His Phe Ala Gln Asn Tyr Gln Leu Phe 25 Ser Ser Ala Lys Glu Asp Leu Lys Arg Leu Gly Ile Gln Lys Glu Ile Phe Tyr Ile Ser Val Asn Glu Glu Asn Glu Lys Ala Leu Leu Asn Cys Tyr Pro Asn Ala Lys Asn Ile Ala Gly Phe Phe His Leu Glu Thr Asp Tyr Val Gly Leu Gly Ile Asp Arg Gln Met Ala Cys Leu Ala Val Asn 90 Asn Gly Val Val Val Asp Ala Gly Ser Ala Ile Thr Ile Asp Leu Ile Lys Glu Gly Lys His Leu Gly Gly Cys Ile Leu Pro Gly Leu Ala Gln Tyr Ile His Ala Tyr Lys Lys Ser Ala Lys Ile Leu Glu Gln Pro Phe 130 Lys Ala Leu Asp Ser Leu Glu Val Leu Pro Lys Ser Thr Arg Asp Ala 155 Val Asn Tyr Gly Met Val Leu Ser Val Ile Ala Cys Ile Gln His Leu 170 Ala Lys Asn Gln Lys Ile Tyr Leu Cys Gly Gly Asp Ala Lys Tyr Leu

185

Ser Ala Phe Leu Pro His Ser Val Cys Lys Glu Arg Leu Val Phe Asp

- 19 -

195 200 205

Gly Met Glu Ile Ala Leu Lys Lys Ala Gly Ile Leu Glu Cys Lys 210 215 220

<210> 18

<211> 267

<212> PRT

<213> Bordetella pertussis

<400> 18

Met Ile Ile Leu Ile Asp Ser Gly Asn Ser Arg Leu Lys Val Gly Trp

1 10 15

Phe Asp Pro Asp Ala Pro Gln Ala Ala Arg Glu Pro Ala Pro Val Ala
20 25 30

Phe Asp Asn Leu Asp Leu Asp Ala Leu Gly Arg Trp Leu Ala Thr Leu 35 40 45

Pro Arg Arg Pro Gln Arg Ala Leu Gly Val Asn Val Ala Gly Leu Ala 50 55 60

Arg Gly Glu Ala Ile Ala Ala Thr Leu Arg Ala Gly Gly Cys Asp Ile 65 70 75 80

Arg Trp Leu Arg Ala Gln Pro Leu Ala Met Gly Leu Arg Asn Gly Tyr 85 90 95

Arg Asn Pro Asp Gln Leu Gly Ala Asp Arg Trp Ala Cys Met Val Gly
100 105

Val Leu Ala Arg Gln Pro Ser Val His Pro Pro Leu Leu Val Ala Ser 115 120 125

Phe Gly Thr Ala Thr Thr Leu Asp Thr Ile Gly Pro Asp Asn Val Phe 130 135 140

Pro Gly Gly Leu Ile Leu Pro Gly Pro Ala Met Met Arg Gly Ala Leu 145 150 155 160

Ala Tyr Gly Thr Ala His Leu Pro Leu Ala Asp Gly Leu Val Ala Asp 165 170 175

Tyr Pro Ile Asp Thr His Gln Ala Ile Ala Ser Gly Ile Ala Ala Ala 180 185 190

Gln Ala Gly Ala Ile Val Arg Gln Trp Leu Ala Gly Arg Gln Arg Tyr 195 200 205

Gly Gln Ala Pro Glu Ile Tyr Val Ala Gly Gly Gly Trp Pro Glu \dot{V} al 210 215 220

Arg Gln Glu Ala Glu Arg Leu Leu Ala Val Thr Gly Ala Ala Phe Gly 225 230 235 240

Ala Thr Pro Gln Pro Thr Tyr Leu Asp Ser Pro Val Leu Asp Gly Leu

245 250 255

Ala Ala Leu Ala Ala Gln Gly Ala Pro Thr Ala 260 265

<210> 19 <211> 777 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(774) <400> 19 ttg tta ctg gtt atc gat gtg ggg aac acc aat act gta ctt ggt gta 4.8 Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val 10 1 tat cat gat gga aaa tta gaa tat cac tgg cgt ata gaa aca agc agg 96 Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg 20 cat aaa aca gaa gat gag ttt ggg atg att ttg cgc tcc tta ttt gat His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp 40 35 cac tcc ggg ctt atg ttt gaa cag ata gat ggc att att att tcg tca His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser 50 gta gtg ccg cca atc atg ttt gcg tta gaa aga atg tgc aca aaa tac 240 Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr 65 . ttt cat atc gag cct caa att gtt ggt cca ggt atg aaa acc ggt tta 288 Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu 85 aat ata aaa tat gac aat ccg aaa gaa gta ggg gca gac aga atc gta 336 Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val 100 aat gct gtc gct gcg ata cac ttg tac ggc aat cca tta att gtt gtc 384 Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 120 115 gat ttc gga acc gcc aca acg tac tgc tat att gat gaa aac aaa caa 432. Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 135 130 480 tac atg ggc ggg gcg att gcc cct ggg att aca att tcg aca gag gcg Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala 155 150 145 ett tac teg egt gea gea aag ett eet egt ate gaa ate ace egg eee Leu Tyr Ser Arg Ala Ala Lys Leu Pro Arg Ile Glu Ile Thr Arg Pro

- 21 -

				165					170					175	•	
gac Asp	aat Asn	att Ile	atc Ile 180	Gly	aa'a Lys	aac Asn	act Thr	gtt Val 185	agc Ser	gcg Ala	atg Met	caa Gln	tct Ser 190	gga Gly	att Ile	576
tta Leu	ttt Phe	ggc Gly 195	tat Tyr	gtc Val	ggc ggc	caa Gln	gtg Val 200	gaa Glu	gga Gly	atc Ile	gtt Val	aag Lys 205	cga Arg	atg Met	aaa Lys	624
tgg Trp	cag Gln 210	gca Ala	aaa Lys	cag Gln	gac Asp	ctc Leu 215	aag Lys	gtc Val	att Ile	gcg Ala	aca Thr 220	gga Gly	ggc Gly	ctg Leu	gcg Ala	672
ccg Pro 225	ctc Leu	att Ile	gcg Ala	aac Asn	gaa Glu 230	tca Ser	gat Asp	tgt Cys	ata Ile	gac Asp 235	atc Ile	gtt Val	gat Asp	cca Pro	ttc Phe 240	720
tta Leu	acc Thr	cta Leu	aaa Lys	ggg Gly 245	ctg Leu	gaa Glu	ttg Leu	att Ile	tat Tyr 250	gaa Glu	aga Arg	aac Asn	cgc Arg	gta Val 255	gga Gly	768
agt Ser	gta Val	tag														777
<211 <212)> 20 L> 96 ?> DN B> Ba	00 NA	lus s	subti	ilis											
	l> CI		(957))												
gtg)> 20 aaa Lys	aat	aaa Lys	gaa Glu 5	ctt Leu	aac Asn	cta Leu	cat His	act Thr 10	tta Leu	tat Tyr	aca Thr	cag Gln	cac His 15	aat Asn	-48
cgg Arg	gag Glu	tct Ser	tgg Trp 20	tct Ser	ggt Gly	ttt Phe	G1y ggg	ggg Gly 25	cat His	ttg Leu	tcg Ser	att Ile	gct Ala 30	gta Val	tct Ser	96
gaa Glu	gaa Glu	gag Glu 35	gca Ala	aaa Lys	gct Ala	gtg Val	gaa Glu 40	gga Gly	ttg Leu	aat Asn	gat Asp	tat Tyr 45	cta Leu	tct Ser	gtt Val	144
gaa Glu	gaa Glu 50	gtg Val	gag Glu	acg Thr	atc Ile	tat Tyr 55	att Ile	ccg Pro	ctt Leu	gtt Val	cgc Arg 60	ttg Leu	ctt Leu	cat His	tta Leu	192
cat His 65	gtc Val	aag Lys	tct Ser	gcg Ala	gct Ala 70	gaa Glu	cgc Arg	aat Asn	aag Lys	cat His 75	gtc Val	aat Asn	gtt Val	ttt Phe	ttg Leu 80	240
aag Lys	cac His	cca Pro	cat His	tca Ser	gcc Ala	aaa Lys	att Ile	ccg Pro	ttt Phe	att Ile	atc Ile	ggc Glv	att Ile	gcc Ala	ggc Glv	288

				85					90					95		
agt (gtc Val	Ala	gtc Val 100	gga Gly	aaa Lys	agċ Ser	Thr	acg Thr 105	gcg Ala	cgg Arg	atc Ile	ttg Leu	cag Gln 110	aag Lys	ctg Leu	336
ctt Leu	tcg Ser	cgt Arg 115	ttg Leu	cct Pro	gac Asp	cgt Arg	cca Pro 120	aaa L <u>y</u> s	gtg Val	agc Ser	ctt Leu	atc Ile 125	acg Thr	aca Thr	gat Asp	384
ggt Gly	ttt Phe 130	tta Leu	ttt Phe	cct Pro	act Thr	gcc Ala 135	gag Glu	ctg Leu	aaa Lys	aag Lys	aaa Lys 140	aat Asn	atg Met	atg Met	tca Ser	432
aga Arg 145	aaa Lys	gga Gly	ttt Phe	cct Pro	gaa Glu 150	agc Ser	tat Tyr	gat Asp	gta Val	aag Lys 155	gcg Ala	ctg Leu	ctc Leu	gaa Glu	ttt Phe 160	480
ttg Leu	aat Asn	gac Asp	tta Leu	aaa Lys 165	tca Ser	gga Gly	aag Lys	gac Asp	agc Ser 170	gta Val	aag Lys	gcc Ala	ccg Pro	gtg Val 175	tat Tyr	528
tcc Ser	cat His	cta Leu	acc Thr 180	tat Tyr	gac Asp	cgc Arg	gag Glu	gaa Glu 185	ggt Gly	gtg Val	ttc Phe	gag Glu	gtt Val 190	gta Val	gaa Glu	576
cag Gln	gcg Ala	gat Asp 195	att Ile	gtg Val	att Ile	att Ile	gaa Glu 200	ggc	att Ile	aat Asn	gtt Val	ctt Leu 205	cag Gln	tcg Ser	ccc Pro	624
acc Thr	ttg Leu 210	Glu	gat Asp	gac Asp	cgg	gaa Glu 215	aac Asn	.ccg Pro	cgt Arg	att Ile	ttt Phe 220	gtt Val	tcc Ser	gat Asp	ttc Phe	672
ttt Phe 225	gat Asp	ttt Phe	tcg Ser	att Ile	tat Tyr 230	Val	gat Asp	gcg Ala	gag Glu	gaa Glu 235	Ser	cgg Arg	att Ile	t to Phe	act Thr 240	720
tgg Trp	tat Tyr	tta Leu	gag Glu	cgt Arg 245	ttt Phe	cgc Arg	-ctg Leu	ctt Leu	cgg Arg 250	Glu	aca Thr	gct Ala	ttt Phe	Caa Glr 255	a aat n Asn	768
cct Pro	gat Asp	tca Ser	tat Tyr 260	Phe	cat His	aaa Lys	ttt Phe	aaa Lys 265	: Asp	tto Leu	tco Ser	gat Asp	cag Glr 270	'Gl	g gct ı Ala	816
gac Asp	gag Glu	atg Met 275	Ala	gcc Ala	tco Sei	att Ile	tgg Trp 280	-Glu	g agt i Sez	gto Val	aac L Asr	c cgg Arg 285	Pro	g aat o Asi	t tta n Leu	864
tat Tyr	gaa Glu 290	ı Asr	att Ile	ttg Leu	cca Pro	a act Thr 295	Lys	tto Phe	ago Aro	j tca j Sei	a gat c Asp 300	Leu	att 1116	tte Le	g cgt u Arg	912
aag Lys 305	Gl	a gad y Asp	G13	g cat / His	aaq 5 Ly:	s Val	gaç Glu	g gaa u Glu	a gto u Val	tte Lev 31:	u Val	a ago	g ago	g gt g Va	a tga 1	960

<210 <211 <212 <213	> 88 > DN	2 A	.us s	ubti	.lis							•			
<220 <221 <222	> CI		(879)												•
<400	> 21														
										gct Ala					48
										atc Ile					96
										gct Ala					144
										gcc Ala 60					192
										aaa Lys					240
		_		_	_		_	-	_	gac Asp	_				288
_										act Thr	-		_		336
_			-	_		_				gaa Glu	_		-	-	384
										tca Ser 140					432
										gac Asp					480
										att Ile					528
									Asp	cgg Arg			Pro		576

Ile Phe Val Ser Asp Phe Phe Asp Phe Ser Ile Tyr Val 1 195 200 205	gat gcg gag 624 Asp Ala Glu	
gaa agc cgg att ttc act tgg tat tta gag cgt ttt cgc (Glu Ser Arg Ile Phe Thr Trp Tyr Leu Glu Arg Phe Arg 210 215 220	Leu Leu Arg	
gaa aca gct ttt caa aat cct gat tca tat ttt cat aaa : Glu Thr Ala Phe Gln Asn Pro Asp Ser Tyr Phe His Lys 225 230 235	ttt aaa gac 720 Phe Lys Asp 240	
ttg tcc gat cag gag gct gac gag atg gca gcc tcg att Leu Ser Asp Gln Glu Ala Asp Glu Met Ala Ala Ser Ile 245 . 250	tgg gag agt 768 Trp Glu Ser 255	
gtc aac cgg ccg aat tta tat gaa aat att ttg cca act Val Asn Arg Pro Asn Leu Tyr Glu Asn Ile Leu Pro Thr 260 265	aaa ttc agg 816 Lys Phe Arg 270	
tca gat ctc att ttg cgt aag gga gac ggg cat aag gtc Ser Asp Leu Ile Leu Arg Lys Gly Asp Gly His Lys Val 275 280 285	gag gaa gtg 864 Glu Glu Val	3
ttg gta agg agg gta tga Leu Val Arg Arg Val 290	882	<u> </u>
<210> 22 <211> 846 <212> DNA <213> Bacillus subtilis		1
	•	
<220> <221> CDS <222> (1)(843)		
<221> CDS	gag acg atc 48 Glu Thr Ile 15	
<221> CDS <222> (1)(843) <400> 22 gtg gaa gga ttg aat gat tat cta tct gtt gaa gaa gtg Met Glu Gly Leu Asn Asp Tyr Leu Ser Val Glu Glu Val	tet geg get 96	
<pre><221> CDS <222> (1)(843) <400> 22 gtg gaa gga ttg aat gat tat cta tct gtt gaa gaa gtg Met Glu Gly Leu Asn Asp Tyr Leu Ser Val Glu Glu Val</pre>	tct gcg gct 96 Ser Ala Ala 30 a cat tca gcc 14 His Ser Ala	5
<pre><221> CDS <222> (1)(843) <400> 22 gtg gaa gga ttg aat gat tat cta tct gtt gaa gaa gtg Met Glu Gly Leu Asn Asp Tyr Leu Ser Val Glu Glu Val</pre>	g tct gcg gct 96 Ser Ala Ala 30 a cat tca gcc 14 His Ser Ala 5	14

- 25 -

65					70					75					80	
	cca Pro															288
	gag Glu															336
	tat Tyr															384
	aag Lys 130															432
	gag Glu															480
	gaa Glu															528
	aac Asn															576
	gat Asp															624
	ctg Leu 210															672
aaa Lys 225	ttt Phe	aaa Lys	gac Asp	ttg Leu	tcc Ser 230	gat Asp	cag Gln	gag Glu	gct Ala	gac Asp 235	gag Glu	atg Met	gca Ala	gcc Ala	tcg Ser 240	720
	tgg Trp															768
act Thr	aaa Lys	ttc Phe	agg Arg 260	tca Ser	gat Asp	ctc Leu	att Ile	ttg Leu 265	cgt Arg	aag Lys	gga Gly	gac Asp	ggg Gly 270	cat His	aag Lys	816
	gag Glu								tga							846

<210> 23 <211> 831

<212> DNA

<213	> Ba	cill	us s	ubti	lis	f								•		
	L> CD	s .)(831)	•												•
atσ)> 23 aaa Lys	aca	aaa Lys	ctg Leu 5	gat Asp	ttt Phe	cta Leu	aaa Lys	atg Met 10	aag Lys	gag Glu	tct Ser	gaa Glu	gaa Glu 15	ecg Pro	48
att Ile	gtc Val	atg Met	ctg Leu 20	acc Thr	gct Ala	tat Tyr	gat Asp	tat Tyr 25	ecg Pro	gca Ala	gct Ala	aaa Lys	ctt Leu 30	gct Ala	gaa Glu	96
caa Gln	gcg Ala	gga Gly 35	gtt Val	gac Asp	atg Met	att Ile	tta Leu 40	gtc Val	ggt Gly	gat Asp	tca Ser	ctt Leu 45	gga Gly	atg Met	gtc Val	144
gtc Val	ctc Leu 50	ggc Gly	ctt Leu	gat Asp	tca Ser	act Thr 55	gtc Val	ggt Gly	gtg Val	aca Thr	gtt Val 60	gcg Ala	gac Asp	atg Met	atc Ile	192
cat His 65	His	aca Thr	aaa Lys	gcc Ala	gtt Val 70	aaa Ļys	agg Arg	ggt Gly	gcg Ala	ccg Pro 75	aat Asn	acc Thr	ttt Phe	att Ile	gtg Val 80	240
aca Thr	gat Asp	atg Met	ccg Pro	ttt Phe 85	atg Met	tct Ser	tat Tyr	cac His	ctg Leu 90	tct Ser	aag Lys	gaa Glu	gat Asp	acg Thr 95	ctg Leu	288
aaa Lys	aat Asn	gca Ala	gcg Ala 100	Ala	atc Ile	gtt Val	cag Gln	gaa Glu 105	Ser	gga Gly	gct Ala	gac Asp	gca Ala 110	ctg Leu	aag Lys	336
ctt Lei	gag Glu	ggc Gly 115	gga Gly	gaa Glu	ggc Gly	gtg Val	ttt Phe 120	gaa Glu	tcc Ser	att Ile	cgc	gca Ala 125	Leu	acg Thr	ctt Leu	384
gga Gly	a ggo y Gly 130	, Ile	Pro	Val	Val	Ser	His	Leu	Gly	Leu	aca Thr 140	Pro	Gln	tca Ser	gtc Val	432
gg: Gl: 14	y Val	ctg Leu	ggc Gly	ggc Gly	tat Tyr 150	Lys	gta Val	.cag	ggc Gly	Lys 155	: Asp	gaa Glu	caa Glr	ago Ser	gcc Ala 160	480
aa. Ly:	a aaa s Lys	a tta s Leu	ata ı Ile	gaa Glu 165	Asp	agt Ser	ata Ile	aaa Lys	tgc Cys 170	Gli	a gaa 1 Glu	gca Ala	a gga	get Ala 175	atg Met	528
at Me	g cti t Lei	t gtg u Val	cto Lev 180	ı Glu	tgt Cys	gtg Val	Pro	gca Ala 189	Glu	cto Lei	c aca	a gco	a Lys	s Ile	gec Ala	576
ga Gl	g accurates	g cta r Lev 199	ı Sei	ata Ile	e Pro	g gto Val	200	Gly	a ato y Ile	gge Gl	g gct y Ala	ggt a Gly 201	y Va.	g aaa l Ly	a gcg s Ala	624

gac Asp	gga Gly 210	GIN	gtt Val	ctc Leu	gtt Val	tat Tyr 215	cat His	gat Asp	att Ile	atc Ile	ggc Gly 220	cac His	ggt Gly	gtt Val	gag Glu	672
aga Arg 225	aca Thr	cct Pro	aaa Lys	ttt Phe	gta Val 230	aag Lys	caa Gln	tat Tyr	acg Thr	cgc Arg 235	att Ile	gat Asp	gaa Glu	acc Thr	atc Ile 240	720
gaa Glu	aca Thr	gca Ala	atc Ile	agc Ser 245	Gly	tat Tyr	gtt Val	cag Gln	gat Asp 250	Val	aga Arg	cat His	cgt Arg	gct Ala 255	ttc Phe	768
cct Pro	gaa Glu	caa Gln	aag Lys 260	cat His	tcc Ser	ttt Phe	caa Gln	atg Met 265	aac Asn	cag Gln	aca Thr	gtg Val	ctt Leu 270	gac Asp	ggc Gly	816
_	tac Tyr															831
<21:	0> 2: 1> 2' 2> P! 3> Ba	77 RT	lus :	subt	ilis							1				
)> 2		_				+ 1					•				
Met 1	гуѕ	Thr	Lys	Leu 5	Asp	Phe	Leu	Lys	Met 10	Lys	Glu	Ser	Glu	Glu 15	Pro	
Ile	Vаl	Met	Leu 20	Thr	Ala	Tyr	Asp	Tyr 25	Pro	Ala	Ala	Lys	Leu 30	Ala	Glu	
Gln	Ala	Gly 35	Val	Asp	Met	Ile	Leu 40	Val	Gly	Asp	Ser	Leu 45	Gly	Met	Val	ı
Val	Leu 50	Gly	Leu	Asp	Ser	Thr 55	Val	Gly	Val	Thr	Val 60	Ala	Asp	Met	Ile	
His 65	His	Thr	Lys	Ala	Val 70	Lys	Arg	Gly	Ala	Pro 75	Asn	Thr	Phe	Ile	Val 80	
Thr	Asp	Met	Pro	Phe 85	Met	Ser	Tyr	His	Leu 90	Ser	Lys	Glu	Asp	Thr 95	Leu	
Lys	Asn	Ala	Ala 100	Ala	Ile	Val	Gln	Glu 105	Ser	Gly	Ala	Asp	Ala 110	Leu	Lys	
Leu	Glu	Gly 115	Gly	Glu	Gly	Val	Phe 120	Glu	Ser	Ile	Arg	Ala 125	Leu	Thr	Leu	
Gly	Gly 130	Ile	Pro	Val	Val	Ser 135	His	Leu	Gly	Leu	Thr 140	Pro	Gln	Ser	Val	
Gly 145	Val	Leu	Gly	Gly	Tyr 150	Lys	Val	Gln	Gly	Lys 155	Asp	Glu	Gln	Ser	Ala 160	

Lys	Lys	Leu	Ile	Glu 165	Asp	Ser	Ile :	Lys 4	Cys 170	Glu	Glu	Ala ·	Gly	Ala 175	Met	
Met	Leu	Val	Leu 180	Glu	Cys	Val	Pro	Ala 185	Glu	Leu	Thr	Ala	Lys 190	Ile	Ala	ı
Glu	Thr	Leu 195	Ser	Ile	Pro	Val	11e 200	Gly	Ile	Gly	Ala	Gly 205	Val	Lys	Ala	
Asp	Gly 210	Gln	Val	Leu	Val	Tyr 215	His	Asp	Ile	Ile	Gly 220	His	Gly	Val	Glu	
Arg 225	Thr	Pro	Lys	Phe	Val 230	Lys	Gln	Tyr	Thr	Arg 235	Ile	Asp	Glu	Thr	11e 240	
				245				•	250				1 -	255		
Pro	Glu	Gln	Lys 260		Ser	Phe	Gln	Met 265	Asn	Gln	Thr	Val	Leu 270	Asp	Gly	
Leu	Tyr	Gly 275		Lys										,	·	
<21 <21	0> 2 1> 8 2> D 3> B	58 NA	.lus	subt	ilis	• •										
	21> 0	DS (1)	(858	3)												
ato Me	00> 2 g aga t Arq 1	cac	g att	e Thi	c gat c Asp	att o Ile	tca Ser	cag Gln	cto Lev	ı Lys	gaa Glu	a gco a Ala	ata Ile	a aa e Ly 1	a caa s Gln 5	48
ta Ty:	c cat	t tca s Sea	a gaq r Gli 20	ı Gl	c aaq y Ly:	g tca s Ser	ato : Ile	gga Gly 25	, Phe	c gtt e Val	t cc	g aco	g ate Med 3	t "GI	g ttt y Phe	96
ct Le	g ca u Hi	t ga s Gl	u Gl	g ca y Hi	t tta s Le	a aco u Thi	tta Lei 40	ı Ala	a ga	c aaa p Ly:	a gc s Al	a aga a Ar	g G1	a ∙ga n •Gl	a aac u Asr	: 144 n
· ga As	p Ål	a Va	t at l Il	t at e Me	g ag t Se	t att	e Pho	t gto	g aa l As	t cc	t gc o Al 6	a Gl	a tt n Ph	e Gl	je eet Ly Pro	192
	5	•														
As	t da	a qa	t tt p Ph	t ga e Gl	a gc u Al 7	а Ту	t cc	g cg	c ga g As	t at p Il 7	e Gl	g cg u Ar	g ga g As	t go p Al	ca gct la Ala 80	a

- 29 -

ga As	t atg p Met	tat Tyr	ccc Pro 100	ggt Gly	gaa Glu	aag Lys	aat Asn	gtc Val 105	acg Thr	att Ile	cat His	gta Val	gaa Glu 110	aga Arg	cgc Arg	336
ac Th	a gac r Asp	gtg Val 115	tta Leu	tgc Cys	GJ À āāā	cgc Arg	tca Ser 120	aga Arg	gaa Glu	gga Gly	cat His	ttt Phe 125	gac Asp	G17 ggg	gtc Val	384
gc Al	g atc a Ile 130	gta Val	ctg Leu	acg Thr	aag Lys	ctt Leu 135	ttc Phe	aat Asn	cta Leu	gtc Val	aag Lys 140	ccg Pro	act Thr	cgt Arg	gcc Ala	432
ta Ty 14	t ttc r Phe 5	ggt Gly	tta Leu	aaa Lys	gat Asp 150	gcg Ala	cag Gln	cag Gln	gta Val	gct Ala 155	gtt Val	gtt Val	gat Asp	ggg Gly	tta Leu 160	480
at Il	c agc e Ser	gac Asp	ttc Phe	ttc Phe 165	atg Met	gat Asp	att Ile	gaa Glu	ttg Leu 170	gtt Val	cct Pro	gtc Val	gat Asp	acg Thr 175	gtc Val	528
ag Ar	a gag g Glu	gaa Glu	gac Asp 180	ggc Gly	tta Leu	gcc Ala	aaa Lys	agc Ser 185	tct Ser	cgc Arg	aat Asn	gta Val	tac Tyr 190	tta Leu	aca Thr	· 576
gc Al	t gag a Glu	gaa Glu 195	aga Arg	aaa Lys	gaa Glu	gcg Ala	cct Pro 200	aag Lys	ctg Leu	tat Tyr	cgg Arg	gcc Ala 205	ctt Leu	caa Gln	aca Thr	624
ag Se	t gcg r Ala 210	gaa Glu	ctt Leu	gtc Val	caa Gln	gcc Ala 215	ggt Gly	gaa Glu	aga Arg	gat Asp	cct Pro 220	gaa Glu	gcg Ala	gtg Val	ata Ile	672
аа Ьу 22	a gct s Ala 5	gca Ala	aaa Lys	gat Asp	atc Ile 230	att Ile	gaa Glu	acg Thr	act Thr	agc Ser 235	gga Gly	acc Thr	ata Ile	gac Asp	tat Tyr 240	720
gt Va	a gag l Glu	ctt Leu	tat Tyr	tcc Ser 245	tat Tyr	ccg Pro	gaa Glu	ctc Leu	gag Glu 250	cct Pro	gtg Val	aat Asn	gaa Glu	att Ile 255	gct Ala	768
gg Gl	a aag y Lys	atg Met	att Ile 260	ctc Leu	gct Ala	gtt Val	gca Ala	gtt Val 265	gct Ala	ttt Phe	tca Ser	aaa Lys	gcg Ala 270	cgt Arg	tta Leu	816
	a gat e Asp															858

<210> 26

<211> 286 <212> PRT

<213> Bacillus subtilis

<400> 26

Met Arg Gln Ile Thr Asp Ile Ser Gln Leu Lys Glu Ala Ile Lys Gln 5 10

Tyr His Ser Glu Gly Lys Ser Ile Gly Phe Val Pro Thr Met Gly Phe 20 25 30

Leu His Glu Gly His Leu Thr Leu Ala Asp Lys Ala Arg Gln Glu Asn
35 40 45

Asp Ala Val Ile Met Ser Ile Phe Val Asn Pro Ala Gln Phe Gly Pro
50 55 60

Asn Glu Asp Phe Glu Ala Tyr Pro Arg Asp Ile Glu Arg Asp Ala Ala 65 70 75 80

Leu Ala Glu Asn Ala Gly Val Asp Ile Leu Phe Thr Pro Asp Ala His

Asp Met Tyr Pro Gly Glu Lys Asn Val Thr Ile His Val, Glu Arg Arg 100 105 110

Thr Asp Val Leu Cys Gly Arg Ser Arg Glu Gly His Phe Asp Gly Val 115 120 125

Ala Ile Val Leu Thr Lys Leu Phe Asn Leu Val Lys Pro Thr Arg Ala 130 135 140

Tyr Phe Gly Leu Lys Asp Ala Gln Gln Val Ala Val Val Asp Gly Leu 145 150 160

Ile Ser Asp Phe Phe Met Asp Ile Glu Leu Val Pro Val Asp Thr Val 165 170 175

Arg Glu Glu Asp Gly Leu Ala Lys Ser Ser Arg Asn Val Tyr Leu Thr 180 185 190

Ala Glu Glu Arg Lys Glu Ala Pro Lys Leu Tyr Arg Ala Leu Gln Thr 195 200 205

Ser Ala Glu Leu Val Gln Ala Gly Glu Arg Asp Pro Glu Ala Val Ile 210 215 220

Lys Ala Ala Lys Asp Ile Ile Glu Thr Thr Ser Gly Thr Ile Asp Tyr 225 230 235 240

Val Glu Leu Tyr Ser Tyr Pro Glu Leu Glu Pro Val Asn Glu Ile Ala 245 250 255

Gly Lys Met Ile Leu Ala Val Ala Val Ala Phe Ser Lys Ala Arg Leu 260 265 270

Ile Asp Asn Ile Ile Ile Asp Ile Arg Glu Met Glu Arg Ile 275 280 285

<210> 27

<211> 381

<212> DNA

<213> Bacillus subtilis

<220> <221> CDS <222> (1)	(381)		ı		
<400> 27 atg tat cga Met Tyr Arg 1	aca atg atg Thr Met Met 5	agc ggc aaa Ser Gly Lys	ctt cac agg g Leu His Arg A 10	ca act gtt acg la Thr Val Thr 15	48
gaa gca aac Glu Ala Asn	ctg aac tat Leu Asn Tyr 20	gtg gga agc Val Gly Ser 25	att aca att g Ile Thr Ile A	at gaa gat ctc sp Glu Asp Leu 30	96 .
att gat gct Ile Asp Ala 35	gtg gga atg Val Gly Met	ctt cct aat Leu Pro Asn 40	Glu Lys Val G	aa att gtg aat ln Ile Val Asn 45	144
aat aat aat Asn Asn Asn 50	gga gca cgt Gly Ala Arg	ctt gaa acg Leu Glu Thr 55	tat att att c Tyr Ile Ile P 60	ct ggt aaa cgg ro Gly Lys Arg	192
gga agc ggc Gly Ser Gly 65	gtc ata tgc Val Ile Cys 70	tta aac ggt Leu Asn Gly	gca gcc gca b Ala Ala Ala A 75	gc ctt gtg cag rg Leu Val Gln 80	240
gaa gga gat Glu Gly Asp	aag gtc att Lys Val Ile 85	att att tcc Ile Ile Ser	tac aaa atg a Tyr Lys Met M 90	tg tct gat caa et Ser Asp Gln 95	288
gaa gcg gca Glu Ala Ala	agc cat gag Ser His Glu 100	ccg aaa gtg Pro Lys Val 105	gct gtt ctg a Ala Val Leu A	at gat caa aac sn Asp Gln Asn 110	336
aaa att gaa Lys Ile Glu 115	caa atg ctg Gln Met Leu	ggg aac gaa Gly Asn Glu 120	cca gcc cgt a Pro Ala Arg Ti	ca att ttg hr Ile Leu 25	381
<210> 28 <211> 127 <212> PRT <213> Bacill	us subtilis				
<400> 28 Met Tyr Arg 1	Thr Met Met 5	Ser Gly Lys	Leu His Arg A	la Thr Val Thr 15	
Glu Ala Asn	Leu Asn Tyr 20	Val Gly Ser 25	Ile Thr Ile A	sp Glu Asp Leu 30	
Ile Asp Ala 35	Val Gly Met	Leu Pro Asn 40		ln Ile Val Asn 45	
Asn Asn Asn 50	Gly Ala Arg	Leu Glu Thr 55	Tyr Ile Ile Pr 60	ro Gly Lys Arg	
Gly Ser Gly 65	Val Ile Cys 70	Leu Asn Gly	Ala Ala Ala Ai 75	rg Leu Val Gln 80	

480

Glu Gly Asp Lys Val Ile Ile Ile Ser Tyr Lys Met Met Ser Asp Gln 90 Glu Ala Ala Ser His Glu Pro Lys Val Ala Val Leu Asn Asp Gln Asn 100 . . . 105 110 Lys Ile Glu Gln Met Leu Gly Asn Glu Pro Ala Arg Thr Ile Leu <210> 29 <211> 894 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(894) <400> 29 atg aaa att gga att atc ggc gga ggc tcc gtt ggt ctt tta tgc gcc Met Lys Ile Gly Ile Ile Gly Gly Gly Ser Val Gly Leu Leu Cys Ala 10 tat tat ttg tca ctt tat cac gac gtg act gtt gtg acg agg cgg caa 96 . Tyr Tyr Leu Ser Leu Tyr His Asp Val Thr Val Val Thr Arg Arg Gln 25 gaa cag gct gcg gcc att cag tct gaa gga atc cgg ctt tat aaa ggc 144 Glu Gln Ala Ala Ala Ile Gln Ser Glu Gly Ile Arg Leu Tyr Lys Gly 40 ggg gag gaa ttc agg gct gat tgc agt gcg gac acg agt atc aat tcg 192 Gly Glu Glu Phe Arg Ala Asp Cys Ser Ala Asp Thr Ser Ile Asn Ser 55 gac ttt gac ctg ctt gtc gtg aca gtg aag cag cat cag ctt caa tct 240 Asp Phe Asp Leu Leu Val Val Thr Val Lys Gln His Gln Leu Gln Ser 70 gtt ttt tcg tcg ctt gaa cga atc ggg aag acg aat ata tta ttt ttg 288 Val Phe Ser Ser Leu Glu Arg Ile Gly Lys Thr Asn Ile Leu Phe Leu 85 caa aac ggc atg ggg cat atc cac gac cta aaa gac tgg cac gtt ggc 336 Gln Asn Gly Met Gly His Ile His Asp Leu Lys Asp Trp His Val Gly 105 cat tee att tat gtt gga ate gtt gag eac gga get gta aga aaa teg 384 His Ser Ile Tyr Val Gly Ile Val Glu His Gly Ala Val Arg Lys Ser 120 gat aca gct gtt gat cat aca ggc cta ggt gcg ata aaa tgg agc gcg 432 Asp Thr Ala Val Asp His Thr Gly Leu Gly Ala Ile Lys Trp Ser Ala 135

ttc gac gat gct gaa cca gac cgg ctg aac atc ttg ttt cag cat aac

Phe 145	Asp	Asp	Ala	Glu	Pro 150	Asp	Arg	Leu	Asn	Ile 155	Leu	Phe	Gln	His	Asn 160	
cat His	tcg Ser	gat Asp	ttt Phe	ccg Pro 165	att Ile	tat Tyr	tat Tyr	gag Glu	acg Thr 170	gat Asp	tgg Trp	tac Tyr	cgt Arg	ctg Leu 175	ctg Leu	528
acg Thr	ggc Gly	aag Lys	ctg Leu 180	att Ile	gta Val	aat Asn	gcg Ala	tgt Cys 185	att Ile	aat Asn	cct Pro	tta Leu	act Thr 190	gcg Ala	tta Leu	576
ttg Leu	caa Gln	gtg Val 195	aaa Lys	aat Asn	gga Gly	gaa Glu	ctg Leu 200	ctg Leu	aca Thr	acg Thr	cca Pro	gct Ala 205	tat Tyr	ctg Leu	gct Ala	624
ttt Phe	atg Met 210	aag Lys	ctg Leu	gta Val	ttt Phe	cag Gln 215	gag Glu	gca Ala	tgc Cys	cgc Arg	att Ile 220	tta Leu	aaa Lys	ctt Leu	gaa Glu	672
aat Asn 225	gaa Glu	gaa Glu	aag Lys	gct Ala	tgg Trp 230	gag Glu	cgg Arg	gtt Val	cag Gln	gcc Ala 235	gtt Val	tgt Cys	Gly ggg	caa Gln	acg Thr 240	720
aaa Lys	gag Glu	aat Asn	cgt Arg	tca Ser 245	tca Ser	atg Met	ctg Leu	gtt Val	gac Asp 250	gtc Val	att Ile	gga Gly	ggc Gly	cgg Arg 255	cag Gln	768
acg Thr	gaa Glu	gct Ala	gac Asp 260	gcc Ala	att Ile	atc Ile	gga Gly	tac Tyr 265	tta Leu	ttg Leu	aag Lys	gaa Glu	gca Ala 270	agt Ser	ctt Leu	816
caa Gln	ggt Gly	ctt Leu 275	gat Asp	gcc Ala	gtc Val	cac His	cta Leu 280	gag Glu	ttt Phe	tta Leu	tat Tyr	ggc Gly 285	agc Ser	atc Ile	aaa Lys	864
	ttg Leu 290															,894
<212 <212	0> 30 1> 29 2> PF 3> Ba	98 RT	.us s	subti	ilis											
	0> 30 Lys		Gly	Ile 5	Ile	Gly	Gly	Gly	Ser 10	Val	Gly	Leu	Leu	Cys 15	Ala	
Tyr	Tyr	Leu	Ser 20	Leu	Tyr	His	Asp	Val 25	Thr	Val	Val	Thr	Arg 30		Gln	
Glu	Gln	Ala 35	Ala	Ala	Ile	Gln	Ser 40	Glu	Gly	Ile	Arg	Leu 45	Tyr	Lys	Gly	
Gly	Glu 50	Glu	Phe	Arg	Ala	Asp 55	Cys	Ser	Ala	Asp	Thr 60	Ser	Ile	Asn	Ser	

WO 01/021772 PCT/US00/25993

- 34 -

Asp Phe Asp Leu Leu Val Val Thr Val Lys Gln His Gln Leu Gln Ser

Val Phe Ser Ser Leu Glu Arg Ile Gly Lys Thr Asn Ile Leu Phe Leu 85 90 95

Gln Asn Gly Met Gly His Ile His Asp Leu Lys Asp Trp His Val Gly
100 105 110

His Ser Ile Tyr Val Gly Ile Val Glu His Gly Ala Val Arg Lys Ser 115 120 125

Asp Thr Ala Val Asp His Thr Gly Leu Gly Ala Ile Lys Trp Ser Ala 130 135 140

Phe Asp Asp Ala Glu Pro Asp Arg Leu Asn Ile Leu Phe Gln His Asn 145 150 155 160

His Ser Asp Phe Pro Ile Tyr Tyr Glu Thr Asp Trp Tyr Arg Leu Leu 165 170 175

Thr Gly Lys Leu Ile Val Asn Ala Cys Ile Asn Pro Leu Thr Ala Leu 180 185 190

Leu Gln Val Lys Asn Gly Glu Leu Leu Thr Thr Pro Ala Tyr Leu Ala 195 200 205

Phe Met Lys Leu Val Phe Gln Glu Ala Cys Arg Ile Leu Lys Leu Glu 210 215 220

Asn Glu Glu Lys Ala Trp Glu Arg Val Gln Ala Val Cys Gly Gln Thr 225 230 235 240

Lys Glu Asn Arg Ser Ser Met Leu Val Asp Val Ile Gly Gly Arg Gln
245 250 255

Thr Glu Ala Asp Ala Ile Ile Gly Tyr Leu Leu Lys Glu Ala Ser Leu 260 265 270

Gln Gly Leu Asp Ala Val His Leu Glu Phe Leu Tyr Gly Ser Ile Lys 275 280 285

Ala Leu Glu Arg Asn Thr Asn Lys Val Phe 290 295

<210> 31 <211> 1725

<212> DNA

<213> Bacillus subtilis

<220>

<221> CDS

<222> (1)..(1722)

<400> 31

atg ggg act aat gta cag gtg gat tca gca tct gcc gaa tgt aca cag Met Gly Thr Asn Val Gln Val Asp Ser Ala Ser Ala Glu Cys Thr Gln

1				5					10					15		
acg Thr	atg Met	agc Ser	gga Gly 20	gca Ala	tta Leu	atg Met	ctg Leu	att Ile 25	gaa Glu	tca Ser	tťa Leu	aaa Lys	aaa Lys 30	gag Glu	aaa Lys	96
gta Val	gaa Glu	atg Met 35	atc Ile	ttc Phe	ggt Gly	tat Tyr	ccg Pro 40	ggc Gly	GJ À āāā	gct Ala	gtg Val	ctt Leu 45	ccg Pro	att Ile	tac Tyr	144
gat Asp	aag Lys 50	cta Leu	tac Tyr	aat Asn	tca Ser	ggg Gly 55	ttg Leu	gta Val	cat His	atc Ile	ctt Leu 60	ccc Pro	cgt Arg	cac His	gaa Glu	192
caa Gln 65	gga Gly	gca Ala	att Ile	cat His	gca Ala 70	gcg Ala	gag Glu	gga Gly	tác Tyr	gca Ala 75	agg Arg	gtc Val	tcc Ser	gga Gly	aaa Lys 80	240
ccg Pro	ggt Gly	gtc Val	gtc Val	att Ile 85	gcc Ala	acg Thr	tca Ser	GJ À daa	ccg Pro 90	gga Gly	gcg Ala	aca Thr	aac Asn	ctt Leu 95	gtt Val	288
aca Thr	ggc	ctt Leu	gct Ala 100	gat Asp	gcc Ala	atg Met	att Ile	gat Asp 105	tca Ser	ttg Leu	ccg Pro	tta Leu	gtc Val 110	gtc Val	ttt Phe	336
aca Thr	Gly ggg	cag Gln 115	gta Val	gca Ala	acc Thr	tct Ser	gta Val 120	atc Ile	Gly ggg	agc Ser	gat Asp	gca Ala 125	ttt Phe	cag Gln	gaa Glu	384
gca Ala	gac Asp 130	att Ile	tta Leu	Gly ggg	att Ile	acg Thr 135	atg Met	cca Pro	gta Val	aca Thr	aaa Lys 140	cac His	agc Ser	tac Tyr	cag Gln	432
gtt Val 145	cgc Arg	cag Gln	ccg Pro	gaa Glu	gat Asp 150	ctg Leu	ccg Pro	cgc Arg	atc Ile	att Ile 155	aaa Lys	gaa Glu	gcg Ala	ttc Phe	cat His 160	480
att Ile	gca Ala	aca Thr	act Thr	gga Gly 165	aga Arg	ccc Pro	gga Gly	cct Pro	gta Val 170	ttg Leu	att Ile	gat Asp	att Ile	ccg Pro 175	aaa Lys	528
gat Asp	gta Val	gca Ala	aca Thr 180	att Ile	gaa Glu	gga Gly	gaa Glu	ttc Phe 185	agc Ser	tac Tyr	gat Asp	cat His	gag Glu 190	atg Met	aat Asn	576
ctc Leu	ccg Pro	gga Gly 195	tac Tyr	cag Gln	ccg Pro	aca Thr	aca Thr 200	gag Glu	ccg Pro	aat Asn	tat Tyr	ttg Leu 205	cag Gln	atc Ile	cgc Arg	624
aag Lys	ctt Leu 210	gtg Val	gaa Glu	gcc Ala	gtg Val	agc Ser 215	agt Ser	gcg Ala	aaa Lys	aaa Lys	ccg Pro 220	gtg Val	atc Ile	ctg Leu	gcg Ala	672
ggt Gly 225	gcg Ala	ggc Gly	gta Val	ctg Leu	cac His 230	gga Gly	aaa Lys	gcg Ala	tca Ser	gaa Glu 235	gaa Glu	tta Leu	aaa Lys	aat Asn	tat Tyr 240	720

gct Ala	gaa Glu	cag Gln	cag Gln	caa Gln 245	atc' Ile	cct Pro	gtg Val	Ala	cac His 250	acc Thr	ctt Leu	ttg Leu	G1 y G3 g	ctc Leu 255	gga Gly	768
ggc Gly	ttc Phe	ccg Pro	gct Ala 260	gac Asp	cat His	ecg Pro	ctt Leu	ttc Phe 265	cta Leu	G1y ggg	atg Met	gcg Ala	gga Gly 270	atg Met	cac His	816
ggt Gly	act Thr	tat Tyr 275	aca Thr	gcc Ala	aat Asn	atg Met	gcc Ala 280	ctt Leu	cat His	gaa Glu	tgt Cys	gat Asp 285	cta Leu	tta Leu	atc Ile	864
agt Ser	atc Ile 290	ggc	gcc Ala	cgt Arg	ttt Phe	gat Asp 295	gac Asp	cgt Arg	gtc Val	aca Th <i>r</i>	gga Gly 300	aac Asn	ctg Leu	aaa Lys	cac His	912
ttt Phe 305	Ala	aga Arg	aac Asn	gca Ala	aag Lys 310	ata Ile	gcc Ala	cac His	atc Ile	gat Asp 315	att Ile	gat, Asp	Pro	gct Ala	gaa Glu 320	960
atc Ile	gga Gly	aaa Lys	atc	atg Met 325	Lys	aca Thr	cag Gln	att Ile	cct Pro 330	gta Val	gtc Val	gga Gly	gac Asp	agc Ser 335	aaa Lys	1008
att Ile	gtc Val	cto Lev	caç Glr 340	gag Glu	ctg Leu	atc Ile	aaa Lys	caa Gln 345	gac Asp	ggc Gly	aaa Lys	⊶caa Gln	ago Ser 350	Asp	tca Ser	1056
ago Ser	gaa Glu	tgg Trp 359	Lys	a aaa s Lys	cag Gln	ctc Leu	gca Ala 360	Glu	tgg Trp	aaa Lys	gaa Glu	gag Glu 365	Tyr	ccg Pro	ctc Leu	1104
tgg Trg	tat Tyi	· Vai	a gat L As _l	c aat o Asr	gaa Glu	gaa Glu 375	Glu	ggt Gly	ttt Phe	aaa Lys	ect Pro	o Glr	aaa Lys	ttg Lev	att Ile	1152
ga: Gl: 38:	ı Tyı	t at	t car e Hi	t caa s Gl	a ttt n Phe 390	e Thr	aaa Lys	a gga s Gly	gaç Glu	g gcc Ala 395	a II	t gto e Val	e gea L Ala	a acg a Thi	gat Asp 400	1200
gt. Va	a gg	c ca y Gl	g ca n Hi	t cas s Gli 40	n Mei	g tgg t Tr	tca Ser	a gcg	g caa Glr 410	n Phe	t ta e Ty	t cco	g tte o Phe	c caa e Gl: 41	a aaa n Lys 5	1248
gc Al	a ga a As	t aa p Ly	a tg s Tr 42	p Va	c ac	g tca r Sea	a ggo r Gl	c gga y Gly 42	y Le	t gg u Gl	a ac y Th	g ate	g gg t Gl 43	λ _δ υ	c ggt e Gly	1296
ct Le	t cc u Pr	g gc o Al 43	a Al	g at a Il	c gg e Gl	c gc y Al	a ca a Gl	n Le	g gce u Ala	c ga a Gl	a aa u Ly	a ga rs As 44	Ь ЧТ	t ac a Th	t gtt r Val	1344
gt Va	c go l Al 45	a Va	t gt il Va	c gg	a ga y As	c gg p Gl 45	y Gl	a tt y Ph	c ca e Gl	a at n Me	g ac t Th	ır re	t ca u Gl	a ga n Gl	a ctc u Leu	1392
ga As	it gt sp Va	t at	t co	ge ga eg Gl	a tt .u Le	a aa u As	t ct n Le	t cc u Pr	g gt o Va	c aa l Ly	g gt 's Va	a gt	g at	t tt e Le	a aat u Asn	1440

- 37 -

465	470	476		
		475	480	
aac got tgt ctc Asn Ala Cys Leu	gga atg gtc ag Gly Met Val Ar 485	ga cag tgg cag gaa gg Gln Trp Gln Glu 490	a att ttc tat gaa : n Ile Phe Tyr Glu 495	1488
gaa cgt tat tca Glu Arg Tyr Ser 500	gaa tct aaa tt Glu Ser Lys Ph	c gct tct cag cct le Ala Ser Gln Pro 505	gac ttc gtc aaa : Asp Phe Val Lys 510	1536
ttg tcc gaa gca Leu Ser Glu Ala 515	tac ggc att aa Tyr Gly Ile Ly 52	a ggc atc aga att s Gly Ile Arg Ile 0	tca tca gaa gcg Ser Ser Glu Ala 525	1584
gaa gca aag gaa Glu Ala Lys Glu 530	aag ctg gaa ga Lys Leu Glu Gl 535	g gca tta aca tca u Ala Leu Thr Ser 540	Arg Glu Pro Val	1632
gtc att gac gtg Val Ile Asp Val 545	cgg gtt gcc ag Arg Val Ala Se 550	c gaa gaa aaa gta r Glu Glu Lys Val 555	ttc ccg atg gtg 1 Phe Pro Met Val , 560	1680
gct ccg ggg aaa Ala Pro Gly Lys	ggg ctg cat ga Gly Leu His Gl 565	a atg gtg ggg gtg u Met Val Gly Val 570	aaa cct tga I	L725
<210> 32 <211> 574 <212> PRT <213> Bacillus s				
Met Gly Thr Asn	Val Gln Val Ası 5	p Ser Ala Ser Ala		
T	3	10	Glu Cys Thr Gln 15	
			15	
Thr Met Ser Gly 20	Ala Leu Met Le	10 u Ile Glu Ser Leu 25 o Gly Gly Ala Val	Lys Lys Glu Lys	
Thr Met Ser Gly 20 Val Glu Met Ile 35	Ala Leu Met Le Phe Gly Tyr Pro 40	10 u Ile Glu Ser Leu 25 o Gly Gly Ala Val	Lys Lys Glu Lys 30 Leu Pro Ile Tyr 45 Pro Arg His Glu	
Thr Met Ser Gly 20 Val Glu Met Ile 35 Asp Lys Leu Tyr 50	Ala Leu Met Le Phe Gly Tyr Pro 40 Asn Ser Gly Leu 55	10 u Ile Glu Ser Leu 25 c Gly Gly Ala Val 0 u Val His Ile Leu	Lys Lys Glu Lys 30 Leu Pro Ile Tyr 45 Pro Arg His Glu	
Thr Met Ser Gly 20 Val Glu Met Ile 35 Asp Lys Leu Tyr 50 Gln Gly Ala Ile 65	Ala Leu Met Leo Phe Gly Tyr Pro 46 Asn Ser Gly Leo 55 His Ala Ala Glo 70	10 u Ile Glu Ser Leu 25 c Gly Gly Ala Val u Val His Ile Leu 60 u Gly Tyr Ala Arg	Lys Lys Glu Lys 30 Leu Pro Ile Tyr 45 Pro Arg His Glu Val Ser Gly Lys 80	
Thr Met Ser Gly 20 Val Glu Met Ile 35 Asp Lys Leu Tyr 50 Gln Gly Ala Ile 65 Pro Gly Val Val	Ala Leu Met Leo Phe Gly Tyr Pro 40 Asn Ser Gly Leo 55 His Ala Ala Glo 70 Ile Ala Thr Ser 85	10 u Ile Glu Ser Leu 25 c Gly Gly Ala Val u Val His Ile Leu 60 u Gly Tyr Ala Arg 75 r Gly Pro Gly Ala	Lys Lys Glu Lys 30 Leu Pro Ile Tyr 45 Pro Arg His Glu Val Ser Gly Lys 80 Thr Asn Leu Val 95	
Thr Met Ser Gly 20 Val Glu Met Ile 35 Asp Lys Leu Tyr 50 Gln Gly Ala Ile 65 Pro Gly Val Val 1 Thr Gly Leu Ala 1	Ala Leu Met Leu Phe Gly Tyr Pro 46 Asn Ser Gly Leu 55 His Ala Ala Glu 70 Ile Ala Thr Sen 85 Asp Ala Met Ile	10 u Ile Glu Ser Leu 25 c Gly Gly Ala Val u Val His Ile Leu 60 u Gly Tyr Ala Arg 75 r Gly Pro Gly Ala 90 e Asp Ser Leu Pro 105	Lys Lys Glu Lys 30 Leu Pro Ile Tyr 45 Pro Arg His Glu Val Ser Gly Lys 80 Thr Asn Leu Val 95 Leu Val Val Phe 110	

	130					135					140				
Val 145	Arg	Gln	Pro	Glu	Asp 150	Leu	Pro	Arg		Ile 155	Lys	Glu	Ala	Phe	His 160
Ile	Ala	Thr	Thr	Gly 165	Arg	Pro	Gly	Pro	Val 170	Leu	Ile	Asp		Pro 175	Lys
Asp	Val	Ala	Thr 180	Ile	Glu	Gly	Glu	Phe 185	Ser	Tyr	Asp	His	Glu 190	Met	Asn
Leu	Pro	Gly 195	Tyr	Gln	Pro	Thr	Thr 200	Glu	Pro	Asn	Tyr	Leu 205	Gln	Ile	Arg
Lys	Leu 210	Val	Glu	Ala	Val	Ser 215	Ser	Ala	Lys	Lys	Pro 220	Val	Ile	Leu	Ala
Gly 225	Ala	Gly	Val	Leu	His 230	Gly	Lys	Ala	Ser	Glu 235	Glu	Leu	Lys	Asn	Tyr 240
,Ala	Glu	Gln	Gln	Gln 245	Ile	Pro	Val	Ala	His 250	Thr	Leu	Leu	-Gly	Leu 255	Gly
Gly	Phe	Pro	Ala 260		His	Pro	Leu	Phe 265	Leu	Gly	Met	Ala	Gly 270	Met	His
Gly	Thr	Tyr 275		Ala	Asn	Met	Ala 280	Leu	His	Glu	Cys	Asp 285		Leu	Ile
Ser	Ile 290		Ala	Arg	Phe	Asp 295		Arg	Val	Thr	Gly 300		Leu	Lys	His
Phe 305		Arg	Asn	Ala	Lys 310	Ile	Ala	His	Ile	Asp 315	Ile	Asp	Pro	Ala	Glu 320
Ile	Gly	Lys	Ile	Met 325		Thr	Gln	Ile	Pro 330		Val	∙Gly	Asp	Ser 335	Lys
Ile	Val	Leu	Gln 340		Leu	Ile	Lys	Gln 345		Gly	Lys	Gln	Ser 350		Ser
Ser	Glu	355	_	Lys	-Gln	Leu	Ala 360		Trp	Lys	~Glu	Glu 365		Pro	Leu
Trp	370		. Asp	Asr	Glu	Glu 375		Gly	·Phe	. Lys	2rc 380		Lys	Leu	Ile
Gl 385	_	: Ile	His	s Glr	390		Lys	-Gly	∙ Glu	395		e Val	. Ala	Thi	400
Val	L Gly	/ Glr	n His	405	_	Trţ	Ser	Ala	Gln 410		туг	Pro	Phe	Glr 415	Lys
Ala	a Ası	p Lys	420		l Thr	: Sea	: Gly	Gly 425		ı Gly	Tha	Met	Gl ₃		e Gly
, Le	ı Pro	Ala 435		a Ile	e Gly	/ Ala	440		Ala	Glu	Lys	445		Th:	val

Val	Ala 450	Val	Val	Gly	Asp	Gly 455	Gly	Phe	Gln	Met	Thr 460	Leu	Gln	Glu	Leu	
Asp 465	Val	Ile	Arg	Glu	Leu 470	Asn	Leu	Pro	Val	Lys 475	Val	Val	Ile	Leu	Asn 480	
Asn	Ala	Cys	Leu	Gly 485	Met	Val	Arg	Gln	Trp 490	Gln	Glu	Ile _.	Phe	Tyr 495	Glu	
Glu	Arg	Tyr	Ser 500	Glu	Ser	Lys	Phe	Ala 505	Ser	Gln	Pro	Asp	Phe 510	Val	Lys	
Leu	Ser	Glu 515	Ala	Tyr	Gly	Ile	Lys 520	Gly	Ile	Arg	Ile	Ser 525	Ser	Glu	Ala	
Glu	Ala 530	Lys	Glu	Lys	Leu	Glu 535	Glu	Ala	Leu	Thr	Ser 540	Arg	Glu	Pro	Val	
Val 545	Ile	Asp	Val	Arg	Val 550	Ala	Ser	Glu	Glu	Lys 555	Val	Phe	Pro	Met	Val 560	
Ala	Pro	Gly	Lys	Gly 565	Leu	His	Glu	Met	Val 570	Gly	Val	Lys	Pro			
<211 <212 <213 <220	0> 33 1> 52 2> DN 3> Ba 0> CI	25 NA Acil:	lus s	subti	ilis											
	2> (1)> 33		(522))												
ttg	aaa Lys	aga	att Ile	atc Ile 5	aca Thr	ttg Leu	act Thr	gtg Val	gtg Val 10	aac Asn	cgc Arg	tcc Ser	GJ À GG À	gtg Val 15	tta Leu	48
aac Asn	cgg Arg	atc Ile	acc Thr 20	ggt Gly	cta Leu	ttc Phe	aca Thr	aaa Lys 25	agg Arg	cat His	tac Tyr	aac Asn	att Ile 30	gaa Glu	agc Ser	96
att Ile	aca Thr	gtt Val 35	gga Gly	cac His	aca Thr	gaa Glu	aca Thr 40	gcc Ala	ggc Gly	gtt Val	tcc Ser	aga Arg 45	atc Ile	acc Thr	ttc Phe	144
gtc Val	gtt Val 50	cat His	gtt Val	gaa Glu	ggt Gly	gaa Glu 55	aat Asn	gat Asp	gtt Val	gaa Glu	cag Gln 60	tta Leu	acg Thr	aaa Lys	cag Gln	192
ctc Leu 65	aac Asn	aaa Lys	cag Gln	att Ile	gat Asp 70	gtg Val	ctg Leu	aaa Lys	gtc Val	aca Thr 75	gac Asp	atc Ile	aca Thr	aat Asn	caa Gln 80	240
tcg	att	gtc	cag	agg	gag	ctg	gcc	tta	atc	aag	gtt	gtc	tcc	gca	cct	288

- 40 -

	85		90	' 9:	5
Ser Thr Arg '	aca gag att Thr Glu Ile 100	aat gga at Asn Gly Il 10	e Ile Glu Pr	g ttt aga gc o Phe Arg Al 110	c tct 336 a Ser
gtc gtt gat Val Val Asp 115	gtc agc aga Val Ser Arg	gac agc at Asp Ser Il 120	c gtt gtt ca e Val Val Gl	g gtg aca gg n Val Thr Gl 125	t gaa 384 y Glu
tct aac aaa Ser Asn Lys 130	att gaa gcg Ile Glu Ala	ctt att ga Leu Ile Gl 135	ag tta tta aa lu Leu Leu Ly 14	aa cct tat gg vs Pro Tyr Gl 10	c att 432 y Ile
aaa gaa atc . Lys Glu Ile 145	gcg aga aca Ala Arg Thr 150	ggt aca ac Gly Thr Th	eg get ttt ge nr Ala Phe Al 155	cg agg gga ac La Arg Gly Th	c agc 480 r Ser 160
aaa agg cgt Lys Arg Arg	cat cca ata His Pro Ile 165	aaa caa ta Lys Gln Ty	at cta ttg ta yr Leu Leu Ty 170	at aaa aca ta yr Lys Thr	a 525
<210> 34 <211> 174 <212> PRT <213> Bacill	lus subtilis			v. 1	. •
<400> 34 Met Lys Arg 1	Ile Ile Thr	Leu Thr V	al Val Asn A	rg Ser Gly Va	al Leu 15
Asn Arg Ile	Thr Gly Leu 20		ys Arg His T 25	yr Asn Ile Gl 30	lu Ser
Ile Thr Val	Gly His Thr	Glu Thr A	la Gly Val S	er Arg Ile Ti 45	nr Phe
Val Val His 50	Val Glu Gly	Glu Asn A 55	sp Val Glu G	ln Leu Thr L 60	ys Gln
Leu Asn Lys 65	Gln Ile Asp 70		ys Val Thr A 75	sp Ile Thr A	sn Gln 80
Ser Ile Val	Gln Arg Glu 85	Leu Ala L	eu Ile Lys V 90	al Val Ser A	la Pro 95
Ser Thr Arg	Thr Glu Ile		le lle Glu E 105	Pro Phe Arg A 110	la Ser
Val Val Asp 115		Asp Ser I 120	Ile Val Val €	Sin Val Thr G 125	ly Glu
Ser Asn Lys 130	Ile Glu Ala	Leu Ile G		Lys Pro Tyr G L40	ly Ile
Lys Glu Ile 145	e Ala Arg Thi 150		Thr Ala Phe A	Ala Arg Gly T	hr Ser 160

Lys Arg Arg His Pro Ile Lys Gln Tyr Leu Leu Tyr Lys Thr 165 170

<210> 35 <211> 1029 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(1026) <400> 35 atg gta aaa gta tat tat aac ggt gat atc aaa gag aac gta ttg gct Met Val Lys Val Tyr Tyr Asn Gly Asp Ile Lys Glu Asn Val Leu Ala gga aaa aca gta gcg gtt atc ggg tac ggt tcg caa ggc cac gca cat Gly Lys Thr Val Ala Val Ile Gly Tyr Gly Ser Gln Gly His Ala His gee etg aac ett aaa gaa age gga gta gae gtg ate gte ggt gtt aga Ala Leu Asn Leu Lys Glu Ser Gly Val Asp Val Ile Val Gly Val Arg caa gga aaa tet tte act caa gee caa gaa gae gga cat aaa gta ttt 192 Gln Gly Lys Ser Phe Thr Gln Ala Gln Glu Asp Gly His Lys Val Phe 50 55 tca gta aaa gaa gcg gca gcc caa gcc gaa atc atc atg gtt ctg ctt Ser Val Lys Glu Ala Ala Ala Gln Ala Glu Ile Ile Met Val Leu Leu 65 ccg gat gag cag caa aaa gta tac gaa gct gaa atc aaa gat gaa 288 Pro Asp Glu Gln Gln Lys Val Tyr Glu Ala Glu Ile Lys Asp Glu ttg aca gca gga aaa tca tta gta ttc gct cat gga ttt aac gtg cat 336 Leu Thr Ala Gly Lys Ser Leu Val Phe Ala His Gly Phe Asn Val His 100 ttc cat caa att gtt cct ccg gcg gat gta gta ttc tta gtg gcc Phe His Gln Ile Val Pro Pro Ala Asp Val Asp Val Phe Leu Val Ala 115 cet aaa ggc eeg gga cac ttg gta aga aga aca tat gag caa gga get 432 Pro Lys Gly Pro Gly His Leu Val Arg Arg Thr. Tyr Glu Gln Gly Ala 130 135 ggc gta cct gca ttg ttc gca atc tat caa gat gtg act gga gaa gca 480 Gly Val Pro Ala Leu Phe Ala Ile Tyr Gln Asp Val Thr Gly Glu Ala 145 150 aga gac aaa gcc ctc gct tat gct aaa gga atc ggc ggc gca aga gcg 528 Arg Asp Lys Ala Leu Ala Tyr Ala Lys Gly Ile Gly Gly Ala Arg Ala 170

ggc Gly	gta Val	tta Leu	gaa Glu 180	acg Thr	aca Thr	ttt Phe	aaa Lys	gaa Glu 185	gaa Glu	aca Thr	gaa Glu	aca Thr	gat Asp 190	ttg Leu	ttc Phe	576
ggt Gly	gag Glu	caa Gln 195	gca Ala	gtt Val	ctt Leu	tgc Cys	ggc Gly 200	gga Gly	tta Leu	agc Ser	gcg Ala	ctt Leu 205	gtc Val	aaa Łys	gcc Ala	624
gga Gly	ttt Phe 210	gaa Glu	acc Thr	tta Leu	act Thr	gaa Glu 215	gca Ala	ggt Gly	tat Tyr	cag Gln	cct Pro 220	gaa Glu	ctt Leu	gca Ala	tac Tyr	672,
ttc Phe 225	gag Glu	tgt Cys	ctt Leu	cat His	gag Glu 230	ctg Leu	aaa Lys	tta Leu	atc Ile	gta Val 235	Asp	ctt Leu	atg Met	tac Tyr	gaa Glu 240	720
gaa Glu	gga Gly	ctt Leu	gca Ala	gga Gly 245	atg Met	aga Arg	tat Tyr	tca Ser	atc Ile 250	tct Ser	gac Asp	aca Thr	gca Ala	cag Gln 255	tgg Trp	768
gga Gly	gat Asp	ttc Phe	gta Val 260	Ser	Gly	cct Pro	cgc Arg	gtt Val 265	Val	gac Asp	gcc Ala	aaa Lys	gta Val 270	. гля	gaa Glu	816
tct Ser	ato Met	aaa Lys 275	s Glu	gta Val	tta Leu	aaa Lys	gat Asp 280	Ile	caa Gln	aac Asr	ggt Gly	Thr 285	Pne	c gca e Ala	aaa Lys	864
gaç Glı	tgi Tri 290	ıl.	c gto e Val	gaa Glu	aac Asn	caa Gln 295	. Val	a aac L Ast	cgt Arg	cct Pro	cgt Arq 300	g Phe	e Ası	e get n Ala	atc Ile	912
aat Asi 30!	n Al	a age	c gaq r Gl	g aad u Asi	gaa Glu 310	His	caa Gli	a ato	gaa e Glu	a gta 1 Va. 31	l Va.	g gga	a aga y Ar	a aaq g Ly:	g ctt s Leu 320	9,60
cg Ar	t ga g Gl	a at u Me	g ate t Me	g cce t Pre 32	o Phe	t gto	g aaa l Ly	a -caa s Gli	a gg n Gl; 33	у Гу	g aa s Ly	g aa s Ly	g ga s Gl	a gc u Al 33	g gtg a Val 5	1008
gt Va	c tc 1 Se	c gt r Va	t gc l Al 34	a Gl	a aai n Asi	t ta: n	ā									1029
<2 <2	10> 11> 12> 13>	342 PRT	illus	s sub	tili	S										

<400> 36 Met Val Lys Val Tyr Tyr Asn Gly Asp Ile Lys Glu Asn Val Leu Ala 1 5 10 15

Gly Lys Thr Val Ala Val Ile Gly Tyr Gly Ser Gln Gly His Ala His 20 25 30

Ala Leu Asn Leu Lys Glu Ser Gly Val Asp Val Ile Val Gly Val Arg · 35 40 45

Gln Gly Lys Ser Phe Thr Gln Ala Gln Glu Asp Gly His Lys Val Phe 50 55 60

Ser Val Lys Glu Ala Ala Ala Gln Ala Glu Ile Ile Met Val Leu Leu 65 70 75 80

Pro Asp Glu Gln Gln Lys Val Tyr Glu Ala Glu Ile Lys Asp Glu 85 90 95

Leu Thr Ala Gly Lys Ser Leu Val Phe Ala His Gly Phe Asn Val His
100 105 110

Phe His Gln Ile Val Pro Pro Ala Asp Val Asp Val Phe Leu Val Ala 115 120 . 125

Pro Lys Gly Pro Gly His Leu Val Arg Arg Thr Tyr Glu Gln Gly Ala 130 135 140

Gly Val Pro Ala Leu Phe Ala Ile Tyr Gln Asp Val Thr Gly Glu Ala 145 150 155 160

Arg Asp Lys Ala Leu Ala Tyr Ala Lys Gly Ile Gly Gly Ala Arg Ala 165 170 175

Gly Val Leu Glu Thr Thr Phe Lys Glu Glu Thr Glu Thr Asp Leu Phe 180 185 190

Gly Glu Gln Ala Val Leu Cys Gly Gly Leu Ser Ala Leu Val Lys Ala 195 200 205

Gly Phe Glu Thr Leu Thr Glu Ala Gly Tyr Gln Pro Glu Leu Ala Tyr 210 215 220

Phe Glu Cys Leu His Glu Leu Lys Leu Ile Val Asp Leu Met Tyr Glu 225 230 235 240

Glu Gly Leu Ala Gly Met Arg Tyr Ser Ile Ser Asp Thr Ala Gln Trp 245 250 255

Gly Asp Phe Val Ser Gly Pro Arg Val Val Asp Ala Lys Val Lys Glu 260 265 270

Ser Met Lys Glu Val Leu Lys Asp Ile Gln Asn Gly Thr Phe Ala Lys 275 280 285

Glu Trp Ile Val Glu Asn Gln Val Asn Arg Pro Arg Phe Asn Ala Ile 290 295 300

Asn Ala Ser Glu Asn Glu His Gln Ile Glu Val Val Gly Arg Lys Leu 305 310 315 320

Arg Glu Met Met Pro Phe Val Lys Gln Gly Lys Lys Glu Ala Val 325 330 335

Val Ser Val Ala Gln Asn

340

<210 <211 <212 <213	> 16' > DN	A	us s	ubti:	lis								ı				•
<220: <221: <222:	> CD		1674)													
<400 atg Met 2	σca	σaa	tta Leu	cgc Arg 5	agt Ser	aat Asn	atg Met	atc Ile	aca Thr 10	caa Gln	gga Gly	atc Ile	gat Asp	aga Arg 15	gct Ala	48	
ccg Pro	cac His	.cgc Arg	agt Ser 20	ttg Leu	ctt Leu	cgt Arg	gca Ala	gca Ala 25	GJ A GGG	gta Val	aaa Lys	gaa Glu	gag Glu 30	gat Asp	ttc Phe	96	
ggc	aag Lys	ccg Pro 35	ttt Phe	att Ile	gcg Ala	gtg Val	tgt Cys 40	aat Asn	tca Ser	tac Tyr	att Ile	gat Asp 45	atc Ile	gtt Val	ccc Pro	144	
ggt Gly	cat His 50	gtt Val	cac His	ttg Leu	cag Gln	gag Glu 55	ttt Phe	GJ À .āāā	aaa Lys	atc Ile	gta Val 60	aaa Lys	gaa Glu	gca Ala	atc Ile	192	•
aga Arg 65	gaa Glu	gca Ala	Gly	ggc Gly	gtt Val 70	ccg Pro	ttt Phe	gaa Glu	ttt Phe	aat Asn 75	acc Thr	att Ile	.GJA aaa	gta Val	gat Asp 80	240	
gat Asp	ggc Gly	at.c Ile	gca Ala	atg Met 85	Gly	cat His	atc Ile	ggt Gly	atg Met 90	aga Arg	tat Tyr	tcg Ser	ctg Leu	cca Pro 95	Ser	288	
cgt Arg	gaa Glu	att Ile	atc Ile 100	Ala	gac Asp	tct Ser	gtg Val	gaa Glu 105	Thr	gtt Val	gta Val	tcc Ser	gca Ala 110	cac His	tgg Trp	33.6	
ttt Phe	gac Asp	gga Gly 115	Met	gtc Val	tgt Cys	att Ile	ccg Pro 120	Asn	tgc Cys	gac Asp	aaa Lys	atc Ile 125	Thr	-ccg Pro	gga Gly	384	
atg Met	ctt Leu 130	Met	gcg Ala	gca Ala	atg Met	cgc Arg 135	Ile	aac	att	ccg Pro	acg Thr 140	Ile	ttt Phe	gto Val	agc Ser	432	
ggc Gly 145	Gly	ccg Pro	g atg Met	gcg Ala	gca Ala 150	Gly	aga Arg	aca Thr	agt Ser	tac Tyr 155	: Gl	r cga / Arg	aaa Lys	ato Ile	tcc Ser 160	480	
ctt Leu	tcc Ser	tca Ser	gta Val	tto Phe 165	Glu	ggg Gly	gta Val	ggc Gly	geo Ala 170	Туг	caa Glr	a gca n Ala	ggg Gly	aaa Lys 175	a atc s Ile	528	
aac	gaa	aaq	gaç	ctt	caa	a gaa	cta	gag	g caç	tto	gga	a tgo	cca	ace	g tgc	576	

_																
Asn	Glu	Asn	Glu 180	Leu	Gln	Glu	Leu	Glu 185	Gln	Phe	Gly	Cys	Pro 190		Cys	•
GJÀ	tct Ser	tgc Cys 195	tca Ser	ggc Gly	atg Met	ttt Phe	acg Thr 200	gcg Ala	aac Asn	tca Ser	atg Met	aac Asn 205	tgt Cys	ctg Leu	tca Ser	624
gaa Glu	gca Ala 210	ctt Leu	ggt Gly	ctt Leu	gct Ala	ttg Leu 215	ccg Pro	ggt Gly	aat Asn	gga Gly	acc Thr 220	att Ile	ctg Leu	gca Ala	aca Thr	672
tct Ser 225	ccg Pro	gaa Glu	cgc Arg	aaa Lys	gag Glu 230	ttt Phe	gtg Val	aga Arg	aaa Lys	tcg Ser 235	gct Ala	gcg Ala	caa Gln	tta Leu	atg Met 240	720
gaa Glu	acg Thr	att Ile	cgc Arg	aaa Lys 245	gat Asp	atc Ile	aaa Lys	ccg Pro	cgt Arg 250	gat Asp	att Ile	gtt Val	aca Thr	gta Val 255	aaa Lys	768
gcg Ala	att Ile	gat Asp	aac Asn 260	gcg Ala	ttt Phe	gca Ala	ctc Leu	gat Asp 265	atg Met	gcg Ala	ctc Leu	gga Gly	ggt Gly 270	tct Ser	aca Thr	816
aat Asn	acc Thr	gtt Val 275	ctt Leu	cat His	acc Thr	ctt Leu	gcc Ala 280	ctt Leu	gca Ala	aac Asn	gaa Glu	gcc Ala 285	ggc Gly	gtt Val	gaa Glu	864
tac Tyr	tct Ser 290	tta Leu	gaa Glu	cgc Arg	att ['] Ile	aac Asn 295	gaa Glu	gtc Val	gct Ala	gag Glu	cgc Arg 300	gtg Val	ccg Pro	cac His	ttg Leu	912
gct Ala 305	aag Lys	ctg Leu	gcg Ala	cct Pro	gca Ala 310	tcg Ser	gat Asp	gtg Val	ttt Phe	att Ile 315	gaa Glu	gat Asp	ctt Leu	cac His	gaa Glu 320	960
gcg Ala	Gly Ggc	ggc Gly	gtt Val	tca Ser 325	gcg Ala	gct Ala	ctg Leu	aat Asn	gag Glu 330	ctt Leu	tcg Ser	aag Lys	aaa Lys	gaa Glu 335	gga Gly	1008
gcg Ala	ctt Leu	cat His	tta Leu 340	gat Asp	gcg Ala	ctg Leu	act Thr	gtt Val 345	aca Thr	gga Gly	aaa Lys	act Thr	ctt Leu 350	gga Gly	gaa Glu	1056
acc Thr	att Ile	gcc Ala 355	gga Gly	cat His	gaa Glu	gta Val	aag Lys 360	gat Asp	tat Tyr	gac Asp	gtc Val	att Ile 365	cac His	ccg Pro	ctg Leu	1104
gat Asp	caa Gln 370	cca Pro	ttc Phe	act Thr	gaa Glu	aag Lys 375	gga Gly	ggc Gly	ctt Leu	gct Ala	gtt Val 380	tta Leu	ttc Phe	ggt Gly	aat Asn	1152
cta Leu 385	gct Ala	ccg Pro	gac Asp	ggc Gly	gct Ala 390	atc Ile	att Ile	aaa Lys	aca Thr	ggc Gly 395	ggc Gly	gta Val	cag Gln	aat Asn	999 Gly 400	1200
att Ile	aca Thr	aga Arg	cac His	gaa Glu 405	G1y ggg	ccg Pro	gct Ala	gtc Val	gta Val 410	ttc Phe	gat Asp	tct Ser	cag Gln	gac Asp 415	gag Glu	1248

										•						
gcg Ala	ctt Leu	gac Asp	ggc Gly 420	att Ile	atc Ile	äac Asn	cga Arg	aaa Lys 425	gta Val	aaa Lys	gaa Glu	ggc Gly	gac Asp 430	gtt Val	gtc Val	1296
atc Ile	atc Ile	aga Arg 435	tac Tyr	gaa Glu	Gly ggg	cca Pro	aaa Lys 440	.eJÀ ààc	gga Gly	cct Pro	ggc Gly	atg Met 445	ccg Pro	gaa Glu	atg Met	1344
ctg Leu	gcg Ala 450	cca Pro	aca Thr	tcc Ser	caa Gln	atc Ile 455	gtt Val	gga Gly	atg Met	gga Gly	ctc Leu 460	GJ A GGG	cca Pro	aaa Lys	gtg Val	1392
gca Ala 465	ttg Leu	att Ile	acg Thr	gac Asp	gga Gly 470	cgt Arg	ttt Phe	tcc Ser	gga Gly	gcc Ala 475	tcc Ser	cgt Arg	ggc Gly	ctc Leu	tca Ser 480	1440
atc Ile	ggc Gly	cac His	gta Val	tca Ser 485	cct Pro	gag Glu	gcc Ala	gct Ala	gag Glu 490	ggc Gly	ggg Gly	ecg Pro	ctt Leu	gcc Ala 495	P.he	1488
gtt Val	gaa Glu	aac Asn	gga Gly 500	Asp	cat His	att Ile	atc Ile	gtt Val 505	gat Asp	att Ile	gaa Glu	aaa Lys	cgc Arg 510	atc Ile	ttg Leu	1536
gat Asp	gta Val	caa Gln 515	- Val	cca Pro	gaa Glu	gaa Glu	gag Glu 520	Trp	gaa Glu	aaa Lys	cga Arg	aaa Lys 525	Ala	aac Asn	tgg Trp	1584
aaa Lys	ggt Gly 530	Phe	gaa Glu	ccg	aaa Lys	gtg Val	. Lys	acc Thr	ggc	tac Tyr	ctg Leu 540	Ala	cgt Arg	tat Tyr	tct Ser	1632
aaa Lys 545	Leu	gtg Val	aca Thr	agt Ser	gcc Ala 550	Asr	acc Thr	ggc Gly	ggt Gly	att Ile 555	Met	aaa Lys	atc : Ile	: :		1674
<21 <21	10> 3 11> 5 12> F	558 PRT	llus	subt	:ilis	3										
Met	00> 3 : Ala	38 a Glu	ı Leı	_	g Sei	c Ası	n Met	: Ile	Th:		ı Gly	y Ile	e Asp	Are	g Ala 5	
Pro	o His	s Ar	g Ser 20	_	ı Let	ı Ar	g Ala	a Ala 25		/ Val	L £y:	s Glu	3 G14		p Phe	
Gl	y Lys	s Pro		e Ile	e Ala	a Va	1 Cy:		n Sei	туз	r Ile	e Ası		e Va	l Pro	
G1;	y Hi:		l Hi	s Le	u Gl	n Gl [.] 5		e Gly	у Ьу:	s Ile	e Va 6		s Gl	u Al	a Ile	
Ar 6	_	u Al	a Gl	y Gl	y Va 7		o Ph	e Gl	ı Pho	e Ası 7		r Il	e Gl	y Va	1 Asp 60	

Asp	Gly	Ile	Ala	Met 85	Gly	His	Ile	Gly	Met 90	Arg	Tyr	Ser	Leu	Pro 95	Ser
Arg	Glu	Ile	Ile 100	Ala	Asp	Ser	Val	Glu 105	Thr	Val	Val	Ser	Ala 110	His	Trp
Phe	Asp	Gly 115	Met	Val	Суѕ	Ile	Pro 120	Asn	Cys	Asp	Lys	Ile 125	Thr	Pro	Gly
Met	Leu 130	Met	Ala	Ala	Met	Arg 135	Ile	Asn	Ile	Pro	Thr 140	Ile	Phe	Val	Ser
Gly 145	Gly	Pro	Met	Ala	Ala 150	Gly	Arg	Thr	Ser	Tyr 155	Gly	Arg	Lys	Ile	Ser 160
Leu	Ser	Ser	Val	Phe 165	Glu	Gly	Val	Gly	Ala 170	Tyr	Gln	Ala	Gly	Lys 175	Ile
Asn	Glu	Asn	Glu 180	Leu	Gln	Glu	Leu	Glu 185	Gln	Phe	Gly	Cys	Pro 190	Thr	Cys
Gly	Ser	Cys 195	Ser	Gly	Met	Phe	Thr 200	Ala	Asn	Ser	Met	Asn 205	Суѕ	Leu	Ser
Glu	Ala 210	Leu	Gly	Leu	Ala	Leu 215	Pro	Gly	Asn	Gly	Thr 220	Ile	Leu	Ala	Thr
Ser 225	Pro	Glu	Arg	Lys	Glu 230	Phe	Val	Arg	Lys	Ser 235	Ala	Ala	Gln	Leu	Met 240
Glu	Thr	Ile	Arg	Lys 245	Asp	Ile	Lys	Pro	Arg 250	Asp	Ile	Val	Thr	Val 255	Lys
Ala	Ile	Asp	Asn 260	Ala	Phe	Ala	Leu	Asp 265	Met	Ala	Leu	Gly	Gly 270	Ser	Thr
Asn	Thr	Val 275	Leu	His	Thr	Leu	Ala 280	Leu	Ala	Asn	Glu	Ala 285	Gly	Val	Glu
Tyr	Ser 290	Leu	Glu	Arg	Ile	Asn 295	Glu	Val	Ala	Glu	Arg 300	Val	Pro	His	Leu
Ala 305	Lys	Leu	Ala	Pro	Ala 310	Ser	Asp	Val	Phe	Ile 315	Glu	Asp	Leu	His	Glu 320
Ala	Gly	Gly	Val	Ser 325	Ala	Ala	Leu	Asn	Glu 330	Leu	Ser	Lys	Lys	Glu 335	Gly
Ala	Leu	His	Leu 340	Asp	Ala	Leu	Thr	Val 345	Thr	Gly	Lys	Thr	Leu 350	Gly	Glu
Thr	Ile	Ala 355	Gly	His	Glu	Val	Lys 360	Asp	Tyr	Asp	Val	Ile 365	His	Pro	Leu
Asp	Gln 370	Pro	Phe	Thr	Glu	Lys 375	Gly	Gly	Leu	Ala	Val	Leu	Phe	Gly	Asn

PCT/US00/25993

Leu Ala Pro Asp Gly Ala Ile Ile Lys Thr Gly Gly Val Gln Asn Gly 385 Ile Thr Arg His Glu Gly Pro Ala Val Val Phe Asp Ser Gln Asp Glu Ala Leu Asp Gly Ile Ile Asn Arg Lys Val Lys Glu Gly Asp Val Val Ile Ile Arg Tyr Glu Gly Pro Lys Gly Gly Pro Gly Met Pro Glu Met Leu Ala Pro Thr Ser Gln Ile Val Gly Met Gly Leu Gly Pro Lys Val Ala Leu Ile Thr Asp Gly Arg Phe Ser Gly Ala Ser Arg Gly Leu Ser 470 Ile Gly His Val Ser Pro Glu Ala Ala Glu Gly Gly Pro Leu Ala Phe Val Glu Asn Gly Asp His Ile Ile Val Asp Ile Glu Lys Arg Ile Leu 500 505 Asp Val Gln Val Pro Glu Glu Glu Trp Glu Lys Arg Lys Ala Asn Trp 520 Lys Gly Phe Glu Pro Lys Val Lys Thr Gly Tyr Leu Ala Arg Tyr Ser Lys Leu Val Thr Ser Ala Asn Thr Gly Gly Ile Met Lys Ile <210> 39 <211> 194 <212> DNA <213> Artificial Sequence <220> <223> Description of Artificial Sequence:promoter sequence <220> <221> -35 signal <222> (136)..(141) <220> <221> -10 signal <222> (159)..(164) · <400> 39 gctattgacg acagctatgg ttcactgtcc accaaccaaa actgtgctca gtaccgccaa 60 tatttctccc ttgaggggta caaagaggtg tccctagaag agatccacgc tgtgtaaaaa 120

010177010 14.

```
gcaaccccgc ctqt
                                                                    194
<210> 40
<211> 163
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence:promoter
      sequence
<220>
<221> -35 signal
<222> (113)..(118)
<220>
<221> -10 signal
<222> (136)..(141)
<400> 40
gcctacctag cttccaagaa agatatccta acagcacaag agcggaaaga tgttttgttc 60
tacatccaga acaacctctg ctaaaattcc tgaaaaattt tgcaaaaagt tgttgacttt 120
atctacaagg tgtggtataa taatcttaac aacagcagga cgc
                                                                    163
<210> 41
<211> 127
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence:promoter
      sequence
<220>
<221> -35_signal
<222> (34)..(39)
<220>
<221> -10_signal
<222> (58)..(63)
<220>
<221> -35_signal
<222> (75)..(80)
<220>
<221> -10_signal
<222> (98)..(103)
gaggaatcat agaattttgt caaaataatt ttattgacaa cgtcttatta acgttgatat 60
aatttaaatt ttatttgaca aaaatgggct cgtgttgtac aataaatgta gtgaggtgga 120
```

PCT/US00/25993

tgcaatg '	. 121
<210> 42	
<211> 24	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence:ribosome binding site	
<400> 42	24
taaacatgag gaggagaaaa catg	
<210> 43	
<211> 28	
<212> DNA <213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: ribosome binding site	•
<400> 43	. 28
attcgagaaa tggagagaat ataatatg	.20
<210> 44	
<211> 13	
<212> DNA <213> Artificial Sequence	
<220>	·
<223> Description of Artificial Sequence:ribosome binding site	
<400> 44	12
agaaaggagg tga	. 13
<210> 45	
<211> 23	
<212> DNA <213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: ribosome binding site	
<400> 45	22
ttaagaaagg aggtgannnn atg	23
<210> 46	
<211> 23	
<212> DNA	
<213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome	
(572) Describition of writingtal pedremontagement	

binding site

<400> 46 ttagaaagga ggtgannnnn atg	23
<210> 47 <211> 23 <212> DNA <213> Artificial Sequence	,
<220> <223> Description of Artificial Sequence:ribosome binding site	,
<400> 47 agaaaggagg tgannnnnn atg	23
<210> 48 <211> 22 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 48 agaaaggagg tgannnnna tg	22
<210> 49 <211> 25 <212> DNA <213> Artificial Sequence	•
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 49 ccctctagaa ggaggagaaa acatg	25
<210> 50 <211> 24 <212> DNA <213> Artificial Sequence	
<220> <223> Description of Artificial Sequence:ribosome binding site	
<400> 50 ccctctagag gaggagaaaa catg	24
<210> 51 <211> 23 <212> DNA	

<213>	Artificial Sequence		
<220> <223>	Description of Artificial Sequence:ribosome binding site		•
<400> ttagaa	51 aagga ggatttaaat atg		23
<210> <211> <212> <213>	23		
<220> <223>	Description of Artificial Sequence:ribosome binding site		•
<400> ttaga	52 aagga ggtttaatta atg		23
<210><211><211><212><213>	· 23	• •	
<220> <223>	Description of Artificial Sequence:ribosome binding site		
<400> ttaga	> 53 aaagga ggtgatttaa atg		23
<220: <223:	> > Description of Artificial Sequence:ribosome binding site		
<400 ttag	> 54 aaagga ggtgtttaaa atg		-23
<220 <223	<pre>>> > Description of Artificial Sequence:ribosome binding site</pre>		
)> 55 :gagaaa ggaggtgaat ataatatg		28
4010	n		

```
<211> 27
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence:ribosome
      binding site
<400> 56
attcgagaaa ggaggtgaat aataatg
                                                                   27
<210> 57
<211> 28
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: ribosome
      binding site
<400> 57
attcgtagaa aggaggtgaa ttaatatg
                                                                   28
<210> 58
<211> 3291
<212> DNA
<213> Bacıllus subtilis
atggggacta atgtacaggt ggattcagca tctgccgaat gtacacagac gatgagcgga 60
gcattaatgc tgattgaatc attaaaaaaa gagaaagtag aaatgatctt cggttatccg 120
ggcggggctg tgcttccgat ttacgataag ctatacaatt cagggttggt acatatcctt 180
ccccgtcacg aacaaggagc aattcatgca gcggagggat acgcaagggt ctccggaaaa 240
ccgggtgtcg tcattgccac gtcagggccg ggagcgacaa accttgttac aggccttgct 300
gatgccatga ttgattcatt gccgttagtc gtctttacag ggcaggtagc aacctctgta 360
atcgggagcg atgcatttca ggaagcagac attttaggga ttacgatgcc agtaacaaaa 420
cacagetace aggttegeea geeggaagat etgeegegea teattaaaga agegtteeat 480
attgcaacaa ctggaagacc cggacctgta ttgattgata ttccgaaaga tgtagcaaca 540
attgaaggag aattcagcta cgatcatgag atgaatctcc cgggatacca gccgacaaca 600
gagccgaatt atttgcagat ccgcaagctt gtggaagccg tgagcagtgc gaaaaaaccg 660
gtgatcctgg cgggtgcggg cgtactgcac ggaaaagcgt cagaagaatt aaaaaattat 720
gctgaacagc agcaaatccc tgtggcacac acccttttgg ggctcggagg cttcccggct 780
gaccateege tttteetagg gatggeggga atgeaeggta ettataeage caatatggee 840
```

cttcatgaat gtgatctatt aatcagtatc ggcgcccgtt ttgatgaccg tgtcacagga 900 aacctgaaac actttgccag aaacgcaaag atagcccaca tcgatattga tccagctgaa 960, atoggaaaaa toatgaaaac acagattoot gtagtoggag acagcaaaat tgtootgcag 1020 gagetgatea aacaagaegg caaacaaage gatteaageg aatggaaaaa acagetegea 1080 gaatggaaag aagagtatcc gctctggtat gtagataatg aagaagaagg ttttaaacct 1140 cagaaattga ttgaatatat tcatcaattt acaaaaggag aggccattgt cgcaacggat 1200 gtaggccage atcaaatgtg gtcagcgcaa ttttatccgt tccaaaaagc agataaatgg 1260 gtcacgtcag gcggacttgg aacgatggga ttcggtcttc cggcggcgat cggcgcacag 1320 ctggccgaaa aagatgctac tgttgtcgcg gttgtcggag acggcggatt ccaaatgacg 1380 cttcaagaac tcgatgttat tcgcgaatta aatcttccgg tcaaggtagt gattttaaat 1440 aacgettgte teggaatggt cagacagtgg caggaaattt tetatgaaga acgttattea 1500 gaatctaaat tegettetea geetgaette gteaaattgt eegaageata eggeattaaa 1560 ggcatcagaa tttcatcaga agcggaagca aaggaaaagc tggaagaggc attaacatca 1620 agagaacctg ttgtcattga cgtgcgggtt gccagcgaag aaaaagtatt cccgatggtg 1680 gctccgggga aagggctgca tgaaatggtg ggggtgaaac cttgaaaaga attatcacat 1740 tgactgtggt gaaccgctcc ggggtgttaa accggatcac cggtctattc acaaaaaggc 1800 attacaacat tgaaagcatt acagttggac acacagaaac agccggcgtt tccagaatca 1860 ccttcgtcgt tcatgttgaa ggtgaaaatg atgttgaaca gttaacgaaa cagctcaaca 1920. aacagattga tgtgctgaaa gtcacagaca tcacaaatca atcgattgtc cagagggagc 1980 tggccttaat caaggttgtc tccgcacctt caacaagaac agagattaat ggaatcatag 2040 aaccgtttag agcctctgtc gttgatgtca gcagagacag catcgttgtt caggtgacag 2100 gtgaatctaa caaaattgaa gcgcttattg agttattaaa accttatggc attaaagaaa 2160 togogagaac aggtacaacg gottttgoga ggggaaccag caaaaggogt catccaataa 2220 aacaatatct attgtataaa acataacaag ggagagattg aaatggtaaa agtatattat 2280 aacggtgata tcaaagagaa cgtattggct ggaaaaacag tagcggttat cgggtacggt 2340 tegeaaggee acgeacatge cetgaacett aaagaaageg gagtagaegt gategteggt 2400 gttagacaag gaaaatcttt cactcaagcc caagaagacg gacataaagt attttcagta 2460 aaagaagegg cageecaage egaaateate atggttetge tteeggatga geageageaa 2520 aaagtatacg aagctgaaat caaagatgaa ttgacagcag gaaaatcatt agtattcgct 2580

- ----- ...-

catggattta acgtgcattt ccatcaatt gttcctccgg cggatgtaga tgtattctta 2640 gtggccccta aaggcccggg acacttggta agaagaacat atgagcaagg agctggcgta 2700 cctgcattgt tcgcaatcta tcaagatgtg actggagaag caagagacaa agccctcgct 2760 tatgctaaag gaatcggcgg cgcaagagcg ggcgtattag aaacgacatt taaagaagaa 2820 acagaaacag atttgttcgg tgagcaagca gttctttgcg gcggattaag cgcgcttgtc 2880 aaagccggat ttgaaacctt aactgaagca ggttatcagc ctgaacttgc atacttcgag 2940 tgtcttcatg agctgaaatt aatcgtagac cttatgtacg aagaaggact tgcaggaatg 3000 agatattcaa tctctgacac agcacagtgg ggagatttcg tatcaagccc tcgcgttgtg 3060 gacgccaaag taaaagaatc tatgaaagaa gtattaaaag atatccaaaa cggtacattc 3120 gcaaaagag ggatcgtcga aaaccaagta aaccgtcctc gtttcaacgc tatcaatgca 3180 agcgagaacg aacatcaaat cgaagtagtg ggaagaaagc ttcgtgaaat gatgccgttt 3240 gtgaaacaag gcaagaagaa ggaagcggtg gtctccgttg cgcaaaatta a 3291

```
<210> 59
<211> 2363
<212> DNA
<213> Bacillus subtilis
<220>
<221> CDS
<222> (242)..(1072)
<220> ·
<221> CDS
<222> (1077)..(1934)
<220>
<221> CDS
<222> (1939)..(2319)
<400> 59
ttggtacaag cccgttgatt ttggtatact tccattgggc aqtatcgcct gcgaactgca 60
cctattatta aaatagatag acattgcagc agtctgcctt gatccaaaaa aggactgqqa 120
cagagggatg aaactcgccg aactttagaa agtgaagaat ccttctcgtt gtaacggaag 180
gttttttggc ttgcagaaga aaacggcaga tcatctcctc taaacatgag gaggagaaaa 240
c atg aaa aca aaa ctg gat ttt cta aaa atg aag gag tct gaa gaa ccg 289
  Met Lys Thr Lys Leu Asp Phe Leu Lys Met Lys Glu Ser Glu Glu Pro
```

att gtc atg ctg acc gct tat gat tat ccg gca gct aaa ctt gct gaa 337 Ile Val Met Leu Thr Ala Tyr Asp Tyr Pro Ala Ala Lys Leu Ala Glu

			20		1	,		25					30			
caa Gln	gcg Ala	gga Gly 35	gtt Val	gac Asp	atg Met	att Ile	tta Leu 40	gtc Val	ggt Gly	gat Asp	tca Ser	ctt Leu 45	gga Gly	atg Met	gt <i>c</i> Val	385
gtc Val	ctc Leu 50	ggc Gly	ctt Leu	gat Asp	tca Ser	act Thr 55	gtc Val	ggt Gly	gtg Val	aca Thr	gtt Val 60	gcg Ala	gac Asp	atg Met	atc Ile	433
cat His 65	cat His	aca Thr	aaa Lys	gcc Ala	gtt Val 70	aaa Lys	agg Arg	ggt Gly	gcg Ala	ccg Pro 75	aat Asn	acc Thr	ttt Phe	att Ile	gtg Val 80	481
aca Thr	gat Asp	atg Met	ccg Pro	ttt Phe 85	atg Met	tct Ser	tat Tyr	cac His	ctg Leu 90	tct Ser	aag Lys	gaa Glu	gat Asp	acg Thr 95	ctg Leu	529
aaa Lys	aat Asn	gca Ala	gcg Ala 100	gct Ala	atc Ile	gtt Val	cag Gln	gaa Glu 105	agc Ser	gga Gly	gct Ala	gac Asp	gca Ala 110	ctg Leu	aag Lys	577
ctt Leu	gag Glu	ggc Gly 115	gga Gly	gaa Glu	ggc Gly	gtg Val	ttt Phe 120	gaa Glu	tcc Ser	att Ile	cgc Arg	gca Ala 125	ttg Leu	acg Thr	ctt Leu	625
gga Gly	ggc Gly 130	att Ile	cca Pro	gta Val	gtc Val	agt Ser 135	cac His	tta Leu	ggt Gly	ttg Leu	aca Thr 140	Pro	cag Gln	tca Ser	gtc Val	673
ggc Gly 145	Val	ctg Leu	ggc	ggc	tat Tyr 150	aaa Lys	gta Val	cag Gln	ggc	aaa Lys 155	Asp	gaa Glu	caa Gln	agc Ser	gcc Ala 160	721
aaa Lys	aaa Lys	tta Leu	ata Ile	gaa Glu 165	gac Asp	agt Ser	ata Ile	aaa Lys	tgc Cys 170	Glu	gaa Glu	gca Ala	gga Gly	gct Ala 175	Met	769
at <u>c</u> Met	ctt Leu	gtg Val	cto Lev 180	ı Glu	tgt Cys	gtg Val	-ccg Pro	gca Ala 185	Glu	ctc Leu	aca Thi	gcc Ala	aaa Lys 190	Ile	gcc Ala	817
gaq Glu	g aco	Leu 195	ı Ser	ata Ile	Pro	gtc Val	att Ile 200	: Gly	ato / Ile	ggg Gly	get Ala	ggt a Gly 205	/ Val	r aaa . Lys	gcg Ala	865
gad Ası	gga Gly 210	/ Glr	a gtt n Val	t ctc l Lev	gtt Val	tat Tyr 215	His	gat Asp	att Ile	ato	gg(Gl) 220	y His	c ggt s Gly	gtt Val	gag Glu	913
aga Are 22	g Thi	a cci	t aaa o Ly:	a ttt s Phe	gta Val 230	Lys	caa Glr	a tat 1 Tyl	acq Thi	g ego Arc 235	; Ile	t gat e Ası	t gaa o Glu	a aco	atc : Ile 240	961
ga Gl	a aca u Thi	a gca r Ala	a ate	c ago e Sei	G13	tat Tyr	gtt Val	caq L Gl	g gat n Asp 250	o Val	a ag	a ca g Hi	t cgi	g Ala	t ttc a Phe	1009

	cct Pro	gaa Glu	caa Gln	aag Lys 260	cat His	tcc Ser	ttt Phe	caa Gln	atg Met 265	aac Asn	cag Gln	aca Tḥr	gtg Val	ctt Leu 270	gac Asp	ggc Gly	1057
ı	ttg Leu	tac Tyr	999 Gly 275	gga Gly	aaa Lys	taa	g ato Me	g aga	a cad g Gli 280	n Ile	t ac e Th	t ga r As	t at p Il	t tc e Se 28	r Gl	g ctg n Leu	1106
	aaa Lys	gaa Glu	gcc Ala 290	ata Ile	aaa Lys	caa Gln	tac Tyr	cat His 295	tca Ser	gag Glu	ggc	aag Lys	tca Ser 300	atc Ile	gga Gly	ttt Phe	1154
	gtt Val	ccg Pro 305	acg Thr	atg Met	ggg Gly	ttt Phe	ctg Leu 310	cat His	gag Glu	G1 y ggg	cat His	tta Leu 315	acc Thr	tta Leu	gca Ala	gac Asp	1202
	aaa Lys 320	gca Ala	aga Arg	caa Gln	gaa Glu	aac Asn 325	gac Asp	gcc Ala	gtt Val	att Ile	atg Met 330	agt Ser	att Ile	ttt Phe	gtg Val	aat Asn 335	1250
	cct Pro	gca Ala	caa Gln	ttc Phe	ggc Gly 340	cct Pro	aat Asn	gaa Glu	gat Asp	ttt Phe 345	gaa Glu	gca Ala	tat 'Tyr	ccg Pro	cgc Arg 350	gat Asp	1298
	att Ile	gag Glu	cgg Arg	gat Asp 355	gca Ala	gct Ala	ctt Leu	gca Ala	gaa Glu 360	aac Asn	gcc Ala	gga Gly	gtc Val	gat Asp 365	att Ile	ctt Leu	1346
	ttt Phe	acg Thr	cca Pro 370	gat Asp	gct Ala	cat His	gat Asp	atg Met 375	tat Tyr	ccc Pro	ggt Gly	gaa Glu	aag Lys 380	aat Asn	gtc Val	acg Thr	1394
	att Ile	cat His 385	gta Val	gaa Glu	aga Arg	cgc Arg	Thr	gac Asp	gtg Val	tta Leu	tgc Cys	ggg Gly 395	cgc Arg	tca Ser	aga Arg	gaa Glu	1442
	gga Gly 400	cat His	ttt Phe	gac Asp	Gly ggg	gtc Val 405	gcg Ala	atc Ile	gta Val	ctg Leu	acg Thr 410	aag Lys	ctt Leu	ttc Phe	aat Asn	cta Leu 415	1490
	gtc Val	aag Lys	ccg Pro	act Thr	cgt Arg 420	Ala	Tyr	ttc Phe	Gly	Leu	Lys	Asp	gcg Ala	Gln	cag Gln 430	gta Val	1538
	gct Ala	gtt Val	gtt Val	gat Asp 435	Gly ggg	tta Leu	atc Ile	agc Ser	gac Asp 440	ttc Phe	ttc Phe	atg Met	gat Asp	att Ile 445	gaa Glu	ttg Leu	1586
	gtt Val	cct Pro	gtc Val 450	gat Asp	acg Thr	gtc Val	aga Arg	gag Glu 455	gaa Glu	gac Asp	ggc Gly	tta Leu	gcc Ala 460	aaa Lys	agc Ser	tct Ser	1634
	cgc Arg	aat Asn 465	gta Val	tac Tyr	tta Leu	aca Thr	gct Ala 470	gag Glu	gaa Glu	aga Arg	aaa Lys	gaa Glu 475	gcg Ala	cct Pro	aag Lys	ctg Leu	1682
	tat Tyr	cgg Arg	gcc Ala	ctt Leu	caa Gln	aca Thr	agt Ser	gcg Ala	gaa Glu	ctt Leu	gtc Val	caa Gln	gcc Ala	ggt Gly	gaa Glu	aga Arg	1730

480					485					490	1				495	
gat Asp	cct Pro	gaa Glu	gcg Ala	gtg Val 500	ata Ile	aaa Lys	gct Ala	gca Ala	aaa Lys 505	gat Asp	atc Ile	att Ile	gaa Glu	acg Thr 510	act Thr	1778
agc Ser	gga Gly	acc Thr	ata Ile 515	gac Asp	tat Tyr	gta Val	gag Glu	ctt Leu 520	tat Tyr	tcc Ser	tat Tyr	ccg Pro	gaa, Glu 525	ctc Leu	gag Glu	1826
cct Pro	gtg Val	aat Asn 530	gaa Glu	att Ile	gct Ala	gga Gly	aag Lys 535	atg Met	att Ile	ctc Leu	gct Ala	gtt Val 540	gca Ala	gtt Val	gct Ala	1874
ttt Phe	tca Ser 545	aaa Lys	gcg Ala	cgt Arg	tta Leu	ata Ile 550	gat Asp	aat Asn	atc Ile	att Ile	att Ile 555	gat Asp	att Ile	cga Arg	gaa Glu	1922
_	gag Glu	_		taat	ate Me	g ta t Ty: 56	r Ar	a ac g Th	a at r Me	g at t Me	g age t Se 57	r Gl	c aa y Ly	a ct s Le	t ∘cac u His	1971
agg Arg 575	Āla	act Thr	gtt Val	acg Thr	gaa Glu 580	Ala	aac Asn	ctg Leu	aac Asn	tat Tyr 585	Val	gga Gly	agc Ser	att Ile	aca Thr 590	2019
att Ile	gat Asp	gaa Glu	gat Asp	ctc Leu 595	att Ile	gat Asp	gct Ala	gtg Val	gga Gly 600	Met	ctt Leu	cct Pro	aat Asn	gaa Glu 605	aaa Lys	2067
gta Val	caa Gln	att Ile	gtg Val 610	Asn	aat Asn	aat Asn	aat Asn	gga Gly 615	Ala	cgt Arc	ctt Leu	gaa Glu	acg Thr 620	Туг	att Ile	2115
att Ile	cct Pro	ggt Gly 625	' Lys	cgg Arg	gga	ago Ser	ggc Gly 630	v Val	ata Ile	tgo Cys	tta Lev	aac Asr 635	GJ?	gca Ala	a gcc a Ala	2163
gca	a cgc Arc 640	Let	gto Val	cag Gln	gaa Glu	gga Gly 645	Asp	aaq Lys	g gto s Val	att I Ile	att = Ile 650	: Ile	te Ser	tac Ty	c aaa c Lys	2211
ato Met 655	: Met	tct Sei	gat Asp	caa Glr	gaa Glu 660	ı Ala	g gca a Ala	a Sei	c cat	gaç 661 661	u Pro	g aaa D Lys	a gto s Val	g gci	t gtt a Val 670	2259
cto Lei	g aat ı Ası	gat n Ası	caa o Glr	a aad n Asr 675	Lys	a att s Ile	t gaa e Glu	a caa	a ato n Med 680	t Le	g ggg u Gl	g aa y Asi	c gaa n Glu	a cc u Pr 68	a gcc o Ala 5	2307
	t aca			ı	gaaga	aaaa	gcc	ccct	tta :	tcgg	gggt	tt t	cttt	taag	a ttti	2363

<210> 60 <211> 293 <212> PRT <213> Bacillus subtilis

<400> 60

Met Ser Ile Ala Val Ser Glu Glu Glu Ala Lys Ala Val Glu Gly Leu
1 10 15

Asn Asp Tyr Leu Ser Val Glu Glu Val Glu Thr Ile Tyr Ile Pro Leu 20 25 30

Val Arg Leu Leu His Leu His Val Lys Ser Ala Ala Glu Arg Asn Lys
35 40 45

His Val Asn Val Phe Leu Lys His Pro His Ser Ala Lys Ile Pro Phe 50 55 60

Ile Ile Gly Ile Ala Gly Ser Val Ala Val Gly Lys Ser Thr Thr Ala 65 70 75 80

Årg Ile Leu Gln Lys Leu Leu Ser Arg Leu Pro Asp Arg Pro Lys Val 85 90 95

Ser Leu Ile Thr Thr Asp Gly Phe Leu Phe Pro Thr Ala Glu Leu Lys
100 105 110

Lys Lys Asn Met Met Ser Arg Lys Gly Phe Pro Glu Ser Tyr Asp Val 115 120 125

Lys Ala Leu Leu Glu Phe Leu Asn Asp Leu Lys Ser Gly Lys Asp Ser 130 135 140

Val Lys Ala Pro Val Tyr Ser His Leu Thr Tyr Asp Arg Glu Glu Gly 145 150 155 160

Val Phe Glu Val Val Glu Gln Ala Asp Ile Val Ile Ile Glu Gly Ile 165 170 175

Asn Val Leu Gln Ser Pro Thr Leu Glu Asp Asp Arg Glu Asn Pro Arg 180 185 190

Ile Phe Val Ser Asp Phe Phe Asp Phe Ser Ile Tyr Val Asp Ala Glu 195 200 205

Glu Ser Arg Ile Phe Thr Trp Tyr Leu Glu Arg Phe Arg Leu Leu Arg 210 215 220

Glu Thr Ala Phe Gln Asn Pro Asp Ser Tyr Phe His Lys Phe Lys Asp 225 230 235 240

Leu Ser Asp Gln Glu Ala Asp Glu Met Ala Ala Ser Ile Trp Glu Ser 245 250 255

Val Asn Arg Pro Asn Leu Tyr Glu Asn Ile Leu Pro Thr Lys Phe Arg 260 265 270

Ser Asp Leu Ile Leu Arg Lys Gly Asp Gly His Lys Val Glu Glu Val 275 280 285

Leu Val Arg Arg Val

290

<210> 61

<211> 281

<212> PRT

<213> Bacillus subtilis

<400> 61

Met Glu Gly Leu Asn Asp Tyr Leu Ser Val Glu Glu Val Glu Thr Ile 1 5 10 15

Tyr Ile Pro Leu Val Arg Leu Leu His Leu His Val Lys Ser Ala Ala 20 25 30

Glu Arg Asn Lys His Val Asn Val Phe Leu Lys His Pro His Ser Ala 35 40 45

Lys Ile Pro Phe Ile Ile Gly Ile Ala Gly Ser Val Ala Val Gly Lys 50 55 60

Ser Thr Thr Ala Arg Ile Leu Gln Lys Leu Leu Ser Arg Leu Pro Asp 65 70 75 80

Arg Pro Lys Val Ser Leu Ile Thr Thr Asp Gly Phe Leu Phe Pro Thr 85 90 95

Ala Glu Leu Lys Lys Lys Asn Met Met Ser Arg Lys Gly Phe Pro Glu 100 105 110

Ser Tyr Asp Val Lys Ala Leu Leu Glu Phe Leu Asn Asp Leu Lys Ser 115 120 125

Gly Lys Asp Ser Val Lys Ala Pro Val Tyr Ser His Leu Thr Tyr Asp 130 135 140

Arg Glu Glu Gly Val Phe Glu Val Val Glu Gln Ala Asp Ile Val Ile 145 150 155 160

Ile Glu Gly Ile Asn Val Leu Gln Ser Pro Thr Leu Glu Asp Asp Arg 165 170 175

Glu Asn Pro Arg Ile Phe Val Ser Asp Phe Phe Asp Phe Ser Ile Tyr 180 185 190

Val Asp Ala Glu Glu Ser Arg Ile Phe Thr Trp Tyr Leu Glu Arg Phe 195 200 205

Arg Leu Leu Arg Glu Thr Ala Phe Gln Asn Pro Asp Ser Tyr Phe His 210 215 220

Lys Phe Lys Asp Leu Ser Asp Gln Glu Ala Asp Glu Met Ala Ala Ser 225 230 235 240

Ile Trp Glu Ser Val Asn Arg Pro Asn Leu Tyr Glu Asn Ile Leu Pro 245 250 255

Thr Lys Phe Arg Ser Asp Leu Ile Leu Arg Lys Gly Asp Gly His Lys

.

260

270

- 61 -

265

Val Glu Val Leu Val Arg Arg Val <210> 62 <211> 1092 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(1089) <400> 62 atg act aaa caa aca att cgc gtt gaa ttg aca tca aca aaa aaa ccg 48 Met Thr Lys Gln Thr Ile Arg Val Glu Leu Thr Ser Thr Lys Lys Pro 10 aaa cca gac cca aat cag ctt tcg ttc gga aga gtg ttt aca gac cac 96 Lys Pro Asp Pro Asn Gln Leu Ser Phe Gly Arg Val Phe Thr Asp His atg ttt gta atg gac tat gcc gca gat aaa ggt tgg tac gat cca aga Met Phe Val Met Asp Tyr Ala Ala Asp Lys Gly Trp Tyr Asp Pro Arg 40 atc att cct tat caa ccc tta tca atg gat cca act gca atg gtc tat 192 Ile Ile Pro Tyr Gln Pro Leu Ser Met Asp Pro Thr Ala Met Val Tyr cac tac ggc caa acc gtg ttt gaa ggg tta aag gct tac gtg tca gag 240 His Tyr Gly Gln Thr Val Phe Glu Gly Leu Lys Ala Tyr Val Ser Glu gat gac cat gtt ctg ctt ttc aga ccg gaa aaa aat atg gaa cgc ctg Asp Asp His Val Leu Leu Phe Arg Pro Glu Lys Asn Met Glu Arg Leu 90 aat caa tca aac gac cgc ctc tgc atc ccg caa att gat gaa gaa cag 336 Asn Gln Ser Asn Asp Arg Leu Cys Ile Pro Gln Ile Asp Glu Glu Gln 100 gtt ctt gaa ggc tta aag cag ctt gtc gca att gat aaa gac tgg att 384 Val Leu Glu Gly Leu Lys Gln Leu Val Ala Ile Asp Lys Asp Trp Ile 120 cca aat gcg gag ggc acg tcc ctt tac atc cgt ccg ttc atc atc gca 432 Pro Asn Ala Glu Gly Thr Ser Leu Tyr Ile Arg Pro Phe Ile Ile Ala acc gag cct ttc ctt ggt gtt gcg gca tct cat acg tat aag ctc ttg 480 Thr Glu Pro Phe Leu Gly Val Ala Ala Ser His Thr Tyr Lys Leu Leu 150 155 atc att ctt tct ccg gtc ggc tct tat tac aaa gaa ggc att aag ccg 528 Ile Ile Leu Ser Pro Val Gly Ser Tyr Tyr Lys Glu Gly Ile Lys Pro

- 62 -

				165					170		•			175		
gtc Val	aaa Lys	atc Ile	gct Ala 180	gtt Val	gaa Glu	agt Ser	gaa Glu	ttt Phe 185	gtc Val	cgt Arg	gcg Ala	gta Val	aaa Lys 190	ggc	gga Gly	576
aca Thr	gga Gly	aat Asn 195	gcc Ala	aaa Lys	acc Thr	gca Ala	gga Gly 200	aac Asn	tat Tyr	gct Ala	tca Ser	agc Ser 205	tta Leu	aaa Lys	gcg Ala	624
cag Gln	cag Gln 210	gta Val	gcc Ala	gaa Glu	gag Glu	aaa Lys 215	gga Gly	ttt Phe	tct Ser	caa Gln	gta Val 220	ctc Leu	tgg Trp	ctg Leu	gac Asp	672
ggc Gly 225	att Ile	gag Glu	aag Lys	aaa Lys	tac Tyr 230	atc Ile	gaa Glu	gaa Glu	gtc Val	gga Gly 235	agc Ser	atg Met	aac Asn	atc Ile	ttc Phe 240	720
ttc Phe	aaa Lys	atc Ile	aac Asn	ggt Gly 245	gaa Glu	atc Ile	gta Val	aca Thr	ccg Pro 250	atg Met	ctg Leu	aac Asn	ggg Gly	agc Ser 255	atc Ile	768
ctg Leu	gaa Glu	ggc	att Ile 260	Thr	ege Arg	aat Asn	tca Ser	Val 265	atc Ile	gcc Ala	ttg Leu	ctt Leu	aag Lys 270	cat His	tgg Trp	816
ggc Gly	ctt Leu	caa Gln 275	Val	tca Ser	gaa Glu	cga Arg	aaa Lys 280	Ile	gcg Ala	atc Ile	gat Asp	gag Glu 285	Val	atc Ile	caa "Gln	864
gcc Ala	cat His 290	Lys	gac Asp	ggc Gly	atc Ile	ctg Leu 295	Glu	gaa Glu	gcc Ala	ttc Phe	gga Gly 300	Thr	ggt Gly	aca Thr	gca Ala	912
gct Ala 305	Val	att Ile	tco Ser	cca Pro	gto Val 310	Gly	gag Glu	ctg Lev	atc Ile	tgg Trp 315	Gla	gat Asp	gaa Glu	aca Thr	Ctt Leu 320	9'60
tcg Ser	ato :Ile	aac Asr	aad Asr	ggt Gly 325	Glu	aca Thr	Gly	gaa Glu	ato 11e 330	Ala	a aaa a Lys	aaa Lys	a cta s Lev	a tat ı Tyr 335	gac Asp	1008
acg Thr	att : Ile	aca Th:	a ggo r Gly 340	y Ile	caa Glr	a aaa a Lys	a ggo s Gly	gct Ala 345	a Val	e gca L Ala	a gad a Asg	gaa Glu	a tto 1 Phe 350	s GT?	tgg Trp	1056
acc Thr	g aco	ga G1: 35	u Va.	c gca l Ala	a gco a Ala	g cto a Leo	g act I Thi 360	c Glu	a ago u Sei	c aaq	g taa	a				1092

<210> 63 <211> 363 <212> PRT

<213> Bacillus subtilis

<400> 63 Met Thr Lys Gln Thr Ile Arg Val Glu Leu Thr Ser Thr Lys Lys Pro

1				5					10					15	
Lys	Pro	Asp	Pro 20	Asn	Gln	Leu	Ser	Phe 25	Gly	Arg	Val	Phe	Thr 30		His
Met	Phe	Val 35	Met	Åsp	Tyr	Ala	Ala 40	Asp	Lys	Gly	Trp	Tyr 45	Asp	Pro	Arg
Ile	Ile 50	Pro	Tyr	Gln	Pro	Leu 55	Ser	Met	Asp	Pro	Thr 60	Ala	Met	Val	Tyr
His 65	Tyr	Gly	Gln	Thr	Val 70	Phe	Glu	Gly	Leu	Lys 75	Ala	Tyr	Val	Ser	Glu 80
Asp	Asp	His	Val	Leu 85	Leu	Phe	Arg	Pro	Glu 90	Lys	Asn	Met	Glu	Arg 95	Leu
Asn	Gln	Ser	Asn 100	Asp	Arg	Leu	Cys	Ile 105	Pro	Gln	Ile	Asp	Glu 110	Glu	Gln
Val	Leu	Glu 115	Gly	Leu	Lys	Gln	Leu 120	Val	Ala	Ile	Asp	Lys 125	Asp	Trp	Ile
Pro	Asn 130	Ala	Glu	Gly	Thr	Ser 135	Leu	Tyr	Ile	Arg	Pro 140	Phe	Ile	Ile	Ala
Thr 145	Glu	Pro	Phe	Leu	Gly 150	Val	Ala	Ala	Ser	His 155	Thr	Tyr	Гуs	Leu	Leu 160
Ile	Ile	Leu	Ser	Pro 165	Val	Gly	Ser	Tyr	Tyr 170	Lys	Glu	Gly	Ile	Lys 175	Pro
Val	Lys	Ile	Ala 180	Val	Glu	Ser	Glu	Phe 185	Val	Arg	Ala	Val	Lys 190	Gly	Gly
Thr	Gly	Asn 195	Ala	Lys	Thr	Ala	Gly 200	Asn	Tyr	Ala	Ser	Ser 205	Leu	Lys	Ala
Gln	Gln 210	Val	Ala	Glu	Glu	Lys 215	Gly	Phe	Ser	Gln	Val 220	Leu	Trp	Leu	Asp
Gly 225	Ile	Glu	Lys	Lys	Tyr 230	Ile	Glu	Glu	Val	Gly 235	Ser	Met	Asn	Ile	Phe 240
Phe	Lys	Ile	Asn	Gly 245	Glu	Ile	Val	Thr	Pro 250	Met	Leu	Asn	Gly	Ser 255	Ile
Leu	Glu	Gly	Ile 260	Thr	Arg	Asn	Ser	Val 265	Ile	Ala	Leu	Leu	Lys 270	His	Trp
Gly	Leu	Gln 275	Val	Ser	Glu	Arg	Lys 280	Ile	Ala	Ile	Asp	Glu 285	Val	Ile	Gln
Ala	His 290	Lys	Asp	Gly	Ile	Leu 295	Glu	Glu	Ala	Phe	Gly 300	Thr	Gly	Thr	Ala
Ala 305	Val	Ile	Ser	Pro	Val 310	Gly	Glu	Leu	Ile	Trp 315	Gln	Asp	Glu	Thr	Leu 320

480

Ser Ile Asn Asn Gly Glu Thr Gly Glu Ile Ala Lys Lys Leu Tyr Asp Thr Ile Thr Gly Ile Gln Lys Gly Ala Val Ala Asp Glu Phe Gly Trp Thr Thr Glu Val Ala Ala Leu Thr Glu Ser Lys <210> 64 <211> 1071 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(1068) <400> 64 ttg aat aag ctt att gaa cga gaa aaa act gta tat tat aag gaa aag Met Asn Lys Leu Ile Glu Arg Glu Lys Thr Val Tyr Tyr Lys Glu Lys ccc gac ccg tot tcc ttg ggg ttt gga caa tat ttt aca gat tat atg Pro Asp Pro Ser Ser Leu Gly Phe Gly Gln Tyr Phe Thr Asp Tyr Met ttt gtg atg gac tac gaa gag ggg att gga tgg cat cat ccg aga att Phe Val Met Asp Tyr Glu Glu Gly Ile Gly Trp His His Pro Arg Ile gcg ccg tac gca ccg ctt acg ctt gat ccg tct tca tct gtt ttt cat 192 Ala Pro Tyr Ala Pro Leu Thr Leu Asp Pro Ser Ser Ser Val Phe His tac ggc cag gct gtt ttt gaa gga tta aaa gca tac aga aca gac gac 240 Tyr Gly Gln Ala Val Phe Glu Gly Leu Lys Ala Tyr Arg Thr Asp Asp ggc agg gtg ctg ctg ttc cgt ccg gat caa aat atc aaa cgg ctg aac 288 Gly Arg Val Leu Leu Phe Arg Pro Asp Gln Asn Ile Lys Arg Leu Asn 90 aga tcg tgt gag cgc atg agc atg ccc cct tta gac gaa gag ctg gtg 336 Arg Ser Cys Glu Arg Met Ser Met Pro Pro Leu Asp Glu Glu Leu Val 105 ctt gag gca ttg acg caa tta gtt gag ctg gag aaa gat tgg gtt cca 384 Leu Glu Ala Leu Thr Gln Leu Val Glu Leu Glu Lys Asp Trp Val Pro aag gaa aaa gga acg tca ctg tat att cgt cct ttt gtc att gcc aca 432 Lys Glu Lys Gly Thr Ser Leu Tyr Ile Arg Pro Phe Val Ile Ala Thr 135

gaa ccg agt ctc ggt gtg aag gca tcc agg agc tat aca ttt atg atc

Glu 145	Pro	Ser	Leu	Gly	Val 150	Lys	Ala	Ser	Arg	Ser 155	Tyr	Thr	Phe	Met	Ile 160	•
gtg Val	ctt Leu	tcg Ser	cct Pro	gtc Val 165	ggc Gly	tcc Ser	tat Tyr	tat Tyr	ggc Gly 170	gac Asp	gat Asp	cag Gln	ctg Leu	aag Lys 175	ccg Pro	528
gtt Val	aga Arg	atc Ile	tat Tyr 180	gtc Val	gaa Glu	gat Asp	gag Glu	tat Tyr 185	gtg Val	agg Arg	gcg Ala	gtc Val	aac Asn 190	gga Gly	gga Gly	576
gtc Val	GJ A Gaa	ttt Phe 195	gca Ala	aaa Lys	acg Thr	gct Ala	gga Gly 200	aac Asn	tat Tyr	gcc Ala	gcc Ala	agt Ser 205	ctt Leu	cag Gln	gca Ala	624
cag Gln	cgg Arg 210	aaa Lys	gcg Ala	aat Asn	gaa Glu	ctg Leu 215	ggc Gly	tat Tyr	gac Asp	cag Gln	gta Val 220	ctg Leu	tgg Trp	ctg Leu	gac Asp	672
gcc Ala 225	atc Ile	gaa Glu	aag Lys	aaa Lys	tat Tyr 230	gtg Val	gaa Glu	gaa Glu	gta Val	ggg Gly 235	agc Ser	atg Met	aac Asn	atc Ile	ttt Phe 240	720
ttc Phe	gtc Val	ata Ile	aac Asn	ggg Gly 245	gaa Glu	gct Ala	gtc Val	aca Thr	cct Pro 250	gct Ala	tta Leu	agc Ser	gga Gly	agc Ser 255	att Ile	768
tta Leu	agc Ser	Gl ^y ggg	gtt Val 260	aca Thr	cgt' Arg	gcg Ala	tct Ser	gcg Ala 265	att Ile	gaa Glu	ttg Leu	att Ile	cga Arg 270	agc Ser	tgg Trp	816
ggc Gly	att Ile	ccg Pro 275	gtt Val	cgt Arg	gaa Glu	gag Glu	aga Arg 280	ata Ile	tcg Ser	att Ile	gat Asp	gag Glu 285	gtg Val	tat Tyr	gcg Ala	864
gcc Ala	tct Ser 290	gca Ala	cgc Arg	gga Gly	gaa Glu	ttg Leu 295	aca Thr	gag Glu	gtc Val	ttt Phe	ggc Gly 300	aca Thr	ggc Gly	acg Thr	gca Ala	912
gca Ala 305	gtc Val	gtt Val	acg Thr	cct Pro	gtc Val 310	ggt Gly	gaa Glu	ctc Leu	aac Asn	atc Ile 315	cat His	gga Gly	aaa Lys	acg Thr	gtg Val 320	960
att Ile	gta Val	ggc	gac Asp	ggg Gly 325	caa Gln	atc Ile	GJ À GGG	gac Asp	ctc Leu 330	tcg Ser	aaa Lys	aag Lys	ctg Leu	tat Tyr 335	gaa Glu	1008
acg Thr	ata Ile	aca Thr	gat Asp 340	att Ile	cag Gln	ctt Leu	ggc Gly	aag Lys 345	gta Val	aaa Lys	ggc Gly	ccg Pro	ttt Phe 350	aac Asn	tgg Trp	1056
	gtg Val	-		tga												1071

<210> 65 <211> 356 <212> PRT <213> Bacillus subtilis

Add to the state of the st

Pro Asp Pro Ser Ser Leu Gly Phe Gly Gln Tyr Phe Thr Asp Tyr Met

Phe Val Met Asp Tyr Glu Glu Gly Ile Gly Trp His His Pro Arg Ile 35 40 45

Ala Pro Tyr Ala Pro Leu Thr Leu Asp Pro Ser Ser Ser Val Phe His
50 55 60

Tyr Gly Gln Ala Val Phe Glu Gly Leu Lys Ala Tyr Arg Thr Asp Asp 65 70 75 80

Gly Arg Val Leu Leu Phe Arg Pro Asp Gln Asn Ile Lys Arg Leu Asn 85 90 95

Arg Ser Cys Glu Arg Met Ser Met Pro Pro Leu Asp Glu Glu Leu Val

Leu Glu Ala Leu Thr Gln Leu Val Glu Leu Glu Lys Asp Trp Val Pro 115 120 125

Lys Glu Lys Gly Thr Ser Leu Tyr Ile Arg Pro Phe Val Ile Ala Thr 130 135 140

Glu Pro Ser Leu Gly Val Lys Ala Ser Arg Ser Tyr Thr Phe Met Ile 145 150 155 160

Val Leu Ser Pro Val Gly Ser Tyr Tyr Gly Asp Asp Gln Leu Lys Pro 165 170 175

Val Arg Ile Tyr Val Glu Asp Glu Tyr Val Arg Ala Val Asn Gly Gly 180 185 190

Val Gly Phe Ala Lys Thr Ala Gly Asn Tyr Ala Ala Ser Leu Gln Ala 195 200 205

Gln Arg Lys Ala Asn Glu Leu Gly Tyr Asp Gln Val Leu Trp Leu Asp 210 215 220

Ala Ile Glu Lys Lys Tyr Val Glu Glu Val Gly Ser Met Asn Ile Phe 225 230 235 240

Phe Val Ile Asn Gly Glu Ala Val Thr Pro Ala Leu Ser Gly Ser Ile 245 250 255

Leu Ser Gly Val Thr Arg Ala Ser Ala Ile Glu Leu Ile Arg Ser Trp 260 265 270

Gly Ile Pro Val Arg Glu Glu Arg Ile Ser Ile Asp Glu Val Tyr Ala 275 280 285

- 67 -

Ala Ser Ala Arg Gly Glu Leu Thr Glu Val Phe Gly Thr Gly Thr Ala ' 290 295 Ala Val Val Thr Pro Val Gly Glu Leu Asn Ile His Gly Lys Thr Val 310 Ile Val Gly Asp Gly Gln Ile Gly Asp Leu Ser Lys Lys Leu Tyr Glu 330 Thr Ile Thr Asp Ile Gln Leu Gly Lys Val Lys Gly Pro Phe Asn Trp 340 345 Thr Val Glu Val 355 <210> 66 <211> 1428 <212> DNA <213> Bacillus subtilis <220> <221> CDS <222> (1)..(1425) <400> 66 atg tta aac ggc caa aaa gaa tat cgc gtg gaa aaa gac ttc ctt ggg Met Leu Asn Gly Gln Lys Glu Tyr Arg Val Glu Lys Asp Phe Leu Gly gaa aaa caa att gaa gca gat gtt tat tac gga att cag acg ctc cgt Glu Lys Gln Ile Glu Ala Asp Val Tyr Tyr Gly Ile Gln Thr Leu Arg gct tct gaa aat ttt ccg atc aca gga tac aaa atc cat gag gaa atg 144 Ala Ser Glu Asn Phe Pro Ile Thr Gly Tyr Lys Ile His Glu Glu Met att aac gca ctg gcg att gtg aaa aaa gct gcg gct ctt gcc aac atg 192 Ile Asn Ala Leu Ala Ile Val Lys Lys Ala Ala Ala Leu Ala Asn Met gac gtg aaa cgg ctg tat gaa gga att ggc caa gct atc gta caa gcc 240 Asp Val Lys Arg Leu Tyr Glu Gly Ile Gly Gln Ala Ile Val Gln Ala 65 gct gac gag att ctg gaa ggc aag tgg cac gat cag ttt atc gtc gat 288 Ala Asp Glu Ile Leu Glu Gly Lys Trp His Asp Gln Phe Ile Val Asp ccg att cag ggc ggt gcc gga act tct atg aac atg aac gcg aat gag 336 Pro Ile Gln Gly Gly Ala Gly Thr Ser Met Asn Met Asn Ala Asn Glu 100 110 gtt atc gga aac cgg gcg ctt gaa atc atg gga cat aaa aag gga gat 384

Val Ile Gly Asn Arg Ala Leu Glu Ile Met Gly His Lys Lys Gly Asp

120

115

tat Tyr	atc Ile 130	cat His	tta Leu	agt Ser	cca' Pro	aac Asn 135	aca Thr	cat His	gtg Val	aac Asn	atg Met 140	tca Ser	cag Gln	tct Ser	cag Gln	432
aac Asn 145	gat Asp	gtg Val	ttc Phe	ccg Pro	act Thr 150	gct Ala	atc Ile	cat His	att Ile	tec Ser 155	aca Thr	ttg Leu	aag Lys	ctc Leu	tta Leu 160	480
gaa Glu	aaa Lys	ctg Leu	ctg Leu	aaa Lys 165	aca Thr	atg Met	gaa Glu	gat Asp	atg Met 170	cat His	agt Ser	gtg Val	ttt Phe	aaa Lys 175	caa Gln	528
aaa Lys	gca Ala	cag Gln	gag Glu 180	ttt Phe	cac His	tct Ser	gtt Val	att Ile 185	aaa Lys	atg Met	ggc Gly	egg Arg	aca Thr 190	cac His	ctt Leu	576
caa Gln	gat Asp	gcg Ala 195	gtt Val	ccg Pro	atc Ile	cgt Arg	ctt Leu 200	Gly	cag Gln	gaa Glu	ttc Phe	gaa Glu 205	gct Ala	tac Tyr	agc Ser	624
cgt Arg	gtt Val 210	ctc Leu	gag Glu	cgt Arg	gat Asp	atc Ile 215	aaa Lys	cga Arg	atc Ile	aag Lys	caa Gln 220	tcg Ser	cgc Arg	cag Gln	cac His	672
ctg Leu 225	tat Tyr	gaa Glu	gtc Val	aac Asn	atg Met 230	Gly	gca Ala	act Thr	gct Ala	gtt Val 235	Gly	aca Thr	GJ y ggg	ctg Leu	aac Asn 240	720
gct Ala	gat Asp	cct Pro	gaa Glu	tat Tyr 245	atc Ile	aaa Lys	cag Gln	gta Val	gta Val 250	aag L ys	cac	ctt Leu	gct Ala	gat Asp 255	Ile.	768
agc Ser	GJ y	ctt Leu	cct Pro 260	Leu	gtc Val	ggc Gly	gct Ala	gat Asp 265	His	ctt Leu	gtt Val	gat Asp	gcg Ala 270	Thr	caa Gln	816
aat Asr	aca Thr	gat Asp 275	Ala	tat Tyr	aca Thr	gag Glu	gta Val 280	Ser	gct Ala	tca Ser	tta Leu	aaa Lys 285	. Val	tgo Cys	atg Met	864
atç Met	aac Asn 290	Met	tcg Ser	j aag Lys	ato Ile	gca Ala 295	Asr	gad Asp	cto Lev	cgc Arg	tta Lei 300	Met	Ala Geo	tco Sei	gga Gly	912
pro 30	Arg	gco JAla	gga Gly	a ctt / Leu	gcq Ala 310	Gli	att i Ile	tct Sea	cto Lev	Pro 315	Ala	a cgt a Arg	caq g Gl	g cco	g ggt o Gly 320	960
tc: Se:	a tct c Sei	att	ato Met	g ccg t Pro 325	G1	g aaa y Lys	gto Val	c aat L Ası	Pro 330	Va.	ato L Med	g gcq t Ala	g gaq a Glu	g cte 1 Lei 33	g atc i Ile 5	1008
aa Asi	c caa n Glr	a att	gc e Ala 340	a Phe	c cae	g gtt n Val	t ato	gg: e -G1; 34;	y Ası	ga n Ası	c aat	t aca	a ate r Ile 350	е Су	c ctt s Leu	1056
gc Al	t tca a Sei	a gaa	a gco u Ala	c ggo a Gly	c ca y Gl	g cti n Lei	t gad ı Gl	g tte	g aa u As	c gto n Val	c at	g ga t Gl	g cc	c gt o Va	g ctt l Leu	1104

- 69 -

		355					360					365					
gtc Val	ttt Phe 370	Asn	ttg Leu	ctt Leu	caa Gln	tcc Ser 375	Ile	agc Ser	atc Ile	atg Met	aac Asn 380	Asn	ggc	ttc Phe	cgt Arg	1152	
tcg Ser 385	ttc Phe	act Thr	gac Asp	aac Asn	tgc Cys 390	tta Leu	aaa Lys	ggc	att Ile	gaa Glu 395	Ala	aac Asn	gaa Glu	aag Lys	cgt Arg 400	1200	٠
atg Met	aag Lys	caa Gln	tac Tyr	gta Val 405	Glu	aaa Lys	agc Ser	gca Ala	ggc Gly 410	gtg Val	atc Ile	aca Thr	gct Ala	gtc Val 415	aat Asn	1248	•
ccg Pro	cat His	ctt Leu	ggg Gly 420	tat Tyr	gaa Glu	gcg Ala	gca Ala	gct Ala 425	aga Arg	att Ile	gcc Ala	agg Arg	gaa Glu 430	gca Ala	att Ile	1'296	
atg Met	aca Thr	ggg Gly 435	caa Gln	tct Ser	gtc Val	cgg Arg	gat Asp 440	ctt Leu	tgt Cys	ctg Leu	cag Gln	cat His 445	gat Asp	gtg Val	ctg Leu	1344	
act Thr	gaa Glu 450	gaa Glu	gaa Glu	ttg Leu	gat Asp	att Ile 455	att Ile	tta Leu	aac Asn	cca Pro	tat Tyr 460	gag Glu	atg Met	acc Thr	aaa Lys	1392	
cca Pro 465	ggt Gly	atc Ile	gca Ala	ggg Gly	aaa Lys 470	gaa Glu	cta Leu	tta Leu	gaa Glu	aaa Lys 475	taa	,			•	1428	
<212 <212	0> 6 l> 4 2> Pl 3> Ba	75	Lus s	subt:	ilis											ı	
)> 6° Leu		Gly	Gln 5	Lys	Glu	Tyr	Arg	Val 10	Glu	Lys	Asp	Phe	Leu 15	Gly	1	
Glu	Lys	Gln	Ile 20	Glu	Ala	Asp	Val	Tyr 25	Tyr	Gly	Ile	Gln	Thr 30	Leu	Arg		
Ala	Ser	Glu 35	Asn	Phe	Pro	Ile	Thr 40	Gly	Tyr	Lys	Ile	His 45	Glu	Glu	Met		
Ile	Asn 50	Ala	Leu	Ala	Ile	Val 55	Lys	Lys	Ala	Ala	Ala 60	Leu	Ala	Asn	Met		
Asp 65	Val	Lys	Arg	Leu	Tyr 70	Glu	Gly	Ile	Gly	Gln 75	Ala	Ile	Val	Gln	Ala 80		
Ala	Asp	Glu	Ile	Leu 85	Glu	Gly	Lys	Trp	His 90	Asp	Gln	Phe	Ile	Val 95	Asp		
Pro	Ile	Gln	Gly 100	Gly	Ala	Gly	Thr	Ser	Met	Asn	Met	Asn	Ala	Asn	Glu		

- Val Ile Gly Asn Arg Ala Leu Glu Ile Met Gly His Lys Lys Gly Asp 115 120 125
- Tyr Ile His Leu Ser Pro Asn Thr His Val Asn Met Ser Gln Ser Gln 130 135 140
- Asn Asp Val Phe Pro Thr Ala Ile His Ile Ser Thr Leu Lys Leu Leu 145 150 155 160
- Glu Lys Leu Leu Lys Thr Met Glu Asp Met His Ser Val Phe Lys Gln
 165 170 175
- Lys Ala Gln Glu Phe His Ser Val Ile Lys Met Gly Arg Thr His Leu 180 185 190
- Gln Asp Ala Val Pro Ile Arg Leu Gly Gln Glu Phe Glu Ala Tyr Ser 195 200 205
- Arg Val Leu Glu Arg Asp Ile Lys Arg Ile Lys Gln Ser Arg Gln His 210 215 220
- Leu Tyr Glu Val Asn Met Gly Ala Thr Ala Val Gly Thr Gly Leu Asn 225 230 235 240
- Ala Asp Pro Glu Tyr Ile Lys Gln Val Val Lys His Leu Ala Asp Ile 245 250 255
- Ser Gly Leu Pro Leu Val Gly Ala Asp His Leu Val Asp Ala Thr Gln 260 265 270
- Asn Thr Asp Ala Tyr Thr Glu Val Ser Ala Ser Leu Lys Val Cys Met 275 280 285
- Met Asn Met Ser Lys Ile Ala Asn Asp Leu Arg Leu Met Ala Ser Gly 290 295 300
- Pro Arg Ala Gly Leu Ala Glu Ile Ser Leu Pro Ala Arg Gln Pro Gly 305 310 315 320
- Ser Ser Ile Met Pro Gly Lys Val Asn Pro Val Met Ala Glu Leu Ile 325 330 335
- Asn Gln Ile Ala Phe Gln Val Ile Gly Asn Asp Asn Thr Ile Cys Leu 340 345 350
- Ala Ser Glu Ala Gly Gln Leu Glu Leu Asn Val Met Glu Pro Val Leu 355 360 . 365
- Val Phe Asn Leu Leu Gln Ser Ile Ser Ile Met Asn Asn Gly Phe Arg 370 375 380
- Ser Phe Thr Asp Asn Cys Leu Lys Gly Ile Glu Ala Asn Glu Lys Arg 385 390 395 400
- Met Lys Gln Tyr Val Glu Lys Ser Ala Gly Val Ile Thr Ala Val Asn 405 410 415
- Pro His Leu Gly Tyr Glu Ala Ala Ala Arg Ile Ala Arg Glu Ala Ile

. - 71 -

			420)				425					430			
Met	Thr	Gly 435	Gln	Ser	Val	Arg	Asp 440	Leu	Cys	Leu	Gin	His 445	Asp	Val	Leu	
Thr	Glu 450	Glu	Glu	Leu	Asp	Ile 455	Ile	Leu	Asn	Pro	Tyr 460		Met	Thr	Lys	
Pro 465	Gly	Ile	Ala	Gly	Lys 470		Leu	Leu	Glu	Lys 475			1			
<21 <21	0> 6 1> 7 2> D 3> B	68 NA	lus	subt	ilis											
	1> C		(765)												
	0> 6											ı				
atg Met 1	aaa Lys	cga Arg	gaa Glu	agc Ser 5	aac Asn	att Ile	caa Gln	gtg Val	ctc Leu 10	agc Ser	cgt Arg	ggt Gly	caa Gln	aaa Lys 15	gat Asp	48
cag Gln	cct Pro	gtg Val	agc Ser 20	cag Gln	att Ile	tat Tyr	caa Gl'n	gta Val 25	tca Ser	aca Thr	atg Met	act Thr	tct Ser 30	cta Leu	tta Leu	96
gac Asp	gga Gly	gta Val 35	tat Tyr	gac Asp	gga Gly	gat Asp	ttt Phe 40	gaa Glu	ctg Leu	tca Ser	gag Glu	att Ile 45	ccg Pro	aaa Lys	tat Tyr	144
gga Gly	gac Asp 50	ttc Phe	ggt Gly	atc Ile	gga Gly	acc Thr 55	ttt Phe	aac Asn	aag Lys	ctt Leu	gac Asp 60	gga Gly	gag Glu	ctg Leu	att Ile	192
ggg Gly 65	ttt Phe	gac Asp	ggc Gly	gaa Glu	ttt Phe 70	tac Tyr	cgt Arg	ctt Leu	cgc Arg	tca Ser 75	gac Asp	gga Glų	acc Thr	gcg Ala	aca Thr 80	240
ccg Pro	gtc Val	caa Gln	aat Asn	gga Gly 85	gac Asp	cgt Arg	tca Ser	ccg Pro	ttc Phe 90	tgt Cys	tca Ser	ttt Phe	acg Thr	ttc Phe 95	ttt Phe	288
aca Thr	ccg Pro	gac Asp	atg Met 100	acg Thr	cac His	aaa Lys	att Ile	gat Asp 105	gcg Ala	aaa Lys	atg Met	aca Thr	cgc Arg 110	gaa Glu	gac Asp	336
ttt Phe	gaa Glu	aaa Lys 115	gag Glu	atc Ile	aac Asn	agc Ser	atg Met 120	ctg Leu	cca Pro	agc Ser	aga Arg	aac Asn 125	tta Leu	ttt Phe	tat Tyr	384
gċa Ala	att Ile 130	cgc Arg	att Ile	gac Asp	gga Gly	ttg Leu 135	ttt Phe	aaa Lys	aag Lys	gtg Val	cag Gln	aca Thr	aga Arg	aca Thr	gta Val	432

				aaa Lys												480
cag Gln	ccg Pro	att Ile	ttc Phe	aac Asn 165	ttc Phe	gac Asp	aac Aşn	gtg Val	aga Arg 170	gga Gly	acg Thr	att Ile	gta Val	ggt Gly 175	ttc Phe	528
				tat Tyr												576
				gaa Glu												624
				tgc Cys												672
				aac Asn												720
				aaa Lys 245											taa	768
<21 <21	0> 6 1> 2 2> P 3> B	55 . RT	lus	subt.	ilis											
<21 <21 <21 <40	1> 2 2> P 3> B 0> 6 Lys	55 RT acil		subt. Ser 5		Ile	Gln	Val	Leu 10		Arg	Gly	Gln	Lys 15		
<21 <21 <21 <40 Met	1> 2 2> P 3> B 0> 6 Lys	55 RT acil 9 Arg	Glu	Ser 5 Gln	Asn				10 Ser					15 Leu		
<21 <21 <21 <40 Met 1 Gln	1> 2 2> P 3> B 0> 6 Lys	55 RT acil 9 Arg Val	Glu Ser 20	Ser 5 Gln	Asn Ile	Tyr	Gln	Val 25 Glu	10 Ser	Thr	Met	Thr	Ser 30	15 Leu	Leu	
<21 <21 <40 Met 1 Gln	1> 2 2> P: 3> B 0> 6 Lys Pro	55 RT acil 9 Arg Val Val 35	Glu Ser 20	Ser 5 Gln Asp	Asn Ile Gly	Tyr Asp	Gln Phe 40	Val 25 Glu	10 Ser Leu	Thr	Met Glu	Thr Ile 45	Ser 30	Leu Lys	Leu	
<21 <21 <21 <40 Met 1 Gln Asp	1> 2 2> P 3> B 0> 6 Lys Pro Gly Asp 50	S5 RT acil 9 Arg Val Val 35	Glu Ser 20 Tyr	Ser 5 Gln Asp	Asn Ile Gly Gly	Tyr Asp Thr 55	Gln Phe 40	Val 25 Glu Asn	10 Ser Leu Lys	Thr Ser Leu	Met Glu Asp 60	Thr Ile 45 Gly	Ser 30 Pro	15 Leu Lys	Leu Tyr	
<21 <21 <21 <40 Met 1 Gln Asp Gly 65	1> 2 2> P 3> B 0> 6 Lys Pro Gly Asp 50	55 RT acil 9 Arg Val Val 35 Phe	Ser 20 Tyr Gly	Ser 5 Gln Asp Ile	Asn Ile Gly Gly Phe 70 Asp	Tyr Asp Thr 55	Gln Phe 40 Phe	Val 25 Glu Asn	Ser Leu Lys Arg	Thr Ser Leu Ser 75	Met Glu Asp 60	Thr Ile 45 Gly Gly	Ser 30 Pro Glu	Leu Lys Leu	Leu Tyr Ile Thr 80	
<21 <21 <21 <40 Met 1 Gln Asp Gly 65	1> 2 2> P 3> B 0> 6 Lys Pro Gly Asp 50 Phe	55 RT acil 9 Arg Val Val 35 Phe Asp	Ser 20 Tyr Gly Gly	Ser 5 Gln Asp Ile Glu Gly 85	Asn Ile Gly Gly Phe 70 Asp	Tyr Asp Thr 55 Tyr	Gln Phe 40 Phe Arg	Val 25 Glu Asn Leu	Lys Arg	Thr Ser Leu Ser 75	Met Glu Asp 60 Asp	Thr Ile 45 Gly Gly	Ser 30 Pro Glu Thr	Leu Lys Leu Ala Phe	Leu Tyr Ile Thr 80	

.,	130	urd	m	Asp	grà	135	Pne	rys	ьуѕ	Val	Gin 140	Thr	Arg	Thr	Val	
Glu 145	Leu	Gln	Glu	Lys	Pro 150	Tyr	Val	Pro	Met	Val 155	Glu	Ala	Val	Lys	Thr 160	
Gln	Pro	Ile	Phe	Asn 165	Phe	Asp	Asn	Val	Arg 170	Gly	Thr	Ile	Val	Gly 175	Phe	,
Leu	Thr	Pro	Ala 180	Tyr	Ala	Asn	Gly	Ile 185	Ala	Val	Ser	Gly	Tyr 190	His	Leu	•
His	Phe	Ile 195	Asp	Glu	Gly	Arg	Asn 200	Ser	Gly	Gly	His	Val 205	Phe	Asp	Tyr	,
Val	Leu 210	Glu	Asp	Суз	Thr	Val 215	Thr	Ile	Ser	Gln	Lys 220	Met	Asn	Met	Asn	
Leu 225	Arg	Leu	Pro	Asn _.	Thr 230	Ala	Asp	Phe	Phe	Asn 235	Ala	Asn	Leu	Asp	Asn 240	
Pro	Asp	Phe	Ala	Lys 245	Asp	Ile	Glu	Thr	Thr 250	Glu	Gly	Ser	Pro	Glu 255		
<211 <212)> 7(l> 12 2> Di	254 NA					. (
\Z1 3)	scuei	cichi	ia co	oli											
<220 <221)'> L> CI	os	(125)		oli											•
<220 <221 <222)'> L> CI	os 1)			oli											•
<220 <221 <222 <400 atg)> L> CI !> (:)> 7(aca	os 1)	(125) tcc	l) ctt	ttt	ggt Gly	gac Asp	aaa Lys	ttt Phe 10	acc Thr	cgc Arg	cac His	tcc Ser	ggc Gly 15	att Ile	48
<220 <221 <222 <400 atg Met 1)> l> CI 2> (:)> 7(aca Thr	OS 1)) ttc Phe ttg	tcc Ser	ctt Leu 5 gaa	ttt Phe	Gly	Asp	Lys gac	Phe 10	Thr	Arg	cac His acg Thr	Ser	Gly 15	Ile	48 96
<220 <221 <222 <400 atg Met 1 acg Thr)> L> CI 2> (I)> 7(aca Thr ctg Leu	OS 1)) ttc Phe ttg Leu	tcc Ser atg Met 20	ctt Leu 5 gaa Glu	ttt Phe gat Asp	Gly ctg Leu aat	Asp aac Asn ccg	gac Asp 25	Phe 10 ggt Gly cag	Thr tta Leu atc	Arg cgc Arg	His	cct Pro 30	Gly 15 ggc Gly	gcg Ala	
<220 <221 <222 <400 atg Met 1 acg Thr att Ile)> CI > CI > 70 aca Thr ctg Leu atg Met	os l) ttc Phe ttg Leu ctc Leu 35	tcc Ser atg Met 20 ggc Gly	ctt Leu 5 gaa Glu ggc Gly	ttt Phe gat Asp ggt Gly	ctg Leu aat Asn	aac Asn ccg Pro 40	gac Asp 25 gcg Ala	Phe 10 ggt Gly cag Gln	Thr tta Leu atc Ile	cgc Arg ccg Pro	acg Thr gaa Glu	cct Pro 30 atg Met	Gly 15 ggc Gly cag Gln	gcg Ala gac Asp	96
<220 <221 <222 <400 atg Met 1 acg Thr att Ile tac Tyr)> CI P> (I P> (I aca Thr ctg Leu atg Met ttc Phe 50 gca	os l) ttc Phe ttg Leu ctc Leu 35 cag Gln	tcc Ser atg Met 20 ggc Gly acg Thr	ctt Leu 5 gaa Glu ggc Gly cta Leu	ttt Phe gat Asp ggt Gly ctg Leu	Ctg Leu aat Asn acc Thr 55	aac Asn ccg Pro 40 gac Asp	gac Asp 25 gcg Ala atg Met	Phe 10 ggt Gly cag Gln ctg Leu cag	Thr tta Leu atc Ile gaa Glu	cgc Arg ccg Pro	His acg Thr gaa Glu 45	cct Pro 30 atg Met aaa Lys	Gly 15 ggc Gly cag Gln gcg Ala	gcg Ala gac Asp	96 144

	85 '	90	.95
cca cag aat att o Pro Gln Asn Ile i 100	gca cta aca Ala Leu Thr	aac ggc agc cag Asn Gly Ser Gln 105	agc gcg ttt ttc tac 336 Ser Ala Phe Phe Tyr , 110
tta ttt aac ctg Leu Phe Asn Leu 115	ttt gcc gga Phe Ala Gly	cgc cgt gcc gat Arg Arg Ala Asp 120	ggt cgg gtc aaa aaa 384 Gly Arg Val Lys Lys 125
gtg ctg ttc ccg Val Leu Phe Pro 130	ctt gca ccg Leu Ala Pro 135	gaa tac att ggc Glu Tyr Ile Gly	tat gct gac gcc gga 432 Tyr Ala Asp Ala Gly 140
ctg gaa gaa gat Leu Glu Glu Asp 145	ctg ttt gtc Leu Phe Val 150	tct gcg cgt ccg Ser Ala Arg Pro 155	aat att gaa ctg ctg 480 Asn Ile Glu Leu Leu 160
ccg gaa ggc cag Pro Glu Gly Gln	ttt aaa tac Phe Lys Tyr 165	cac gtc gat ttt His Val Asp Phe 170	gag cat ctg cat att 528 Glu His Leu His Ile 175
ggc gaa gaa acc Gly Glu Glu Thr 180	ggg atg att Gly Met Ile	tgc gtc tcc cgg Cys Val Ser Arg 185	g ccg acg aat cca aca 576 g Pro Thr Asn Pro Thr 190
ggc aat gtg att Gly Asn Val Ile 195	act gac gaa Thr Asp Glu	gag ttg ctg aac Glu Leu Leu Lys 200	g ctt gac gcg ctg ggc 624 s Leu Asp Ala Leu Gly 205
aat caa cac ggc Asn Gln His Gly 210	att ccg ctg Ile Pro Leu 215	gtg att gat aad Val Ile Asp Asr	c gct tat ggc gtc ccg 672 n Ala Tyr Gly Val Pro 220
ttc ccg ggt atc Phe Pro Gly Ile 225	atc ttc agt Ile Phe Ser 230	gaa gcg cgc ccc Glu Ala Arg Pro 23	g cta tgg aat ccg aat 720 o Leu Trp Asn Pro Asn 5 240
atc gtg ctg tgc Ile Val Leu Cys	atg agt ctt Met Ser Leu 245	tcc aag ctg gg Ser Lys Leu Gl 250	t cta cct ggc tcc cgc 768 y Leu Pro Gly Ser Arg 255
tgc ggc att atc Cys Gly Ile Ile 260	Ile Ala Asn	gaa aaa atc at Glu Lys Ile Il 265	c acc gcc atc acc aat 816 e Thr Ala Ile Thr Asn 270
atg aac ggc att Met Asn Gly Ile 275	atc agc ctg lle Ser Leu	gca ect ggc gg Ala Pro Gly Gl 280	t att ggt ccg gcg atg 864 y Ile Gly Pro Ala Met 285
atg tgt gaa atg Met Cys Glu Met 290	att aag cgt : Ile Lys Arg 295	Asn Asp Leu Le	eg ege etg tet gaa aca 912 eu Arg Leu Ser Glu Thr 300
gtc atc aaa ccg Val Ile Lys Pro 305	ttt tac tac Phe Tyr Tyr 310	c cag cgt gtt ca c Gln Arg Val Gl 31	ng gaa act atc gcc atc 960 In Glu Thr Ile Ala Ile 15 320

Ile	cgc Arg	cgc	tat Tyr	tta Leu 325	Pro	gaa Glu	aat Asn	cgc Arg	tgc Cys 330	Leu	att Ile	cat His	aaa Lys	Pro	gaa Glu	1008
gga Gly	gcc Ala	att Ile	ttc Phe 340	ctc Leu	tgg Trp	cta Leu	tgg Trp	ttt Phe 345	aag Lys	gat Asp	ttg Leu	ecc Pro	att Ile 350	Thr	acc Thr	1056
aag Lys	cag Gln	ctc Leu 355	Tyr	cag Gln	cgc Arg	ctg Leu	aaa Lys 360	Ala	cgc Arg	ggc Gly	gtg Val	ctg Leu 365	atg Met	gtg Val	ccg	1104
ggg Gly	cac His 370	ASI	ttc Phe	ttc Phe	cca Pro	ggg Gly 375	ctg Leu	gat Asp	aaa Lys	ccg Pro	tgg Trp 380	ccg Pro	cat His	acg Thr	cat	1152
caa Gln 385	Cys	atg Met	cgc Arg	atg Met	aac Asn 390	tac Tyr	gta Val	cca Pro	gag Glu	ccg Pro 395	gag Glu	aaa Lys	att Ile	gag Glu	gcg Ala 400	1200
GJA	gtg Val	aag Lys	att Ile	ctg Leu 405	gcg Ala	gaa Glu	gag Glu	ata Ile	gaa Glu 410	aga Arg	gcc Ala	tgg Trp	gct Ala	gaa Glu 415	agt Ser	1248
cac His	taa						1					ı				1254
<21	0> 7	1														•
<21 <21	1> 4 2> P	17 RT	richi	ia co	oli											
<21 <21 <21 <40	1> 4 2> P1 3> E3	17 RT schei	richi Ser			Gly	Asp	Lys	Phe 10	Thr	Arg	His	Ser	Gly 15	Ile	
<21 <21 <21 <40 Met	1> 4 2> P 3> E 0> 7 Thr	17 RT schei l Phe		Leu 5	Phe				10					15		
<21 <21 <21 <40 Met 1	1> 4. 2> P 3> E 0> 7 Thr	17 RT scher l Phe Leu	Ser Met	Leu 5 Glu	Phe Asp	Leu	Asn	Asp 25	10 Gly	Leu	Arg	Thr	Pro 30	15 Gly	Ala	
<21 <21 <40 Met 1 Thr	1> 4. 2> P. 3> E. 0> 7. Thr Leu	17 RT scher 1 Phe Leu Leu 35	Ser Met 20	Leu 5 Glu Gly	Phe Asp Gly	Leu Asn	Asn Pro 40	Asp 25 Ala	Gly Gln	Leu Ile	Arg Pro	Thr Glu 45	Pro 30 Met	15 Gly Gln	Ala Asp	
<21 <21 <21 <40 Met 1 Thr Ile	1> 4. 2> P! 3> E: 0> 7: Thr Leu Met	17 RT schen l Phe Leu 35 Gln	Ser Met 20 Gly	Leu 5 Glu Gly Leu	Phe Asp Gly Leu	Leu Asn Thr 55	Asn Pro 40 Asp	Asp 25 Ala Met	Gly Gln Leu	Leu Ile Glu	Arg Pro Ser 60	Thr Glu 45 Gly	Pro 30 Met	15 Gly Gln Ala	Ala Asp Thr	
<21 <21 <40 Met 1 Thr Ile Tyr Asp 65	1> 4 2> P 3> E 0> 7: Thr Leu Met Phe 50	17 RT schei 1 Phe Leu 35 Gln	Ser Met 20 Gly	Leu 5 Glu Gly Leu Asn	Phe Asp Gly Leu Tyr 70	Leu Asn Thr 55 Asp	Asn Pro 40 Asp Gly	Asp 25 Ala Met	Gly Gln Leu Gln	Leu Ile Glu Gly 75	Arg Pro Ser 60 Lys	Thr Glu 45 Gly Thr	Pro 30 Met Lys Glu	Gly Gln Ala Leu	Ala Asp Thr Leu 80	
<21 <21 <40 Met 1 Thr Ile Tyr Asp 65 Thr	1> 4 2> P1 3> E3 0> 7: Thr Leu Met Phe 50 Ala	17 RT schei l Phe Leu 35 Gln Leu Leu	Ser Met 20 Gly Thr	Leu 5 Glu Gly Leu Asn Gly 85	Phe Asp Gly Leu Tyr 70 Met	Leu Asn Thr 55 Asp	Asn Pro 40 Asp Gly Arg	Asp 25 Ala Met Pro	Gly Gln Leu Gln Lys 90	Leu Ile Glu Gly 75 Leu	Arg Pro Ser 60 Lys	Thr Glu 45 Gly Thr Trp	Pro 30 Met Lys Glu Asp	Gly Gln Ala Leu Ile 95	Ala Asp Thr Leu 80 Glu	

- Val Leu Phe Pro Leu Ala Pro Glu Tyr Ile Gly Tyr Ala Asp Ala Gly 130 135 140
- Leu Glu Glu Asp Leu Phe Val Ser Ala Arg Pro Asn Ile Glu Leu Leu 145 150 155 160
- Pro Glu Gly Gln Phe Lys Tyr His Val Asp Phe Glu His Leu His Ile 165 170 175
- Gly Glu Glu Thr Gly Met Ile Cys Val Ser Arg Pro Thr Asn Pro Thr 180 185 190
- Gly Asn Val Ile Thr Asp Glu Glu Leu Leu Lys Leu Asp Ala Leu Gly
 195 200 205
- Asn Gln His Gly Ile Pro Leu Val Ile Asp Asn Ala Tyr Gly Val Pro
- Phe Pro Gly Ile Ile Phe Ser Glu Ala Arg Pro Leu Trp Asn Pro Asn 225 230 235 240
- Ile Val Leu Cys Met Ser Leu Ser Lys Leu Gly Leu Pro Gly Ser Arg
- Cys Gly Ile Ile Ile Ala Asn Glu Lys Ile Ile Thr Ala Ile Thr Asn 260 265 270
- Met Asn Gly Ile Ile Ser Leu Ala Pro Gly Gly Ile Gly Pro Ala Met 275 280 285
- Met Cys Glu Met Ile Lys Arg Asn Asp Leu Leu Arg Leu Ser Glu Thr 290 295 300
- Val Ile Lys Pro Phe Tyr Tyr Gln Arg Val Gln Glu Thr Ile Ala Ile 305 310 315 320
- Ile Arg Arg Tyr Leu Pro Glu Asn Arg Cys Leu Ile His Lys Pro Glu 325 330 335
- Gly Ala Ile Phe Leu Trp Leu Trp Phe Lys Asp Leu Pro Ile Thr Thr 340 345 350
- Lys Gln Leu Tyr Gln Arg Leu Lys Ala Arg Gly Val Leu Met Val Pro 355 360 365
- Gly His Asn Phe Phe Pro Gly Leu Asp Lys Pro Trp Pro His Thr His 370 380
- Gln Cys Met Arg Met Asn Tyr Val Pro Glu Pro Glu Lys Ile Glu Ala 385 390 395 400
- Gly Val Lys Ile Leu Ala Glu Glu Ile Glu Arg Ala Trp Ala Glu Ser 405 410 415

His

<210> 72 <211> 8803 <212> DNA <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
pAN294 plasmid

<400> 72 tgcgccgcta cagggcgcgt ccattcgcca ttcaggctgc gcaactgttg ggaagggcga 60 tcggtgcggg cctcttcgct attacgccag ctggcgaaag ggggatgtgc tgcaaggcga 120 ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagtgaa 180 ttgtaatacg actcactata gggcgaattg ggcccgacgt cgcatgctgg atgaaaagcc 240 gatgaccgct tttcaggtct gtcagcagct ttttcctgct gtatatgaaa aggaattgtt 300 tttaacgatg tcagaaacgg caggtcacct tgatgtgttg gaggctgaag aagccatcac 360 gtcatattgg gaaggaaata ccgtatactt taaaacaatg aagaggtgaa atgggtgaaa 420 catatagcgg gaaaaaggat ttggataacc ggcgcttcag gagggcttgg agaaagaatc 480 gcatacttat gcgcggctga aggagcccat gtcctgctgt cggctagacg cgaggatcgt 540 ttgatagaaa tcaaaaggaa aataaccgag gaatggagcg gacagtgtga gattttcct 600 ctggatgtcg gccgcctaga ggatatcgcc cgggtccgcg atcagatcgg ctcgattgat 660 gtactgatta acaatgcagg cttcggtata tttgaaacgg ttttagactc tacattggat 720 gacatgaaag cgatgtttga tgtgaatgtc ttcggcctga tcgcctgtac aaaagcggtg 780 cttccgcaaa tgcttgagca aaaaaaggga catatcatca atatcgcctc tcaagcgggg 840 aaaatcgcca caccgaagtc tagcctgtat tccgcgacca aacatgccgt gttaggttac 900 tcaaacgett tgcggatgga getttcggga accggcattt atgtgacaac agtcaacccg 960 ggcccgattc agacggactt tttttccatt gctgataaag gcggggacta cgccaaaaat 1020 gtcggccgct ggatgcttga tcctgatgac gtggcagctc aaattacagc tgcaattttt 1080 acgaaaaagc gggagatcaa tcttccgcgt ttaatgaatg ccggcactaa gctgtatcag 1140 ctgtttccag ctcttgtaga aaagctggca ggacgcgcgc tcatgaaaaa ataatgatag 1200 aactgcctgt ggtggagtgg cttgtttctc acggggcagt ttttgatagt ggaagggaga 1260 gattgttgaa tgtcagttca ttcagaagtc cttcatgctc tgcttaaaga tccgtttatt 1320 cagaaactga ttgatgcaga gcctgtattc tgggcaaatt caggcaagaa agaggggcca 1380 ttaccccgtg cagatgagtg ggcaaccgag atagcggaag cggaaaaaag aatgcagcgg 1440

tttgcacctt acattgccga ggtgtttcct gagacgaaag gcgctaaagg aatcatcgag 1500 tctccgcttt ttgaggtgca gcatatgaag ggaaagctgg aagcggcata tcagcagcca 1560 tttcccggaa gatggctttt aaagtgcgac catgagcttc cgatttcagg atcgattaaa 1620 gcgaggggcg ggatttatga agtgttaaag tatgctgaaa atctcgcgct tcaagaagga 1680 atgetteagg aaacegatga ttacegeate ttacaggaag ageggtttae egggtttte 1740 tocogotatt cgattgctgt cggttcgaca ggaaatctag gtttaagcat cggcatcatc 1800 ggcgcggcac tcgggtttcg cgtgacagtg catatgtccg ccgatgctaa gcagtggaaa 1860 aaggatetee teegeeaaaa gggagteaet gttatggagt acgaaacaga ttacagtgaa 1920 gcggtgaacg aagggagacg gcaggcggaa caagatccat tctgttattt tattgatgat 1980 gaacattete gteagetgtt ettaggatat getgttgetg caageegatt aaaaacaeag 2040 cttgactgta tgaatataaa gccaagtctt gagacgccct tgtttgtgta tctgccgtgc 2100 ggagtcggcg gaggaccggg cggtgtagca tttgggctga agcttttata cggagatgat 2160 gttcatgtgt ttttcgcaga accaactcat tcaccttgta £gctgttagg gctttattca 2220 ggacttcacg agaagatctc cgtccaggat atcggcctgg ataatcagac ggctgctgac 2280 ggacttgccg tagggaggcc gtcaggattt gtcggcaagc tgattgaacc gcttctgagc 2340 ggctgttata cggtagagga caatacgctt tatactttgc ttcatatgct ggctgtatct 2400 gaagataaat atttagagee etetgetett getggeatgt tegggeeggt teagettttt 2460 togacagaag agggaaggog ctatgotoag aaatataaga tggaacatgo ogtacatgto 2520 gtctggggaa cgggaggaag catggttcca aaagatgaaa tggctgcgta taaccgaatc 2580 ggtgctgatt tgctaaaaaa acgaaatgga aaataagcag acagtgaaaa ggttttccgt 2640 tacaatettt gtaagggttt taacetacag agagtcaggt gtaaacagtg aaaaataaag 2700 aacttaacct acatacttta tatacacagc acaatcggga gtcttggtct ggttttgggg 2760 ggcatttgtc gattgctgta tctgaagaag aggcaaaagc tgtggaagga ttgaatgatt 2820 atctatctgt tgaagaagtg gagacgatct atattccgct tgttcgcttg cttcatttac 2880 atgtcaagtc tgcggctgaa cgcaataagc atgtcaatgt ttttttgaag cacccacatt 2940 cagecaaaat teegtttatt ateggeattg eeggeagtgt egeagtegga aaaageaega 3000 cggcgcggat cttgcagaag ctgctttcgc gtttgcctga ccgtccaaaa gtgagcctta 3060 tcacgacaga tggtttttta tttcctactg ccgagctgaa aaagaaaaat atgatgtcaa 3120 gaaaaggatt tootgaaago tatgatgtaa aggogotgot ogaatttttg aatgaottaa 3180

aatcaggaaa ggacagcgta aaggccccgg tgtattccca tctaacctat gaccgcgagg 3240 aaggtgtgtt cgaggttgta gaacaggcgg atattgtgat tattgaaggc attaatgttc 3300 ttcagtcgcc caccttggag gatgaccggg aaaacccgcg tatttttgtt tccgatttct 3360 ttgattttc gatttatgtg gatgcggagg aaagccggat tttcacttgg tatttagagc 3420 gttttcgcct gcttcgggaa acagcttttc aaaatcctga ttcatatttt cataaattta 3480 aagacttgtc cgatcaggag gctgacgaga tggcagcctc gatttgggag agtgtcaacc 3540 ggccgaattt atatgaaaat attttgccaa ctaaattcag gtcagatctc attttgcgta 3600 agggagacgg gcataaggtc gaggaagtgt tggtaaggag ggtatgaaat gtgctgcagc 3660 tcgagcaata gttaccctta ttatcaagat aagaaagaaa aggatttttc gctacgctca 3720 aatcetttaa aaaaacacaa aagaccacat tttttaatgt ggtetttatt etteaactaa 3780 ttatgttaca gtaatattga cttttaaaaa aggattgatt ctaatgaaga aagcagacaa 3900 gtaagcctcc taaattcact ttagataaaa atttaggagg catatcaaat gaactttaat 3960 aaaattgatt tagacaattg gaagagaaaa gagatattta atcattattt gaaccaacaa 4020 acgactitta gtataaccac agaaattgat attagtgttt tataccgaaa cataaaacaa 4080 gaaggatata aattttaccc tgcatttatt ttcttagtga caagggtgat aaactcaaat 4140 acagctttta gaactggtta caatagcgac ggagagttag gttattggga taagttagag 4200 ccactttata caatttttga tggtgtatct aaaacattct ctggtatttg gactcctgta 4260 aagaatgact tcaaagagtt ttatgattta tacctttctg atgtagagaa atataatggt 4320 tcggggaaat tgtttcccaa aacacctata cctgaaaatg ctttttctct ttctattatt 4380 ccatggactt catttactgg gtttaactta aatatcaata ataatagtaa ttaccttcta 4440 cccattatta cagcaggaaa attcattaat aaaggtaatt caatatattt accgctatct 4500 ttacaggtac atcattctgt ttgtgatggt tatcatgcag gattgtttat gaactctatt 4560 caggaattgt cagataggcc taatgactgg cttttataat atgagataat gccgactgta 4620 ctttttacag teggttttct aatgteacta acctgeeceg ttagttgaag aaggttttta 4680 tattacaget gtcgactcgt gatcttcgga caggctgttc agctttttct caatgcgatc 4740 cagctgcgct tttcggtttt tcgcatactt gaagcctgta acagccgcaa agacgacagc 4800 ggcaaatata ataaatacaa acagetgaaa cateacatea eetatattea tgttetteae 4860 ctcatgtttg cgggagagat tcattctctt ccgtttttta tttaaagcgg cttttccaga 4920

cgggaacggt gttttgtggt ctccattttc atttgccgat aggcgaacgc taaaaatggc 4980 aggeegagea gggtaatgee geteaggaea gaaaaaatat aaateggeeg geeagegeea 5040 aacaggtota tacatatoco cocgacocaa gggoogatga cgtttocgag otgtggaaaa 5100 ccgattgccc cgaaataagt gccttttaat cctggttttg caatctggtc tacatacaaa 5160 tocatcatag agaataaaag cacttogoog attgtaaatg tgatgacaat catcacaatt 5220 gatggaacac cgtgtgatac ggtgaaaatg gccatgctga tgctaaccat cacattaccg 5280 agcatcagag aacaaagcgg cgaaaaccgt tttgcaaaat ggacaatggg aaattgcgtc 5340 qccaacacaa cgattgcgtt taatgtcagc atcagcccat acagcttcgt tccattgccg 5400 atcaaggggt totgogccat atactgaggg aatgtggaac tgaattgtga gtagccgaag 5460 gtgcatagcg taatgccgac caaagcaatg gtaaaaagat aatccttttg cgtgaccata \$520 , aacgetteee geaegeteat atttegggae tgggetggtg etgataagga tggatgtttt 5580 ttaaattgga gggcaagcac aattccgtat agtccgtaaa tgactgcagg caccaaaaag 5640 ggcgtagteg attgegatga geegaaatat aggeeaagea eaggteegaa gaeaaegeeg 5700 atattaatag cegeatageg taaattaaaa actageagte tegittitte tietgicata 5760 tcagacaaca aggcctttga agcgggctca aacagtgatt tgcaaagacc gtttaatgcg 5820 tttactacaa aaaacaccca gagattagat gctgccgcaa agcctgcaaa taccagcatc 5880 catccgaaaa tcgatacaag catcatgttt tttctgccga atttatctga gatatatccg 5940 ccgtaaaagc ttgcgaggat gccgactgat gagctcgcgg cgatgaccag ccctgcatag 6000 gaagctgatg cgccttggac ggctgtcaaa taaatcgcta aaaaaggaat gctcatcgat 6060 qttqccattc tgccgaaaat ggttccgatt ataattgtac gcgttggatg catagcttga 6120 qtattctata gtgtcaccta aatagcttgg cgtaatcatg gtcatagctg tttcctgtgt 6180 qaaattgtta teegeteaca atteeacaca acataegage eggaageata aagtgtaaag 6240 cctggggtgc ctaatgagtg agctaactca cattaattgc gttgcgctca ctgcccgctt 6300 tocaqtoggg aaacotgtog tgccagotgo attaatgaat oggocaacgo goggggagag 6360 geggtttgeg tattgggege tetteegett cetegeteae tgaetegetg egeteggteg 6420 ttcggctgcg gcgagcggta tcagctcact caaaggcggt aatacggtta tccacagaat 6480 caggggataa cgcaggaaag aacatgtgag caaaaaggcca gcaaaaggcc aggaaccgta 6540 aaaaggccgc gttgctggcg tttttcgata ggctccgccc ccctgacgag catcacaaaa 6600 atcgacgete aagteagagg tggegaaace egacaggaet ataaagatae eaggegttte 6660

cccctggaag ctccctcgtg cgctctcctg ttccgaccct gccgcttacc ggatacctgt 6720 cogectttet ceetteggga 'agegtggege ttteteatag etcaegetgt aggtatetea 6780 gttcggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaacccccc gttcagcccg 6840 accyctycyc cttatccygt aactatcytc ttyaytccaa cccyytaaga cacyacttat 6900 cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta ggcggtgcta 6960 cagagttett gaagtggtgg eetaactaeg getacaetag aaggacagta tttggtatet 7020 gcgctctgct gaagccagtt accttcggaa aaagagttgg tagctcttga tccggcaaac 7080 aaaccaccgc tggtagcggt ggtttttttg tttgcaagca gcagattacg cgcagaaaaa 7140 aaggatetea agaagateet ttgatetttt etaeggggte tgaegeteag tggaaegaaa 7200 actcacgtta agggattttg gtcatgagat tatcaaaaag gatcttcacc tagatccttt 7260 taaattaaaa atgaagtttt aaatcaatct aaagtatata tgagtaaact tggtctgaca 7320 gttaccaatg cttaatcagt gaggcaccta tctcagcgat ctgtctattt cgttcatcca 7380 tagttgcctg actccccgtc gtgtagataa ctacgatacg ggagggctta ccatctggcc 7440 ccagtgctgc aatgataccg cgagacccac gctcaccggc tccagattta tcagcaataa 7500 accagecage eggaagggee gagegeagaa gtggteetge aactttatee geeteeatee 7560 agtctattaa ttgttgccgg gaagctagag taagtagttc gccagttaat agtitgcgca 7620 acgttgttgg cattgctaca ggcatcgtgg tgtcacgctc gtcgtttggt atggcttcat 7680 tcagctccgg ttcccaacga tcaaggcgag ttacatgatc ccccatgttg tgcaaaaaag 7740 eggttagete etteggteet eegategttg teagaagtaa gttggeegea gtgttateae 7800 tcatggttat ggcagcactg cataattctc ttactgtcat gccatccgta agatgctttt 7860 ctgtgactgg tgagtactca accaagtcat tctgagaata ccgcgcccgg cgaccgagtt 7920 gctcttgccc ggcgtcaata cgggataata gtgtatgaca tagcagaact ttaaaagtgc 7980 tcatcattgg aaaacgttct tcggggcgaa aactctcaag gatcttaccg ctgttgagat 8040 ccagttcgat gtaacccact cgtgcaccca actgatcttc agcatctttt actttcacca 8100 gcgtttctgg gtgagcaaaa acaggaaggc aaaatgccgc aaaaaaggga ataagggcga 8160 cacggaaatg ttgaatactc atactcttcc tttttcaata ttattgaagc atttatcagg 8220 gttattgtct catgagcgga tacatatttg aatgtattta gaaaaataaa caaatagggg 8280 tteegegeac attteecega aaagtgeeac etgtatgegg tgtgaaatac egeacagatg 8340 cgtaaggaga aaataccgca tcaggcgaaa ttgtaaacgt taatattttg ttaaaattcg 8400

cgttaaatat ttgttaaatc agctcattt ttaaccaata ggccgaaatc ggcaaaatcc 8460 cttataaatc aaaagaatag accgagatag ggttgagtgt tgttccagtt tggaacaaga 8520 gtccactatt aaagaacgtg gactccaacg tcaaagggcg aaaaaccgtc tatcagggcg 8580 atggcccact acgtgaacca tcacccaaat caagttttt gcggtcgagg tgccgtaaag 8640 ctctaaatcg gaaccctaaa gggagccccc gatttagagc ttgacggga aagccggcga 8700 acgtggcgag aaaggaaggg aagaaagcga aaggagcggg cgctagggcg ctggcaagtg 8760 tagcggtcac gctgcggta accaccacc ccgccgcgct taa 8803

<210> 73

<211> 8320

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN296 plasmid

<400> 73 tgcgccgcta cagggcgcgt ccattcgcca ttcaggctgc gcaactgttg ggaagggcga 60 teggtgeggg cetetteget attacgecag etggegaaag ggggatgtge tgcaaggega 120 ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagtgaa 180 ttgtaatacg actcactata gggcgaattg ggcccgacgt cgcatgctgg atgaaaagcc 240 qatqaccqct tttcaggtct gtcagcagct ttttcctgct gtatatgaaa aggaattgtt 300 tttaacgatg tcagaaacgg caggtcacct tgatgtgttg gaggctgaag aagccatcac 360 qtcatattqq gaaqgaaata ccqtatactt taaaacaatq aagaggtgaa atgggtgaaa 420 catatagcgg gaaaaaggat ttggataacc ggcgcttcag gagggcttgg agaaagaatc 480 gcatacttat gcgcggctga aggagcccat gtcctgctgt cggctagacg cgaggatcgt 540 ttgatagaaa tcaaaaggaa aataaccgag gaatggagcg gacagtgtga gattttcct 600 ctggatgtcg gccgcctaga ggatatcgcc cgggtccgcg atcagatcgg ctcgattgat 660 gtactgatta acaatgcagg cttcggtata tttgaaacgg ttttagactc tacattggat 720 qacatqaaaq cqatqtttga tgtgaatgtc ttcggcctga tcgcctgtac aaaagcggtg 780 cttccgcaaa tgcttgagca aaaaaaggga catatcatca atatcgcctc tcaagcgggg 840 aaaatcgcca caccgaagtc tagcctgtat tccgcgacca aacatgccgt gttaggttac 900 tcaaacgctt tgcggatgga gctttcggga accggcattt atgtgacaac agtcaacccg 960 ggcccgattc agacggactt tttttccatt gctgataaag gcggggacta cgccaaaaat 1020 gtcggccgct ggatgcttga tcctgatgac gtggcagctc aaattacagc tgcaattttt 1080 acgaaaaagc gggagatcaa tcttccgcgt ttaatgaatg ccggcactaa .gctgtatcag 1140 ctgtttccag ctcttgtaga aaagctggca ggacgcgcgc tcatgaaaaa ataatgatag 1200 aactgcctgt ggtggagtgg cttgtttctc acggggcagt ttttgatagt ggaagggaga 1260 gattgttgaa tgtcagttca ttcagaagtc cttcatgctc tgcttaaaga tccgtttatt 1320 cagaaactga ttgatgcaga gcctgtattc tgggcaaatt caggcaagaa agaggggcca 1380 ttaccccgtg cagatgagtg ggcaaccgag atagcggaag cggaaaaaag aatgcagcgg 1440 tttgcacctt acattgccga ggtgtttcct gagacgaaag gcgctaaagg aatcatcgag 1500 tctccgcttt ttgaggtgca gcatatgaag ggaaagctgg aagcggcata tcagcagcca 1560 tttcccggaa gatggctttt aaagtgcgac catgagcttc cgatttcagg atcgattaaa 1620 gcgaggggcg ggatttatga agtgttaaag tatgctgaaa atctcgcgct tcaagaagga 1680 atgettcagg aaaccgatga ttaccgcatc ttacaggaag agcggtttac cgggttttc 1740 tcccgctatt cgattgctgt cggttcgaca ggaaatctag gtttaagcat cggcatcatc 1800 ggcycggcac tcgggtttcg cgtgacagtg catatgtccg ccgatgctaa gcagtggaaa 1860 aaggatetee teegeeaaaa gggagteaet gttatggagt acgaaacaga ttacagtyaa 1920 gcggtgaacg aagggagacg gcaggcggaa caagatccat tctgttattt tattqatgat 1980 gaacattctc gtcagctgtt cttaggatat gctgttgctg caagccgatt aaaaacacag 2040 cttgactgta tgaatataaa gccaagtctt gagacgccct tgtttgtgta tctgccgtgc 2100 ggagtcggcg gaggaccggg cggtgtagca tttgggctga agcttttata cggagatgat 2160 gttcatgtgt ttttcgcaga accaactcat tcaccttgta tgctgttagg gctttattca 2220 ggacttcacg agaagatctc cgtccaggat atcggcctgg ataatcagac ggctgctgac 2280 ggacttgccg tagggaggcc gtcaggattt gtcggcaagc tgattgaacc gcttctgagc 2340 ggctgttata cggtagagga caatacgctt tatactttgc ttcatatgct ggctgtatct 2400 gaagataaat atttagagcc ctctgctctt gctggcatgt tcgggccggt tcagcttttt 2460 tcgacagaag agggaaggcg ctatgctcag aaatataaga tggaacatgc cgtacatgtc 2520 gtctggggaa cgggaggaag catggttcca aaagatgaaa tggctgcgta taaccgaatc 2580 ggtgctgatt tgctaaaaaa acgaaatgga aaataagcag acagtgaaaa ggttttccgt 2640 tacaatcttt gtaagggttt taacctacag agagtcaggt gtaaacagtg aaaaataaag 2700 aacttaacct acatacttta tatacacagc acaatcggga gtcttctgca gctcgagcaa 2760

tagttaccct tattatcaag ataagaaaga aaaggatttt tcgctacgct caaatccttt 2820 aaaaaaacac aaaagaccac attttttaat gtggtcttta ttcttcaact aaagcaccca 2880 ttagttcaac aaacgaaaat tggataaagt gggatatttt taaaatatat atttatgtta 2940 cagtaatatt gacttttaaa aaaggattga ttctaatgaa gaaagcagac aagtaagcct 3000 cctaaattca ctttagataa aaatttagga ggcatatcaa atgaacttta ataaaattga 3060 tttagacaat tggaagagaa aagagatatt taatcattat ttgaaccaac aaacgacttt 3120 tagtataacc acagaaattg atattagtgt tttataccga aacataaaac aagaaggata 3180 taaattttac cctgcattta ttttcttagt gacaagggtg ataaactcaa atacagcttt 3240 tagaactggt tacaatagcg acggagagtt aggttattgg gataagttag agccacttta 3300 tacaattttt gatggtgtat ctaaaacatt ctctggtatt tggactcctg taaagaatga 3360 cttcaaagag ttttatgatt tatacctttc tgatgtagag aaatataatg gttcggggaa 3420 attgtttccc aaaacaccta tacctgaaaa tgcttttct ctttctatta ttccatggac 3480 ttcatttact gggtttaact taaatatcaa taataatagt aattactttc tacccattat 3540 tacagcagga aaattcatta ataaaggtaa ttcaatatat ttaccgctat ctttacaggt 3600 acatcattct gtttgtgatg gttatcatgc aggattgttt atgaactcta ttcaggaatt 3660 gtcagatagg cctaatgact ggcttttata atatgagata atgccgactg tactttttac 3720 agtoggtttt ctaatgtoac taacotgooc ogttagttga agaaggtttt tatattacag 3780 ctgtcgacta aggtcgagga agtgttggta aggagggtat gaaatgtgca tcatattgaa 3840 ctgtatgtct ctgatttgga ggcgtctagg cggttttggg gctggttctt aaaagaactt 3900 ggttataaag agtatcaaaa atggagctca ggcatcagct ggaagaaaga tcgtttttac 3960 ctagtgattg tgcaggcgaa agagccattt ctagagccgg aataccatag atgccgagtc 4020 ggtctgaacc atctcgcatt tcatgctgaa tccaagcttc aagtcgatca gatgactgaa 4080 aaattgacgg caaaaggcta tcgtgtgttg taccgagaca ggcatccttt tgccggagga 4140 qacqqqcatt atqcaqtctt ttqtqaqqat ccaqaccqqa ttaaqqtaqa qctcqttqcc 4200 ccaagetgtt aategtgate tteggacagg etgtteaget tttteteaat gegateeage 4260 tgcqcttttc ggtttttcgc atacttgaag cctgtaacag ccgcaaagac gacagcggca 4320 aatataataa atacaaacag ctgaaacatc acatcaccta tattcatgtt cttcacctca 4380 tgtttgcggg agagattcat tctcttccgt tttttattta aagcggcttt tccagacggg 4440 aacggtgttt tgtggtctcc attttcattt gccgataggc gaacgctaaa aatggcaggc 4500

cgagcagggt aatgccgctc aggacagaaa aaatataaat cggccggcca gcgccaaaca 4560 ggtctataca tatccccccg acccaagggc cgatgacgtt tccgagctgt ggaaaaccga 4620 ttgccccgaa ataagtgcct tttaatcctg gttttgcaat ctggtctaca tacaaatcca 4,680 tcatagagaa taaaagcact tcgccgattg taaatgtgat gacaatcatc acaattgatg 4740 gaacacegtg tgatacggtg aaaatggcca tgctgatgct aaccatcaca ttaccgagca 4800 tcagagaaca aagcggcgaa aaccgttttg caaaatggac aatgggaaat tgcgtcgcca 4860 acacaacgat tgcgtttaat gtcagcatca gcccatacag cttcgttcca ttgccgatca 4920 aggggttctg cgccatatac tgagggaatg tggaactgaa ttgtgagtag ccgaaggtgc 4980 atagcgtaat gccgaccaaa gcaatggtaa aaagataatc cttttgcgtg accataaacg 5040 cttcccgcac gctcatattt cgggactggg ctggtgctga taaggatgga tgttttttaa 5100 attggagggc aagcacaatt ccgtatagtc cgtaaatgac tgcaggcacc aaaaagggcg 5160 tagtcgattg cgatgagccg aaatataggc caagcacagg tccgaagaca acgccgatat 5220 taatageege atagegtaaa ttaaaaacta geagtetegt tttttettet gteatateag 5280 acaacaagge etttgaageg ggeteaaaca gtgatttgea aagacegttt aatgegttta 5340 🔆 ctacaaaaaa cacccagaga ttagatgctg ccgcaaagcc tgcaaatacc agcatccatc 5400 cgaaaatcga tacaagcatc atgttttttc tgccgaattt atctgagata tatccgccgt 5460 aaaagettge gaggatgeeg aetgatgage tegeggegat gaccageect geataggaag 5520 · ctgatgcgcc ttggacggct gtcaaataaa tcgctaaaaa aggaatgctc atcgatgttg 5580 ccattetgcc gaaaatggtt ccgattataa ttgtaacgcg ttggatgcat agettgagta 5640 ttctatagtg tcacctaaat agcttggcgt aatcatggtc atagctgttt cctgtgtgaa 5700 attgttatcc gctcacaatt ccacacaaca tacgagccgg aagcataaag tgtaaagcct 5760 ggggtgccta atgagtgagc taactcacat taattgcgtt gcgctcactg cccgctttcc 5820 agtcgggaaa cctgtcgtgc cagctgcatt aatgaatcgg ccaacgcgcg gggagaggcg 5880 gtttgcgtat tgggcgctct tccgcttcct cgctcactga ctcgctgcgc tcggtcgttc 5940 ggctgcggcg agcggtatca gctcactcaa aggcggtaat acggttatcc acagaatcag 6000 gggataacgc aggaaagaac atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa 6060 aggccgcgtt gctggcgttt ttcgataggc tccgccccc tgacgagcat cacaaaaatc 6120 gacgctcaag tcagaggtgg cgaaacccga caggactata aagataccag gcgtttcccc 6180 ctggaagete eetegtgege teteetgtte egaceetgee gettaeegga taeetgteeg 6240

cctttctccc ttcgggaage gtggcgcttt ctcatagete acgctgtagg tatctcagtt 6300 cggtgtaggt cgttcgctcc aagctgggct gtgtgcacga accccccgtt cagcccgacc 6360 gctgcgcctt atccggtaac tatcgtcttg agtccaaccc ggtaagacac gacttatcgc 6420 cactggcagc agecactggt aacaggatta gcagagcgag gtatgtaggc ggtgctacag 6480. agttettgaa gtggtggeet aactaegget acactagaag gacagtattt ggtatetgeg 6540 ctctgctgaa gccagttacc ttcggaaaaa gagttggtag ctcttgatcc ggcaaacaaa 6600 ccaccgctgg tagcggtggt ttttttgttt gcaagcagca gattacgcgc agaaaaaag 6660 gateteaaga agateetttg atetttteta eggggtetga egeteagtgg aaegaaaaet 6720 cacgttaagg gattttggtc atgagattat caaaaaggat cttcacctag atccttttaa 6780 attaaaaatg aagttttaaa tcaatctaaa gtatatatga gtaaacttgg tctgacagtt 6840 accaatgctt aatcagtgag gcacctatct cagcgatctg tctatttcgt tcatccatag 6900 ttgcctgact ccccgtcgtg tagataacta cgatacggga gggcttacca tctggcccca 6960. gtgctgcaat gataccgcga gacccacgct caccggctcc agatttatca gcaataaacc 7020 · · agccagccgg aagggccyag cgcagaagtg gtcctgcaac tttatccgcc tccatccagt 7080 ctattaattg ttgccgggaa gctagagtaa gtagttcgcc agttaatagt ttgcgcaacg 7140. ttgttggcat tgctacaggc atcgtggtgt cacgctcgtc gtttggtatg gcttcattca 7200. gctccggttc ccaacgatca aggcgagtta catgatcccc catgttgtgc aaaaaagcgg 7260: ttageteett eggteeteeg ategttgtea gaagtaagtt ggeegeagtg ttateactea 7320 tggttatggc agcactgcat aattctctta ctgtcatgcc atccgtaaga tgctttctg 7380 tgactggtga gtactcaacc aagtcattct gagaataccg cgcccggcga ccgagttgct 7440 cttycccggc gtcaatacgg gataatagtg tatgacatag cagaacttta aaagtgctca 7500 tcattggaaa acgttcttcg gggcgaaaac tctcaaggat cttaccgctg ttgagatcca 7560 gttcgatgta acceactcgt geacceaact gatetteage atettttact tteaceageg 7620 tttctgggtg agcaaaaaca ggaaggcaaa atgccgcaaa aaagggaata agggcgacac 7680 ggaaatgttg aatactcata ctcttccttt ttcaatatta ttgaagcatt tatcagggtt 7740 attgtctcat gagcggatac atatttgaat gtatttagaa aaataaacaa ataggggttc 7800 cgcgcacatt teccegaaaa gtgccacetg tatgcggtgt gaaatacege acagatgcgt 7860 aaggagaaaa taccgcatca ggcgaaattg taaacgttaa tattttgtta aaattcgcgt 7920 taaatatttg ttaaatcagc tcattttta accaataggc cgaaatcggc aaaatccctt 7980

ataaatcaaa agaatagac gagatagggt tgagtgttgt tecagtttgg aacaagagtc 8040 cactattaaa gaacgtggac tecaacgtca aagggcgaaa aaccgtctat cagggcgatg 8100 geccactacg tgaaccatca eccaaatcaa gttttttgeg gtegaggtge egtaaagete 8160 taaateggaa ecctaaaggg agececegat ttagagettg aeggggaaag ecggegaacg 8220 tggegagaaa ggaagggaag aaagegaaag gagegggege tagggegetg geaagtgtag 8280 eggteaeget gegegtaace accacaccg ecgegttaa

<210> 74

<211> 250

<212> PRT

<213> Clostridium acetobutylicum

<400> 74

Asn Lys Arg Ala Ala Phe Met Leu Leu Phe Leu Arg Ser Val Leu 1 5 10 15

Lys Val Ile Leu Val Leu Asp Val Gly Asn Thr Asn Ile Val Leu Gly 20 25 30

Ile Tyr Asn Asp Thr Lys Leu Thr Ala Glu Trp Arg Leu Ser Thr Asp 35 40 45

Val Leu Arg Ser Ala Asp Glu Tyr Gly Ile Gln Val Met Asn Leu Phe 50 55 60

Gln Gln Asp Lys Leu Asp Pro Thr Leu Val Glu Gly Val Ile Ile Ser 65 70 75 80

Ser Val Val Pro Asn Ile Met Tyr Ser Leu Glu His Met Ile Arg Lys 85 90 95

Tyr Phe Lys Ile Asn Pro Leu Val Val Gly Pro Gly Ile Lys Thr Gly 100 105 110

Ile Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile
115 120 125

Val Asn Ala Val Ala Ala His Glu Ile Tyr Lys Arg Ser Leu Ile Ile 130 135 140

Ile Asp Phe Gly Thr Ala Thr Thr Phe Cys Ala Val Arg Glu Asn Gly 145 150 155 160

Asp Tyr Leu Gly Gly Ala Ile Cys Pro Gly Ile Lys Val Ser Ser Glu 165 170 175

Ala Leu Phe Glu Lys Ala Ala Lys Leu Pro Arg Val Glu Leu Ile Lys 180 185 190

Pro Ala Tyr Ala Ile Cys Lys Asn Thr Ile Ser Ser Ile Gln Ser Gly
195 200 205

- 88 -

Ile Val Tyr Arg Tyr Leu Arg Gln Val Lys Tyr Leu Phe Glu Lys Leu

Lys Glu Asn Leu Pro Asp Gly Arg Arg Thr Arg Thr Ser Leu Val Leu 235

Ala Thr Gly Gly Leu Ala Lys Leu Ile Asn 245

<210>.75

<211> 258

<212> PRT

<213> Rhodobacter capsulatus

<400> 75

Met Leu Leu Cys Ile Asp Cys Gly Asn Thr Asn Thr Val Phe Ser Val

Trp Asp Gly Thr Asp Phe Ala Ala Thr Trp Arg Ile Ala Thr Asp His 25

Arg Arg Thr Ala Asp Glu Tyr Phe Val Trp Leu Asn Thr Leu Met Gln 40 45

Leu Lys Gly Leu Gln Gly Arg Ile Ser Glu Ala Ile Ile Ser Ser Thr

Ala Pro Arg Val Val Phe Asn Leu Arg Val Leu Cys Asn Arg Tyr Phe

Asp Cys Arg Pro Tyr Val Val Gly Lys Pro Gly Cys Glu Leu Pro Val

Ala Pro Arg Val Asp Pro Gly Thr Thr Val Gly Pro Asp Arg Leu Val

Asn Thr Val Ala Gly Tyr Asp Arg His Gly Gly Asp Leu Ile Val Val

Asp Phe Gly Thr Ala Thr Thr Phe Asp Val Val Ala Pro Asp Gly Ala

Tyr Ile Gly Gly Val Ile Ala Pro Gly Val Asn Leu Ser Leu Glu Ala 150 155

Leu His Met Ala Ala Ala Ala Leu Pro His Val Asp Val Thr Lys Pro

Gln Gly Val Ile Gly Thr Asn Thr Val Ala Cys Ile Gln Ser Gly Val 185

Tyr Trp Gly Tyr Ile Gly Leu Val Glu Gly Ile Val Arg Gln Ile Arg

Met Glu Arg Asp Arg Pro Met Lys Val Ile Ala Thr Gly Gly Leu Ala 215

Ser Leu Phe Asp Leu Gly Phe Asp Leu Phe Asp Lys Val Glu Asp Asp 225 230 235 240

Leu Thr Met His Gly Leu Arg Leu Ile Phe Asp Tyr Asn Lys Gly Leu 245 250 255

Gly Ala

<210> 76

<211> 10801

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
pAN240 plasmid

<400> 76

gaattttgcg gccgcttcga aagctgtaat ataaaaacct tcttcaacta acggggcagg 60 ttagtgacat tagaaaaccg actgtaaaaa gtacagtcgg cattatctca tattataaaa 120 gccagtcatt aggcctatct gacaattcct gaatagagtt cataaacaat cctgcatgat 180 aaccatcaca aacagaatga tgtacctgta aagatagcgg taaatatatt gaattacctt 240 tattaatgaa tittcctgct gtaataatgg gtagaaygta attactatta ttattgatat 300 ttaagttaaa cccagtaaat gaagtccatg gaataataga aagagaaaaa gcattttcag 360 gtataggtgt tttgggaaac aatttccccg aaccattata tttctctaca tcagaaaggt 420 ataaatcata aaactctttg aagtcattct ttacaggagt ccaaatacca gagaatgttt 480 tagatacacc atcaaaaatt gtataaagtg gctctaactt atcccaataa cctaactctc 540 cgtcgctatt gtaaccagtt ctaaaagctg tatttgagtt tatcaccctt gtcactaaga 600 aaataaatgc agggtaaaat ttatatcctt cttgttttat gtttcggtat aaaacactaa 660 tatcaatttc tgtggttata ctaaaagtcg tttgttggtt caaataatga ttaaatatct 720 cttttctctt ccaattgtct aaatcaattt tattaaagtt catttgatat gcctcctaaa 780 tttttatcta aagtgaattt aggaggetta ettgtetget ttetteatta gaatcaatee 840 ttttttaaaa gtcaatatta ctgtaacata aatatatatt ttaaaaaatat cccactttat 900 ccaattttcg tttgttgaac taatgggtgc tttagttgaa gaataaagac cacattaaaa 960 aatgtggtct tttgtgtttt tttaaaggat ttgagcgtag cgaaaaatcc ttttctttct 1020 tatcttgata ataagggtaa ctattgaatt cggtaccaag agtttgtaga aacgcaaaaa 1080 ggccatccgt caggatggcc ttctgcttaa tttgatgcct ggcagtttat ggcgggcgtc 1140

ctgcccgcca ccctccgggc cgttgcttcg caacgttcaa atccgctccc ggcggatttg 1200 tectactcag gagagegtte accgacaaac aacagataaa acgaaaggce cagtettteg 1280 actgageett tegttttatt tgatgeetgg cagtteeeta etetegeatg gggagaeece 1320 acactaccat cggcgctacg gcgtttcact tctgagttcg gcatggggtc aggtgggacc 1380 accgcgctac tgccgccagg caaattctgt tttatcagac cgcttctgcg ttctgattta 1440 atotgtatca ggotgaaaat ottototoat cogocaaaac aggatootac ggaaatggag 1500 cggcaaaacc gttttactct caaaatctta aaagaaaacc cccgataaag ggggcttttc 1560 ttctacaaaa ttgtacgggc tggttcgttc cccagcattt gttcaatttt gttttgatca 1620 ttcagaacag ccactttcgg ctcatggctt gccgcttctt gatcagacat cattttgtag 1680 gaaataataa tgaccttatc teetteetge acaaggegtg eggetgeace gtttaageat 1740 atgacgccgc ttccccgttt accaggaata atatacgttt caagacgtgc tccattatta 1800 ttattoacaa tttgtacttt ttcattagga agcattocca cagcatcaat gagatottca 1860 tcaattgtaa tgcttcccac atagttcagg tttgcttccg taacagttgc cctgtgaagt 1920 ttgccgctca tcattgttcg atacatatta tattctctcc atttctcgaa tatcaataat 1980 gatattatet attaaacgeg ettttgaaaa ageaactgea acagegagaa teatetttee 2040 agcaatttca ttcacaggct cgagttccgg ataggaataa agctctacat agtctatggt 2100 tecgetagte gtttcaatga tatettttge agettttate accgetteag gatetettte 2160 accggcttgg acaagttccg cacttgtttg aagggcccga tacagcttag gcgcttcttt 2220 tettteetea getgttaagt atacattgeg agagettttg getaageegt etteetete 2280 gaccgtatcg acaggaacca attcaatatc catgaagaag tegetgatta acccatcaac 2340 aacagetace tgctgcgcat ettttaaace gaaataggca egagtegget tgactagatt 2400 gaaaagette gteagtaega tegegaeeee gteaaaatgt eettetettg agegeeegea 2460 taacacgtct gtgcgtcttt ctacatgaat cgtgacattc ttttcaccgg gatacatatc 2520 atgagcatet ggcgtaaaaa gaatategae teeggegttt tetgeaagag etgeateeeg 2580 ctcaatatcg cgcggatatg cttcaaaatc ttcattaggg ccgaattgtg caggattcac 2640 aaaaatactc ataataacgg cgtcgttttc ttgtcttgct ttgtctgcta aggttaaatg 2700 cccctcatgc agaaacccca tcgtcggaac aaatccgatt gacttgccct ctgaatggta 2760 ttgttttatg gettettea getgtgaaat ateagtaate tgteteatet tatttteece 2820 cgtacaagee gteaageact gtetggttea tttgaaagga atgettttgt teagggaaag 2880 cacgatgtct tacatcctga acatatccgc tgattgctgt ttcgatggtt tcatcaatgc 2940 gcgtatattg ctttacaaat ttaggtgttc tctcaacacc gtggccgata atatcatgat 3000 aaacgagaac ttgtccgtcc gctttcacac cagccccgat tccaatgacc ggtatgctta 3060 gcgtctcggc aattttggct gtgagttctg ccggcacaca ttccagcaca agcatcatag 3120 ctcctgcttc ttcgcatttt atactgtctt ctattaattt tttggcgctt tgttcgtctt 3180 tgccctgtac tttatagccg cccagtacgc cgactgactg cggtgtcaaa cctaagtgac 3240 tgactactgg aatgcctcca agcgtcaatg cgcgaatgga ttcaaacacg ccttctccgc 3300 cctcaagett cagtgegtea gctccgcttt cctgaacgat agccgctgca tttttcagcg 3360 tatcttcctt agacaggtga taagacataa acggcatatc tgtcacaata aaggtattcg 3420 gcgcacccct tttaacggct tttgtatgat ggatcatgtc cgcaactgtc acaccgacag 3480 ttgaatcaag gccgaggacg accattccaa gtgaatcacc gactaaaatc atgtcaactc 3540 ccgcttgttc agcaagttta gctgccggat aatcataagc ggtcagcatg acaatcggtt 3600 gagegteetg etgttgttaa gattattata ecacacettg tagataaagt caacaacttt 3720 . ttgcaaaatt tttcaggaat tttagcagag gttgttctgg atgtagaaca aaacatcttt 3780 ccgctcttgt gctgttagga tatctttctt ggaagctagg taggcctcga gttatggcag 3840. ttggttaaaa ggaaacaaaa agaccgtttt cacacaaaac ggtctttttc gatttctttt 3900 tacagtcaca gccacttttg caaaaaccgg acagcttcat gccttataac tgctgtttcg 3960 gtcgacgatg atctgccgtt ttcttctgca agccaaaaaa ccttccgtta caacgagaag 4020 gattetteae tttetaaagt teggegagtt teatecetet gteecagtee ttttttggat 4080 caaggcagac tgctgcaatg tctatctatt ttaataatag gtgcagttcg caggcgatac 4140 tgcccaatgg aagtatacca aaatcaacgg gcttgtacca acacattagc ccaattcgat 4200 atcggcagaa tagattttt taatgccttc gttcgtttct aaaagcagaa cgccttcatc 4260 atctatacct aacgccttac cgtaaaaggt tccgtttaac gttctggctc tcatattagt 4320 gccaataccg agcgcatage tttcccataa aagcttaatc ggcgtaaatc cgtgcgtcat 4380 ataatcccgg taccgtttct caaagcatag taaaatatgc tggatgacgc cggcccgatc 4440 aattttttcc ccagcagctt ggctgaggct tgtcgcgatg tccttcaatt catctggaaa 4500 atcattagge tgctggttaa cgttaatgce gatcccaatg atcactgaac gtacgcggtc 4560 ttcttcagcc tgcatttccg ttaggatacc gactgttttt tttccgttaa tcaaaatatc 4620

atttggccat ttaatatccg tttggatgcc tgctgcctct tctattccct gcacaacagc 4680 tactgcagca agcagagtca gctgcggtgt tttttggagc ggaatgtcag gccgcaaaat 4740 caggeteate caaacacegt tteettettg agaatgeeat accetagaca tteggeeect 4800 teeggetgtt tgtttgteag ceaceaeag ggtgeettee ggtgegttat tattegegag 4860 ctcatgagcc gttttttgcg tgcttgaaag aacgtcatgg taaataagat gctggcccat 4920 cacttccgtt tttaatccaa aacgaatttc gctttcactg agttttccgg gttttttgat 4980 gagoogatat cotttette taacggotte tactteataa ecetettee gaagetette 5040 aatatgette cacacageag ttettgaaca geegagagea teaetgattt tttggeegga 5100 aataaattca ttgccggcct gagaaaataa ttcaataagg tctttctta atgttgaccg 5160 catgtettea gecaeteete tatgtgttte ttttgattgg agagetteee tgteacaaca 5220 geotgetega tecaetgtaa ttettetgae acceattte eggeeggeeg gtttegaage 5280 gcaagcaagt cettaccegt gatatcaaga teettaagge tittgategg caggittiga 5340 taagegtaet gaatgteett eagtttettt teateeagtt titegtittg eegaageige 5400. gatattttgg ccgctgagag cagtgctttt ttcccagctc tgtacattgt cattgcgtca 5460 aggetetgge caaacgtate ggcaatgtga atggetteet tgateaettt teeegggage 5520 ttccaggctt tcaggaaaag gggcgcgtct ttcaaaacta tgccaaggtt aattaaaaga 5580 gcagcccaaa gctcctcacg ggatgttaaa gagaagaatg gaaactcact cgttgaaatc 5640 aggttttctc gtttatgata aaaaccagga agctcttcat acaatctcgt ttgaatgagt 5700 gtttgaageg cetggegaga agetetteee tgeageaatt teteaaaete tatagttttt 5760 cgttcgactg aaacatggga gaggagtgat ttttctttcg caatggcttc ttctgtttcc 5820 ggtgaaagcg taaagccaag ctggctcata aagcgtacgg ctctcagcat acgaagcgca 5880 tectettgaa atetateete aggettteea aeggttegaa teaetttetg ateaatatet 5940 ttcttgccgc caaaataatc aagcaccttc ccgtccgctg tcatggccat cgcattgatc 6000 gttaaatctc tgcgttttag atcctcttct aatgatgaga taaattgcac ttctgacggt 6060 cttctgaaat caacataatc agattcagtc cggaatgtcg tgacttcata ggtttcatcc 6120 toccagagoa caataatggt coogtgotot ttgoctacat caacagtoog otgaaacago 6180 cgttctactt gatcaggtgc cgcatctgtc gcgatatcga catctccgat cgttcgtttc 6240 atatagetgt cacgaactge geceeegaca aaataageet gatggeeege ttegattaag 6300 atgeggagea egggaagtge tttgataaaa aettttteea tgtgateaet eeggttetge 6360

taaatcggca taaatctgtt catactggct gacaattttt ttagaagaaa attcattttc 6420 aagcatetet attgeegeet tigitaaaaeg attgettage igiteatett étaaaatget 6480 categegegg getgttgegg eegtaacate acegacatee aceaaaaate egeteacatt 6540 gttttttata acctcaggga taccgccaat gtttgttcca atacaaggca ctccgcaagc 6600 catcgcttca agcaggacaa ggccaaagct ttcttttca gatagcagca gcttcaaatc 6660 gctaatagaa taaagatett caacaeggte ttgattteca agcattaaga ettggtette 6720 caagecatat tttetgataa getegeagge tgtegattte teeggacegt eteegactaa 6780 aagcagette getttegttt tgeeagegat attgeggaac acaeggatga cateetgeae 6840 gcgtgcagcc actggtaaca ggattagcag agcgaggtat gtaggcggtg ctacagagtt 6900 cttgaagtgg tggcctaact acggctacac tagaaggaca gtatttggta tctgcgctct 6960 gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca aacaaaccac 7020 cgctggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc 7080 tcaagaagat cetttgatet tttetaeggg gtetgaeget cagtggaacg aaaactcaeg 7140 ttaagggatt ttggtcatga gattatcaaa aaggatcttc acctagatcc ttttaaatta 7200'. aaaatgaagt tttaaatcaa tctaaagtat atatgagtaa acttggtctg acagttacca 7260 . atgcttaatc agtgaggcac ctatctcagc gatctgtcta tttcgttcat ccatagttgc 7320. ctgactcccc gtcgtgtaga taactacgat acgggagggc ttaccatctg gccccagtgc 7380 tgcaatgata ccgcgagacc cacgctcacc ggctccagat ttatcagcaa taaaccagcc 7440 ageeggaagg geegagegea gaagtggtee tgeaacttta teegeeteea teeagtetat 7500 taattgttgc cgggaagcta gagtaagtag ttcgccagtt aatagtttgc gcaacgttgt 7560 tgccattgct acaggcatcg tggtgtcacg ctcgtcgttt ggtatggctt cattcagctc 7620 cggttcccaa cgatcaaggc gagttacatg atcccccatg ttgtgcaaaa aagcggttag 7680 ctccttcggt cctccgatcg ttgtcagaag taagttggcc gcagtgttat cactcatggt 7740 tatggcagca ctgcataatt ctcttactgt catgccatcc gtaagatgct tttctgtgac 7800 tggtgagtac tcaaccaagt cattctgaga atagtgtatg cggcgaccga gttgctcttg 7860 cccggcgtca atacgggata ataccgcgcc acatagcaga actttaaaag tgctcatcat 7920 tggaaaacgt tcttcggggc gaaaactctc aaggatctta ccgctgttga gatccagttc 7980 gatgtaaccc actcgtgcac ccaactgatc ttcagcatct tttactttca ccagcgtttc 8040 tgggtgagca aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggg cgacacggaa 8100

atgttgaata ctcatactct teetttttca atattattga agcatttate agggttattg 8160 teteatgage ggatacatat tigaatgtat tiagaaaaat aaacaaatag gggtteegeg 8220 cacatttece egaaaagtge cacetgaegt etaagaaace attattatea tgacattaac 8280 ctataaaaat aggcgtatca cgaggccctt tcgtctcgca tgcggatcag tgagggtttq 8340 caactgcggg tcaaggatct ggatttcgat cacggcacga tcatcgtgcg ggagggcaag 8400 ggctccaagg atcgggcctt gatgttaccc gagagcttgg cacccagcct gcgcqaqcaq 8460 gggaattgat ccggtggatg accttttgaa tgacctttaa tagattatat tactaattaa 8520 ttggggaccc tagaggtccc cttttttatt ttaaaaattt tttcacaaaa cggtttacaa 8580 gcataacggg ttttgctgcc cgcaaacggg ctgttctggt gttgctagtt tgttatcaga 8640 ategeagate eggetteagg titgeegget gaaagegeta titetteeag aattgeeatg 8700 attttttccc cacgggaggc gtcactggct cccgtgttgt cggcagcttt gattcgataa 8760 gcagcatcgc ctgtttcagg ctgtctatgt gtgactgttg agctgtaaca agttgtctca 8820 ggtgttcaat ttcatgttct agttgctttg ttttactggt ttcacctgtt ctattaggtg 8880. ttacatgctg ttcatctgtt acattgtcga tctgttcatg gtgaacagct ttaaatgcac 8940 caaaaactcg taaaagctct gatgtatcta tcttttttac accgttttca tctgtgcata 9000 tggacagttt tecetttgat atetaaeggt gaacagttgt tetaettttg tttgttagte 9060 ttgatgcttc actgatagat acaagagcca taagaacctc agatecttcc gtatttagcc 9120 aqtatgttct ctagtgtggt tcgttgtttt tgcgtgagcc atgagaacga accattgaga 9180 tcatgcttac tttgcatgtc actcaaaaat tttgcctcaa aactggtgag ctgaattttt 9240 gcagttaaag catcgtgtag tgtttttctt agtccgttac gtaggtagga atctgatqta 9300 atggttgttg gtattttgtc accattcatt tttatctggt tgttctcaag ttcggttacq 9360 agatecattt gtetatetag tteaacttgg aaaateaacg tateagtegg geggeetege 9420 ttatcaacca ccaatttcat attgctgtaa gtgtttaaat ctttacttat tggtttcaaa 9480 acceattggt taageetttt aaacteatgg tagttatttt caageattaa catgaactta 9540 aattcatcaa ggctaatctc tatatttgcc ttgtgagttt tcttttgtgt tagttctttt 9600 aataaccact cataaatcct catagagtat ttgttttcaa aagacttaac atgttccaga 9660 ttatatttta tgaatttttt taactggaaa aqataaqqca atatctcttc actaaaaact 9720 aattctaatt tttcgcttga gaacttggca tagtttgtcc actggaaaat ctcaaagcct 9780 ttaaccaaag gattootgat ttocacagtt ctcgtcatca gctctctggt tgctttagct 9840

aatacaccat aagcattttc cctactgatg ttcatcatct gagcgtattg gttataagtg 9900 aacgataccg tccgttcttt ccttgtaggg ttttcaatcg tggggttgag tagtgccaca 9960 cagcataaaa ttagcttggt ttcatgctcc gttaagtcat agcgactaat cgctagttca 10020 tttgctttga aaacaactaa ttcagacata catctcaatt ggtctaggtg attttaatca 10080 ctataccaat tgagatgggc tagtcaatga taattactag teetttteet ttgagttgtg 10140 ggtatctgta aattctgcta gacctttgct ggaaaacttg taaattctgc tagaccctct 10200 gtaaattccg ctagaccttt gtgtgttttt tttgtttata ttcaagtggt tataatttat 10260 agaataaaga aagaataaaa aaagataaaa agaatagate eeageeetgt gtataaetea 10320 ctactttagt cagttccgca gtattacaaa aggatgtcgc aaacgctgtt tgctcctcta 10380 caaaacagac cttaaaaccc taaaggctta agtagcaccc tcgcaagctc gggcaaatcg 10440 ctgaatattc cttttgtctc cgaccatcag gcacctgagt cgctgtcttt ttcgtgacat 10500 tcagttcgct gcgctcacgg ctctggcagt gaatgggggt aaatggcact acaggcgcct 10560 tttatggatt catgcaagga aactacccat aatacaagaa aagcccgtca cgggcttctc 10620 agggcgtttt atggcgggtc tgctatgtgg tgctatctga cttttcgctg ttcagcagtt 10680 cetgeeetet gatttteeag tetgaceact teggattate cegtgaeagg teatteagae 10740 . tggctaatgc acccagtaag gcagcggtat catcaacagg cttacccgtc ttactgtcaa 10800 С 10801

<400> 77

ctcgaggcct acctagcttc caagaaagat atcctaacag cacaagagcg gaaagatgtt 60 ttgttctaca tccagaacaa cctctgctaa aattcctgaa aaattttgca aaaagttgtt 120 gactttatct acaaggtgtg gtataataat cttaacaaca gcaggacgct ctagaggagg 180 agacatcatg aaaattggaa ttatcggcgg aggctccgtt ggtcttttat gcgcctatta 240 tttgtcactt tatcacgacg tgactgttgt gacgaggcgg caagaacagg ctgcggccat 300 tcagtctgaa ggaatccggc tttataaagg cggggaggaa ttcagggctg attgcagtgc 360

<210> 77

<211> 8654

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN236 plasmid

ggacacgagt atcaattcgg actttgacct gcttgtcgtg acagtgaagc agcatcagct 420 tcaatctgtt ttttcgtcgc ttgaacgaat cgggaagacg aatatattat ttttgcaaaa 480 cggcatgggg catatccacg acctaaaaga ctggcacgtt ggccattcca tttatgttgg 540 aatcqttgag cacggagctg taagaaaatc ggatacagct gttgatcata caggcctagg 600 tgcqataaaa tggagcgcgt tcgacgatgc tgaaccagac cggctgaaca tcttgtttca 660 gcataaccat teggattite egattiatta tgagaeggat tggtaeegte tgetgaeggg 720 caagetgatt gtaaatgegt gtattaatee tttaactgeg ttattgeaag tgaaaaatgg 780 agaactgctg acaacgccag cttatctggc ttttatgaag ctggtatttc aggaggcatg 840 ccgcatttta aaacttgaaa atgaagaaaa ggcttgggag cgggttcagg ccgtttgtgg 900 gcaaacgaaa gagaatcgtt catcaatgct ggttgacgtc attggaggcc ggcagacgga 960 agctgacgcc attatcggat acttattgaa ggaagcaagt cttcaaggtc ttgatgccgt 1020 ccacctagag tttttatatg gcagcatcaa agcattggag cgaaatacaa acaaagtctt 1080 ttgagctttt tcggtaacat gctatactca tttcggatca ctaactattt attggaggat 1140 cctgttttgg cggatgagag aagattttca gcctgataca gattaaatca gaacgcagaa 1200 geggtetgat aaaacagaat ttgeetggeg geagtagege ggtggteeca eetgaeeeca 1260 tgccgaactc agaagtgaaa cgccgtagcg ccgatggtag tgtgggggtct ccccatgcga 1320 gagtagggaa ctgccaggca tcaaataaaa cgaaaggctc agtcgaaaga ctgggccttt 1380 cgttttatct gttgtttgtc ggtgaacgct ctcctgagta ggacaaatcc gccgggagcg 1440 gatttgaacg ttgcgaagca acggcccgga gggtggcggg caggacgccc gccataaact 1500 qccaqgcatc aaattaagca gaaggccatc ctgacggatg gcctttttgc gtttctacaa 1560 actettggta eecagaaaaa geggeaaaag eggetgttaa aaaagegaaa tegaagaage 1620 tgtctgccgc taagacggaa tatcaaaagc gttctgctgt tgtgtcatct ttaaaaagtca 1680 cagccgatga atcccagcaa gatgtcctaa aatacttgaa cacccagaaa gataaaggaa 1740 atgcagacca aattcattct tattatgtgg tgaacgggat tgctgttcat gcctcaaaag 1800 aggttatgga aaaagtggtg cagtttcccg aagtggaaaa ggtgcttcct aatgagaaac 1860 ggcagctttt taagtcatcc tccccattta atatgaaaaa agcacagaaa gctattaaag 1920 caactgacgg tgtggaatgg aatgtagacc aaatcgatgc cccaaaagct tgggcacttg 1980 gatatgatgg aactggcacg gttgttgcgt ccattgatac cggggtggaa tggaatcatc 2040 cggcattaaa agagaaatat cgcggatata atccggaaaa tcctaatgag cctgaaaatg 2100

aaatgaactg gtatgatgee gtageaggeg aggeaageee ttatgatgat ttggeteatg 2160 gaacccacgt gacaggcacg atggtgggct ctgaacctga tggaacaaat caaatcggtg 2220 tagcacctgg cgcaaaatgg attgctgtta aagcgttctc tgaagatggc ggcactgatg 2280 ctgacatttt ggaagctggt gaatgggttt tagcaccaaa ggacgcggaa ggaaatcccc 2340 acceggaaat ggctectgat gttgtcaata actcatgggg agggggctct ggacttgatg 2400 aatggtacag agacatggtc aatgcctggc gttcggccga tattttccct gagttttcag 2460 cggggaatac ggatetettt atteceggeg ggeetggtte tategeaaat eeggeaaact 2520 atccagaatc gtttgcaact ggagcgactg agaattccaa ttccccatgg agagaaaaga 2580 aaatcgctaa tgttgattac tttgaacttc tgcatattct tgaatttaaa aaggctgaaa 2640 gagtaaaaga ttgtgctgaa atattagagt ataaacaaaa tcgtgaaaca ggcgaaagaa 2700 agttgtatcg agtgtggttt tgtaaatcca ggctttgtcc aatgtgcaac tggaggagag 2760 caatgaaaca tggcattcag tcacaaaagg ttgttgctga agttattaaa caaaagccaa 2820 cagttcgttg gttgtttctc acattaacag ttaaaaatgt ttatgatggc gaagaattaa 2880 ataagagttt gtcagatatg gctcaaggat ttcgccgaat gatgcaatat aaaaaaatta 2940 ataaaaatct tgttggtttt atgcgtgcaa cggaagtgac aataaataat aaagataatt 3000 cttataatca gcacatgcat gtattggtat gtgtggaacc aacttatttt aagaatacag 3060 aaaactacgt gaatcaaaaa caatggattc aattttggaa aaaggcaatg aaattagact 3120 atgatecaaa tgtaaaagtt caaatgatte gaeegaaaaa taaatataaa teggatatae 3180 aatcggcaat tgacgaaact gcaaaatatc ctgtaaagga tacggatttt atgaccgatg 3240 atgaagaaaa gaatttgaaa cgtttgtctg atttggagga aggtttacac cgtaaaaggt 3300 taatctccta tggtggtttg ttaaaagaaa tacataaaaa attaaacctt gatgacacag 3360 aagaaggcga tttgattcat acagatgatg acgaaaaagc cgatgaagat ggattttcta 3420 ttattgcaat gtggaattgg gaacggaaaa attattttat taaagagtag ttcaacaaac 3480 gggccatatt gttgtataag tgatgaaata ctgaatttaa aacttagttt atatgtggta 3540 caattaaaag agggaagcgt atcattaacc ctataaacta cgtctgccct cattattgga 3660 gggtgaaatg tgaatacatc ctattcacaa tcgaatttac gacacaacca aattttaatt 3720 tggctttgca ttttatcttt ttttagcgta ttaaatgaaa tggttttgaa cgtctcatta 3780 cctgatattg caaatgattt taataaacca cctgcgagta caaactgggt gaacacagcc 3840

tttatgttaa ccttttccat tggaacagct gtatatggaa agctatctga tcaattaggc 3900 atcaaaaggt tactcctatt tggaattata ataaattgtt tcgggtcggt aattgggttt 3960 qttqqccatt ctttcttttc cttacttatt atggctcgtt ttattcaagg ggctggtgca 4020 getgeattte cageactegt aatggttgta gttgegeget atattecaaa ggaaaatagg 4080 qqtaaagcat ttggtcttat tggatcgata gtagccatgg gagaaggagt cggtocagcg 4140 attgqtggaa tgatagccca ttatattcat tggtcctatc ttctactcat tcctatgata 4200 acaattatca ctqttccqtt tcttatgaaa ttattaaaga aagaagtaag gataaaaggt 4260 cattttgata tcaaaggaat tatactaatg tctgtaggca ttgtatttt tatgttgttt 4320 acaacatcat atagcatttc ttttcttatc gttagcgtgc tgtcattcct gatatttgta 4380 aaacatatca ggaaagtaac agatcctttt gttgatcccg gattagggaa aaatatacct 4440 tttatgattg gagttctttg tgggggaatt atatttggaa cagtagcagg gtttgtctct 4500 atggtteett atatgatgaa agatgtteac cagetaagta etgeegaaat eggaagtgta 4560 attattttcc ctggaacaat gagtgtcatt attttcggct acattggtgg gatacttgtt 4620 qatagaagag gtcctttata cgtgttaaac atcggagtta catttctttc tgttagcttt 4680 ttaactgctt cctttctttt agaaacaaca tcatggttca tgacaattat aatcgtattt 4740. qttttaggtg ggctttcgtt caccaaaaca gttatatcaa caattgtttc aagtagcttg 4800 aaacagcagg aagctggtgc tggaatgagt ttgcttaact ttaccagctt tttatcagag 4860. ggaacaggta ttgcaattgt aggtggttta ttatccatac ccttacttga tcaaaggttg 4920 ttacctatgg aagttgatca gtcaacttat ctgtatagta atttgttatt actttttca 4980 qgaatcattg tcattagttg gctggttacc ttgaatgtat ataaacattc tcaaagggat 5040 ttctaaatcg ttaagggatc aactttggga gagagttcaa aattgatcct ttttttataa 5100 cagttcgaag cggccgcaat tettgaagac gaaagggeet egtgataege etatttttat 5160 aggttaatgt catgataata atggtttett agaegteagg tggeaetttt eggggaaatg 5220 tgcgcggaac ccctatttgt ttatttttct aaatacattc aaatatgtat ccgctcatga 5280 gacaataacc ctgataaatg cttcaataat attgaaaaag gaagagtatg agtattcaac 5340 atttccgtgt cgcccttatt cccttttttg cggcattttg ccttcctgtt tttgctcacc 5400 cagaaacgct ggtgaaagta aaagatgctg aagatcagtt gggtgcacga gtgggttaca 5460 tcqaactqqa tctcaacagc ggtaagatcc ttgagagttt tcgccccgaa gaacgttttc 5520 caatgatgag cacttttaaa gttctgctat gtggcgcggt attatcccgt attgacgccg 5580

ggcaagagca acteggtege egcatacact atteteagaa tgaettggtt gagtaeteae 5640 cagtcacaga aaagcatett aeggatggea tgacagtaag agaattatge agtgetgeea 5700 taaccatgag tgataacact gcggccaact tacttctgac aacgatcgga ggaccgaagg 5760 agetaacege tttttgcae aacatggggg atcatgtaae tegeettgat egttgggaae 5820 cggagctgaa tgaagccata ccaaacgacg agcgtgacac cacgatgcct gcagcaatgg 5880 caacaacgtt gcgcaaacta ttaactggcg aactacttac tctagcttcc cggcaacaat 5940 taatagactg gatggaggcg gataaagttg caggaccact tetgcgcteg geeetteegg 6000 ctggctggtt tattgctgat aaatctggag ccggtgagcg tgggtctcgc ggtatcattg 6060 cagcactggg gccagatggt aagccctccc gtatcgtagt tatctacacg acggggagtc 6120 aggcaactat ggatgaacga aatagacaga tcgctgagat aggtgcctca ctgattaagc 6180 attggtaact gtcagaccaa gtttactcat atatacttta gattgattta aaacttcatt 6240 tttaatttaa aaggatctag gtgaagatcc tttttgataa tctcatgacc aaaatccctt 6300 aacgtgagtt ttcgttccac tgagcgtcag accccgtaga aaagatcaaa ggatcttctt 6360 gagateettt ttttetgege gtaatetget gettgeaaac aaaaaaacca eegetaccag 6420 eggtggtttg tttgeeggat caagagetae caactetttt teegaaggta aetggettea 6480 gcagagegea gataceaaat actgteette tagtgtagee gtagttagge caccaettea 6540 agaactetgt ageacegeet acataceteg etetgetaat eetyttacea gtggetgetg 6600 ccagtggcga taagtcgtgt cttaccgggt tggactcaag acgatagtta ccggataagg 6660 cgcagcggtc gggctgaacg gggggttcgt gcacacagcc cagcttggag cgaacgacct 6720 acaccgaact gagataccta cagcgtgagc tatgagaaag cgccacgctt cccgaaggga 6780 gaaaggcgga caggtateeg gtaageggea gggteggaae aggagagege acgagggage 6840 ttccaggggg aaacgcctgg tatctttata gtcctgtcgg gtttcgccac ctctgacttg 6900 agcgtcgatt tttgtgatgc tcgtcagggg ggcggagcct atggaaaaac gccagcaacg 6960 cggccttttt acggttcctg gccttttgct ggccttttgc tcacatgttc tttcctgcgt 7020 tateceetga ttetgtggat aacegtatta eegeetttga gtgagetgat aeegetegee 7080 gcagccgaac gaccgagcgc agcgagtcag tgagcgagga agcggaagag cgcctgatgc 7140 ggtattttct ccttacgcat ctgtgcggta tttcacaccg catatggtgc actctcagta 7200 caatctgctc tgatgccgca tagttaagcc agtatacact ccgctatcgc tacgtgactg 7260 ggtcatggct gcgccccgac acccgccaac acccgctgac gcgccctgac gggcttgtct 7320

geteceggea teegettaca gacaagetgt gacegtetee gggagetgea tgtgtcagag 7380 gttttcaccg tcatcaccga aacgcgcgag gcagctgcgg taaagctcat cagcgtggtc 7440 gtgaagcgat teacagatgt etgeetgtte atecgegtee agetegttga gttteteeag 7500 aagegttaat gtetggette tgataaageg ggeeatgtta agggeggttt ttteetgttt 7560 ggtcacttga tgcctccgtg taagggggaa tttctgttca tgggggtaat gataccgatg 7620 aaacgagaga ggatgctcac gatacgggtt actgatgatg aacatgcccg gttactggaa 7680 cgttgtgagg gtaaacaact ggcggtatgg atgcggcggg accagagaaa aatcactcag 7740 ggtcaatgcc agcgcttcgt taatacagat gtaggtgttc cacagggtag ccagcagcat 7800 cctgcgatgc agatccggaa cataatggtg cagggcgctg acttccgcgt ttccagactt 7860 tacgaaacac ggaaaccgaa gaccattcat gttgttgctc aggtcgcaga cgttttgcag 7920 cagcagtcgc ttcacgttcg ctcgcgtatc ggtgattcat tctgctaacc agtaaggcaa 7980 ccccgccagc ctagccgggt cctcaacgac aggagcacga tcatgcgcac ccgtggccag 8040 gacccaacgc tgcccgagat gcgccgcgtg cggctgctgg agatggcgga cgcgatggat 8100 atgttctgcc aagggttggt ttgcgcattc acagttctcc gcaagaattg attggctcca 8160 attettggag tggtgaatce gttagegagg tgeegeegge ttecattcag gtegaggtgg 8220 cccggctcca tycaccgcga cycaacgcgg ggaggcagac aaggtatagg gcggcgccta 8280 caatccatgo caaccogtto catgtgotog cogaggoggo ataaatogoo gtgacgatoa 8340 🦠 geggtecagt gategaagtt aggetggtaa gageegegag egateettga agetgteeet 8400 gatggtcgtc atctacctgc ctggacagca tggcctgcaa cgcgggcatc ccgatgccgc 8460 cggaagcgag aagaatcata atggggaagg ccatccagcc tcgcgtcggc ggccgcttcg 8520 tcgaccgaaa cagcagttat aaggcatgaa gctgtccggt ttttgcaaaa gtggctgtga 8500 ctgtaaaaag aaatcgaaaa agaccgtttt gtgtgaaaac ggtctttttg tttcctttta 8640 8654 accaactgcc ataa

<210> 78

<211> 8093

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN423 plasmid

<400> 78 ggcggccgct tcgtcgaccg aaacagcagt tataaggcat gaagctgtcc ggtttttgca 60

aaagtggctg tgactgtaaa aagaaatcga aaaagaccgt tttgtgtgaa aacggtcttt 120 ttgtttcctt ttaaccaact gccataactc gaggcctacc tagcttccaa gaaagatatc 180 ctaacagcac aagagcggaa agatgttttg ttctacatcc agaacaacct ctgctaaaat 240 tcctgaaaaa ttttgcaaaa agttgttgac tttatctaca aggtgtggta taataatctt 300 aacaacagca ggacgctcta gaaaaggagg aatttaaatg tatcgtacga tgatgagcgg 360 caaacttcac agggcaactg ttacggaagc aaacctgaac tatgtgggaa gcattacaat 420 tgatgaagat ctcattgatg ctgtgggaat gcttcctaat gaaaaagtac aaattgtgaa 480 taataataat ggagcacgtc ttgaaacgta tattattcct ggtaaacggg gaagcggcgt 540 catatgetta aacggtgcag cegeacgeet tgtgcaggaa ggagataagg teattattat 600 ttcctacaaa atgatgtctg atcaagaagc ggcaagccat gagccgaaag tggctgttct 660 gaatgatcaa aacaaaattg aacaaatgct ggggaacgaa ccagcccgta caattttgta 720 aaggateetg ttttggegga tgagagaaga tttteageet gatacagatt aaatcagaae 780 gcagaagcgg tctgataaaa cagaatttgc ctggcggcag tagcgcggtg gtcccacctg 840 accecatgee gaacteagaa gtgaaaegee gtagegeega tggtagtgtg gggteteece 900 atgcgagagt agggaactgc caggcatcaa ataaaacgaa aggctcagtc gaaagactgg 960 gcctttcgtt ttatctgttg tttgtcggtg aacgctctcc tgagtaggac aaatccgccg 1020 ggagcggatt tgaacgttgc gaagcaacgg cccggagggt ggcgggcagg acgcccgcca 1080 taaactgcca ggcatcaaat taagcagaag gccatcctga cggatggcct ttttgcgttt 1140 ctacaaactc ttggtaccca gaaaaagcgg caaaagcggc tgttaaaaaa gcgaaatcga 1200 agaagetgte tgccgctaag acggaatate aaaagegtte tgctgttgtg tcatetttaa 1260 aagtcacage egatgaatee cageaagatg teetaaaata ettgaacace cagaaagata 1320 aaggaaatgc agaccaaatt cattcttatt atgtggtgaa cgggattgct gttcatgcct 1380 caaaagaggt tatggaaaaa gtggtgcagt ttcccgaagt ggaaaaggtg cttcctaatg 1440 agaaacggca gctttttaag tcatcctccc catttaatat gaaaaaagca cagaaagcta 1500 ttaaagcaac tgacggtgtg gaatggaatg tagaccaaat cgatgcccca aaagcttggg 1560 cacttggata tgatggaact ggcacggttg ttgcgtccat tgataccggg gtggaatgga 1620 atcatccggc attaaaagag aaatatcgcg gatataatcc ggaaaatcct aatgagcctg 1680 aaaatgaaat gaactggtat gatgccgtag caggcgaggc aagcccttat gatgatttgg 1740 ctcatggaac ccacgtgaca ggcacgatgg tgggctctga acctgatgga acaaatcaaa 1800

teggtgtage acctggegea aaatggattg etgttaaage gttetetgaa gatggeggea 1860 ctgatgctga cattttggaa gctggtgaat gggttttagc accaaaggac gcggaaggaa 1920 atccccaccc ggaaatggct cctgatgttg tcaataactc atggggaggg ggctctggac 1980 ttgatgaatg gtacagagac atggtcaatg cctggcgttc ggccgatatt ttccctgagt 2040 tttcagcggg gaatacggat ctctttattc ccggcgggcc tggttctatc gcaaatccgg 2100 caaactatcc agaatcgttt gcaactggag cgactgagaa ttccaattcc ccatggagag 2160 aaaagaaaat cgctaatgtt gattactttg aacttctgca tattcttgaa tttaaaaagg 2220 ctgaaagagt aaaagattgt gctgaaatat tagagtataa acaaaatcgt gaaacaggcg 2280 aaagaaagtt gtatcgagtg tggttttgta aatccagget ttgtccaatg tgcaactgga 2340 ggagagcaat gaaacatggc attcagtcac aaaaggttgt tgctgaagtt attaaacaaa 2400 agccaacagt togttggttg tttctcacat taacagttaa aaatgtttat gatggcgaag 2460 aattaaataa gagtttgtca gatatggctc aaggatttcg ccgaatgatg caatataaaa 2520 aaattaataa aaatettgtt ggttttatge gtgeaaegga agtgaeaata aataataaag 2580 ataattotta taatcagcac atgcatgtat tggtatgtgt ggaaccaact tattttaaga 2640 atacagaaaa ctacgtgaat caaaaacaat ggattcaatt ttggaaaaag gcaatgaaat 2700 tagactatga tecaaatgta aaagtteaaa tgattegaee gaaaaataaa tataaategg 2760 atatacaatc ggcaattgac gaaactgcaa aatatcctgt aaaggatacg gattttatga 2820 ccgatgatga agaaaagaat ttgaaacgtt tgtctgattt ggaggaaggt ttacaccgta 2880 aaaggttaat ctcctatggt ggtttgttaa aagaaataca taaaaaatta aaccttgatg 2940 acacagaaga aggcgatttg attcatacag atgatgacga aaaagccgat gaagatggat 3000 tttctattat tgcaatgtgg aattgggaac ggaaaaatta ttttattaaa gagtagttca 3060 acaaacgggc catattgttg tataagtgat gaaatactga atttaaaact tagtttatat 3120 gtggtaaaat gttttaatca agtttaggag gaattaatta tgaagtgtaa tgaatgtaac 3180 agggttcaat taaaagaggg aagcgtatca ttaaccctat aaactacgtc tgccctcatt 3240 attggagggt gaaatgtgaa tacatcctat tcacaatcga atttacgaca caaccaaatt 3300 ttaatttggc tttgcatttt atctttttt agcgtattaa atgaaatggt tttgaacgtc 3360 tcattacctg atattgcaaa tgattttaat aaaccacctg cgagtacaaa ctgggtgaac 3420 acagoottta tgttaacott ttocattgga acagotgtat atggaaagot atctgatcaa 3480 ttaggcatca aaaggttact cctatttgga attataataa attgtttcgg gtcggtaatt 3540

gggtttgttg gccattcttt cttttcctta cttattatgg ctcgttttat tcaaggggct 3600 ggtgcagctg catttccagc actcgtaatg gttgtagttg cgcgctatat tccaaaggaa 3660 aataggggta aagcatttgg tottattgga togatagtag ocatgggaga aggagtoggt 3720 ccagcgattg gtggaatgat agcccattat attcattggt cctatcttct actcattcct 3780 atgataacaa ttatcactgt tccgtttctt atgaaattat taaagaaaga agtaaggata 3840 aaaggtcatt ttgatatcaa aggaattata ctaatgtctg taggcattgt attttttatg 3900 ttgtttacaa catcatatag catttctttt cttatcgtta gcgtgctgtc attcctgata 3960 tttgtaaaac atatcaggaa agtaacagat ccttttgttg atcccggatt agggaaaaat 4020 atacctttta tgattggagt tctttgtggg ggaattatat ttggaacagt agcagggttt 4080 gtctctatgg ttccttatat gatgaaagat gttcaccagc taagtactgc cgaaatcgga 4140 agtgtaatta ttttccctgg aacaatgagt gtcattattt tcggctacat tggtgggata 4200 cttgttgata gaagaggtcc tttatacgtg ttaaacatcg gagttacatt tctttctgtt 4260 agctttttaa ctgcttcctt tettttagaa acaacatcat ggttcatgac aattataatc 4320 gtatttgttt taggtgggct ttcgttcacc aaaacagtta tatcaacaat tgtttcaagt 4380 / agettgaaac agcaggaage tggtgetgga atgagtttge ttaaetttae cagettttta 4440 tcagagggaa caggtattgc aattgtaggt ggtttattat ccataccett acttgatcaa 4500 aggitgitac ctatggaagt tgatcagtca acttatctgt atagtaattt gitattacit 4560. ttttcaggaa tcattgtcat tagttggctg gttaccttga atgtatataa acattctcaa 4620 agggatttct aaatcgttaa gggatcaact ttgggagaga gttcaaaatt gatcetttt 4680 ttataacagt tcgaagcggc cgcaattett gaagacgaaa gggcctcgtg atacgcctat 4740 ttttataggt taatgtcatg ataataatgg tttcttagac gtcaggtggc acttttcggg 4800 gaaatgtgcg cggaacccct atttgtttat ttttctaaat acattcaaat atgtatccgc 4860 tcatgagaca ataaccctga taaatgcttc aataatattg aaaaaggaag agtatgagta 4920 ttcaacattt ccgtgtcgcc cttattccct tttttgcggc attttgcctt cctgtttttg 4980 ctcacccaga aacgctggtg aaagtaaaag atgctgaaga tcagttgggt gcacgagtgg 5040 gttacatcga actggatctc aacagcggta agatccttga gagttttcgc cccgaagaac 5100 gttttccaat gatgagcact tttaaagttc tgctatgtgg cgcggtatta tcccgtattg 5160 acgccgggca agagcaactc ggtcgccgca tacactattc tcagaatgac ttggttgagt 5220 actcaccagt cacagaaaag catcttacgg atggcatgac agtaagagaa ttatgcagtg 5280

ctgccataac catgagtgat aacactgcgg ccaacttact tctgacaacg atcggaggac 5340 cgaaggaget aacegetttt ttgcacaaca tgggggatca tgtaactege ettgategtt 5400 gggaaccgga gctgaatgaa gccataccaa acgacgagcg tgacaccacg atgcctgcag 5460 caatggcaac aacgttgcgc aaactattaa ctggcgaact acttactcta gcttcccggc 5520 aacaattaat agactggatg gaggcggata aagttgcagg accacttctg cgctcggccc 5580 ttccqqctqq ctqqtttatt gctgataaat ctggagccgg tgagcgtggg tctcgcggta 5640 tcattgcage actggggcca gatggtaage cetecegtat egtagttate taeaegaegg 5700 ggagtcaggc aactatggat gaacgaaata gacagatcgc tgagataggt gcctcactga 5760 ttaagcattg gtaactgtca gaccaagttt actcatatat actttagatt gatttaaaac 5820 ttcattttta atttaaaagg atctaggtga agatcctttt tgataatctc atgaccaaaa 5880 tecettaacg tgagtttteg ttecactgag egteagacce egtagaaaag atcaaaggat 5940 cttcttgaga tcctttttt ctgcgcgtaa tctgctgctt gcaaacaaaa aaaccaccgc 6000 taccageggt ggtttgtttg coggatcaag agetaccaac tettttteeg aaggtaactg 6060 getteageag agegeagata ceaaatactg teettetagt gtageegtag ttaggecace 6120 acttcaagaa ctctgtagca ccgcctacat acctcgctct gctaatcctg ttaccagtgg 6180 ctgctgccag tggcgataag tcgtgtctta ccgggttgga ctcaagacga tagttaccgg 6240 ataaggcgca gcggtcgggc tgaacggggg gttcgtgcac acagcccagc ttggagcgaa 6300 cgacctacac cgaactgaga tacctacage gtgagetatg agaaagegee aegetteeeg 6360 aagggagaaa ggcggacagg tatccggtaa gcggcagggt cggaacagga gagcgcacga 6420 gggagettee agggggaaae geetggtate tttatagtee tgtegggttt egeeacetet 6480 gacttgagcg tcgatttttg tgatgctcgt caggggggcg gagcctatgg aaaaacgcca 6540 gcaacgcggc ctttttacgg ttcctggcct tttgctggcc ttttgctcac atgttctttc 6600 ctgcgttatc ccctgattct gtggataacc gtattaccgc ctttgagtga gctgataccg 6660 ctcgccgcag ccgaacgacc gagcgcagcg agtcagtgag cgaggaagcg gaagagcgcc 6720 tgatgcggta ttttctcctt acgcatctgt gcggtatttc acaccgcata tggtgcactc 6780 tcagtacaat ctgctctgat gccgcatagt taagccagta tacactccgc tatcgctacg 6840 tqactqqqtc atqqctqcqc cccqacaccc qccaacaccc gctqacqcqc cctqacqqqc 6900 ttgtctgctc ccggcatccg cttacagaca agctgtgacc gtctccggga gctgcatgtg 6960 tcaqaggttt tcaccgtcat caccgaaacg cgcgaggcag ctgcggtaaa gctcatcagc 7020

gtggtcgtga agcgattcac agatgtctgc ctgttcatcc gcgtccagct cgttgagttt 7080 ctccagaagc gttaatgtct ggcttctgat aaagcgggcc atgttaaggg cggtttttc 7140 ctgtttggtc acttgatgcc tccgtgtaag ggggaatttc tgttcatggg ggtaatgata 7200 ccgatgaaac gagagaggat gctcacgata cgggttactg atgatgaaca tgcccggtta 7260 ctggaacgtt gtgagggtaa acaactggcg gtatggatgc ggcgggacca gagaaaaatc 7320 actcagggtc aatgccagcg cttcgttaat acagatgtag gtgttccaca gggtagccag 7380 cagcatectg cgatgcagat ceggaacata atggtgcagg gegetgaett eegegtttee 7440 agactttacg aaacacggaa accgaagacc attcatgttg ttgctcaggt cgcagacgtt 7500 ttgcagcagc agtcgcttca cgttcgctcg cgtatcggtg attcattctg ctaaccagta 7560 aggeaacccc gecagectag cegggteete aacgacagga geacgateat gegeacccqt 7620 ggccaggacc caacgctgcc cgagatgcgc cgcgtgcggc tgctggagat ggcggacgcg 7680 atggatatgt tctgccaagg gttggtttgc gcattcacag ttctccgcaa gaattgattg 7740 gctccaattc ttggagtggt gaatccgtta gcgaggtgcc gccggcttcc attcaggtcg 7800 : aggtggcccg gctccatgca ccgcgacgca acgcggggag gcagacaagg tatagggcgg 7860 . egectacaat ccatgecaac ccgttecatg tgetegecga ggeggcataa ategeegtga 7920 . cgatcagcgg tccagtgatc gaagttaggc tggtaagagc cgcgagcgat ccttgaagct 7980 gtccctgatg gtcgtcatct acctgcctgg acagcatggc ctgcaacgcg ggcatcccga 8040 tgccgccgga agcgagaaga atcataatgg ggaaggccat ccagcctcgc gtc .8093

<210> 79

<211> 8098

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
 pAN429 plasmid

<400> 79

ggcggccgct tcgtcgaccg aaacagcagt tataaggcat gaagctgtcc ggttttgca 60 aaagtggctg tgactgtaaa aagaaatcga aaaagaccgt tttgtgtgaa aacggtcttt 120 ttgtttcctt ttaaccaact gccataactc gaggcctacc tagcttccaa gaaagatatc 180 ctaacagcac aagagcggaa agatgttttg ttctacatcc agaacaacct ctgctaaaat 240 tcctgaaaaa ttttgcaaaa agttgttgac tttatctaca aggtgtggta taataatctt 300

aacaacagca ggacgctcta gattagaaag gaggtttaat taatgtatcg tacgatgatg 360 ageggeaaac tteacaggge aactgttaeg gaageaaace tgaactatgt gggaageatt 420 acaattgatg aagatotoat tgatgotgtg ggaatgotto otaatgaaaa agtacaaatt 480 gtgaataata ataatggagc acgtcttgaa acgtatatta ttcctggtaa acggggaagc 540 ggcgtcatat gcttaaacgg tgcagccgca cgccttgtgc aggaaggaga taaggtcatt 600 attatttcct acaaaatgat gtctgatcaa gaagcggcaa gccatgagcc gaaagtggct 660 gttctgaatg atcaaaacaa aattgaacaa atgctgggga acgaaccagc ccgtacaatt 720 ttgtaaagga tcctgttttg gcggatgaga gaagattttc agcctgatac agattaaatc 780 agaacgcaga agcggtctga taaaacagaa tttgcctggc ggcagtagcg cggtggtccc 840 acctgacccc atgccgaact cagaagtgaa acgccgtagc gccgatggta gtgtggggtc 900 tececatgeg agagtaggga actgecagge ateaaataaa acgaaagget cagtegaaag 960 actgggcctt tcgttttatc tgttgtttgt cggtgaacgc tctcctgagt aggacaaatc 1020 cyccgggagc ggatttgaac gttgcgaagc aacggcccgg agggtggcgg gcaggacgcc 1080 cgccataaac tgccaggcat caaattaagc agaaggccat cctgacggat ggcctttttg 1140 ; cgtttctaca aactcttggt acccagaaaa agcggcaaaa gcggctgtta aaaaagcgaa 1200 . ategaagaag etgtetgeeg etaagaegga atateaaaag egttetgetg ttgtgteate 1260 tttaaaagtc acagccgatg aatcccagca agatgtccta aaatacttga acacccagaa 1320 agataaagga aatgcagacc aaattcattc ttattatgtg gtgaacggga ttgctgttca 1380 tgcctcaaaa gaggttatgg aaaaagtggt gcagtttccc gaagtggaaa aggtgcttcc 1440 taatgagaaa cggcagcttt ttaagtcatc ctccccattt aatatgaaaa aagcacagaa 1500 agctattaaa gcaactgacg gtgtggaatg gaatgtagac caaatcgatg ccccaaaagc 1560 ttgggcactt ggatatgatg gaactggcac ggttgttgcg tccattgata ccggggtgga 1620 atggaatcat ccggcattaa aagagaaata tcgcggatat aatccggaaa atcctaatga 1680 gcctgaaaat gaaatgaact ggtatgatgc cgtagcaggc gaggcaagcc cttatgatga 1740 tttggctcat ggaacccacg tgacaggcac gatggtgggc tctgaacctg atggaacaaa 1800 tcaaatcggt gtagcacctg gcgcaaaatg gattgctgtt aaagcgttct ctgaagatgg 1860 cggcactgat gctgacattt tggaagctgg tgaatgggtt ttagcaccaa aggacgcgga 1920 aggaaatccc cacccggaaa tggctcctga tgttgtcaat aactcatggg gagggggctc 1980 tggacttgat gaatggtaca gagacatggt caatgcctgg cgttcggccg atattttccc 2040

tgagttttca gcggggaata cggatctctt tattcccggc gggcctggtt ctatcgcaaá 2100 teeggcaaac tatecagaat egtttgcaac tggagegaet gagaatteea atteeceatg 2160 gagagaaaag aaaatcgcta atgttgatta ctttgaactt ctgcatattc ttgaatttaa 2220 aaaggctgaa agagtaaaag attgtgctga aatattagag tataaacaaa atcgtgaaac 2280 aggegaaaga aagttgtate gagtgtggtt ttgtaaatee aggetttgte caatgtgeaa 2340 ctggaggaga gcaatgaaac atggcattca gtcacaaaag gttgttgctg aagttattaa 2400 acaaaagcca acagttegtt ggttgtttet cacattaaca gttaaaaatg tttatgatgg 2460 cgaagaatta aataagagtt tgtcagatat ggctcaagga tttcgccgaa tgatgcaata 2520 taaaaaaatt aataaaaatc ttgttggttt tatgcgtgca acggaagtga caataaataa 2580 taaagataat tottataato agoacatgoa tgtattggta tgtgtggaac caacttattt 2640 taagaataca gaaaactacg tgaatcaaaa acaatggatt caattttgga aaaaggcaat 2700 gaaattagac tatgatccaa atgtaaaagt tcaaatgatt cgaccgaaaa ataaatataa 2760 atcggatata caatcggcaa ttgacgaaac tgcaaaatat cctgtaaagg atacggattt 2820 tatgaccgat gatgaagaaa agaatttgaa acgtttgtct gatttggagg aaggtttaca 2880; ccgtaaaagg ttaatctcct atggtggttt gttaaaagaa atacataaaa aattaaacct 2940 tgatgacaca gaagaaggcg atttgattca tacagatgat gacgaaaaag ccgatgaaga 3000. tggattttct attattgcaa tgtggaattg ggaacggaaa aattatttta ttaaagagta 3060 gttcaacaaa cgggccatat tgttgtataa gtgatgaaat actgaattta aaacttagtt 3120 tatatgtggt aaaatgtttt aatcaagttt aggaggaatt aattatgaag tgtaatgaat 3180 gtaacagggt tcaattaaaa gagggaagcg tatcattaac cctataaact acgtctgccc 3240 tcattattgg agggtgaaat gtgaatacat cctattcaca atcgaattta cgacacaacc 3300 aaattttaat tiggctitgc attitatcit titttagcgt attaaatgaa atggtittga 3360 acgtctcatt acctgatatt gcaaatgatt ttaataaacc acctgcgagt acaaactggg 3420 tgaacacage ctttatgtta accttttcca ttggaacage tgtatatgga aagetatetg 3480 atcaattagg catcaaaagg ttactcctat ttggaattat aataaattgt ttcgggtcgg 3540 taattgggtt tgttggccat tetttetttt eettaettat tatggetegt tttatteaag 3600 gggctggtgc agctgcattt ccagcactcg taatggttgt agttgcgcgc tatattccaa 3660 aggaaaatag gggtaaagca tttggtctta ttggatcgat agtagccatg ggagaaggag 3720 teggtecage gattggtgga atgatagece attatattea ttggtectat ettetaetea 3780

ttcctatgat	aacaattatc	actgttccgt	ttcttatgaa	attattaaag	aaagaagtaa	3840
ggataaaagg	tcattttgat	atcaaaggaa	ttatactaat	gtctgtaggc	attgtatttt	3900
ttatgttgtt	tacaacatca	tatagcattt	cttttcttat	cgttagcgtg	ctgtcattcc	3960
tgatatttgt	aaaacatatc	aggaaagtaa	cagatccttt	tgttgatccc	ggattaggga	4020
aaaatatacc	ttttatgatt	ggagttcttt	gtgggggaat	tatatttgga	acagtagcag	4080
ggtttgtctc	tatggttcct	tatatgatga	aagatgttca	ccagctaagt	actgccgaaa	41.4.0
tcggaagtgt	aattatttc	cctggaacaa	tgagtgtcat	tattttcggc	tacattggtg	4200
ggatacttgt	tgatagaaga	ggtcctttat	acgtgttaaa	catcggagtt	acatttcttt	4260
ctgttagctt	tttaactgct	tcctttcttt	tagaaacaac	atcatggttc	atgacaatta	4320
taatcgtatt	tgttttaggt	gggctttcgt	tcaccaaaac	agttatatca	acaattgttt	4380
caagtagctt	gaaacagcag	gaagctggtg	ctggaatgag	tttgcttaac	tttaccagct	4440
ttttatcaga	gggaacaggt	attgcaattg	taggtggttt	attatccata	eccttacttg	4500
atcaaaggtt	gttacctatg	gaagttgatc	agtcaactta	tctgtatagt	aatttgttat	4560
tacttttttc	aggaatcatt	gtcattagtt	-ggctggttac	cttgaatgta	tataaacatt	4620
ctcaaaggga	tttctaaatc	gttaagggat	caactttggg	agagayttca	aaattgatcc	4680
ttcttttata	acagttcgaa	geggeegeaa	ttcttgaaga	cgaaagggcc	tegtgatacg	4740
cctattttta	taggttaatg	tcatgataat	aatggtttct	tagacgtcag	gtggcacttt	4800
tcggggaaat	gtgcgcggaa	cccctatttg	tttattttc	taaatacatt	. caaatatgta	4860
tccgctcatg	agacaataac	cctgataaat	gcttcaataa	tattgaaaaa	ggaagagtat	4920
gagtattcaa	catttccgtg	tegecettat	tecettett	gcggcatttt	geetteetgt	4980
ttttgctcac	: ccagaaacgc	tggtgaaagt	aaaagatgct	gaagatcagt	tgggtgcacg	5040
agtgggttac	: atcgaactgg	atctcaacag	g cggtaagatc	cttgagagtt	ttegeceega	5100
agaacgtttt	ccaatgatga	gcactttaa	agttctgcta	tgtggcgcgg	g tattateceg	5160
tattgacgcc	gggcaagagc	: aactcggtc	g ccgcatacac	: tattctcaga	atgacttggt	5220
tgagtactca	ccagtcacag	g aaaagcatct	tacggatggc	: atgacagtaa	a gagaattatg	5280
cagtgctgcc	ataaccatga	gtgataacad	tgcggccaac	ttacttctga	a caacgatcgg	5340
aggaccgaag	g gagetaaeeg	ctttttgc	a caacatgggg	gatcatgta:	a ctcgccttga	5400
tcgttgggaa	a ccggagctga	atgaagccat	accaaacgac	gagegtgae	a ccacgatgcc	54.60
tacaacaata	r gcaacaacgt	tococaaaci	attaactoo	gaactactt	a ctctagette	5520

ccggcaacaa ttaatagact ggatggaggc ggataaagtt gcaggaccac ttctgcgctc 5580 ggcccttccg gctggctggt trattgctga taaatctgga gccggtgagc gtgggtctcg 5640 cggtatcatt gcagcactgg ggccagatgg taagccctcc cgtatcgtag ttatctacac 5700 gacggggagt caggcaacta tggatgaacg aaatagacag atcgctgaga taggtgcctc 5760 actgattaag cattggtaac tgtcagacca agtttactca tatatacttt agattgattt 5820 aaaacttcat ttttaattta aaaggatcta ggtgaagatc ctttttgata atctcatgac 5880 caaaatccct taacgtgagt tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa 5940 aggatettet tgagateett tttttetgeg egtaatetge tgettgeaaa caaaaaaace 6000 accgctacca gcggtggttt gtttgccgga tcaagagcta ccaactcttt ttccgaaggt 6060 aactggette agcagagege agataceaaa tactgteett etagtgtage egtagttagg 6120 ccaccacttc aagaactctg tagcaccgcc tacatacctc gctctgctaa tcctgttacc 6180 agtggctgct gccagtggcg ataagtcgtg tcttaccggg ttggactcaa gacgatagtt 6240 accggataag gcgcagcggt cgggctgaac ggggggttcg tgcacacagc ccagcttgga 6300 . gegaaegaee tacaeegaae tgagataeet acagegtgag etatgagaaa gegeeaeget 6360 . 🕟 tecegaaggg agaaaggegg acaggtatee ggtaagegge agggteggaa caggagageg 6420 $_{\odot}$ cacgagggag cttccagggg gaaacgcctg gtatctttat agtcctgtcg ggtttcgcca 6480 🛒 cctctyactt gagcgtcgat ttttgtgatg ctcgtcaggg gggcggagcc tatggaaaaa 6540 egecageaac geggeetttt taeggtteet ggeettttge tggeettttg etcacatgtt 6600 ctttcctgcg ttatcccctg attctgtgga taaccgtatt accgcctttg agtgagctga 6660 tacegetege egeageegaa egaeegageg çagegagtea gtgagegagg aageggaaga 6720 gegeetgatg eggtatttte teettaegea tetgtgeggt attteacace geatatggtg 6780 cacteteagt acaatetget etgatgeege atagttaage eagtatacae teegetateg 6840 ctacgtgact gggtcatggc tgcgccccga cacccgccaa cacccgctga cgcgccctga 6900 egggettgte tgeteeegge ateegettae agacaagetg tgacegtete egggagetge 6960 atgtgtcaga ggttttcacc gtcatcaccg aaacgcgcga ggcagctgcg gtaaagctca 7020 tcagcgtggt cgtgaagcga ttcacagatg tctgcctgtt catccgcgtc cagctcgttg 7080 agtttctcca gaagcgttaa tgtctggctt ctgataaagc gggccatgtt aagggcggtt 7140 ttttcctgtt tggtcacttg atgcctccgt gtaaggggga atttctgttc atgggggtaa 7200 tgataccgat gaaacgagag aggatgctca cgatacgggt tactgatgat gaacatgccc 7260

agattactga acgttgtaag ggtaaacaac tggcggtatg gatgcggcgg gaccagagaa 7320
aaatcactca gggtcaatgc cagcgcttcg ttaatacaga tgtaggtgtt ecacagggta 7380
gccagcagca tcctgcgatg cagatccgga acataatggt gcagggcgct gacttccgcg 7440
tttccagact ttacgaaaca cggaaaccga agaccattca tgttgttgct caggtcgcag 7500
acgttttgca gcagcagtcg cttcacgttc gctcgcgtat cggtgattca ttctgctaac 7560
cagtaaggca accccgccag cctagccggg tcctcaacga caggagcacg atcatgcgca 7620
cccgtggcca ggacccaacg ctgcccgaga tgcgccgcgt gcggctgctg gagatggcgg 7680
acgcgatgga tatgttctgc caagggttgg tttgcgcatt cacagttctc cgcaagaatt 7740
gattggctcc aattcttgga gtggtgaatc cgttagcgag gtgccgccgg cttccattca 7800
ggtcgaggtg gcccggctcc atgcaccgc acgcaacgcg gggaggcaga caaggtatag 7860
ggcggcgcct acaatccatg ccaacccgtt ccatgtgctc gccgaggcgg cataaatcgc 7920
cgtgacgatc agcggtccag tgatcgaagt taggctgta agagccgcag gcgatccttg 7980
aagctgtccc tgatggtcgt catctacctg cctggacagc atggcctgca acgcggcat 8040
cccgatgccg ccggaagcga gaagaatcat aatggggaag gccatccagc ctcgctc 8098

<400> 80
ccatcgaatg gccagatgat taattcctaa tttttgttga cactctatca ttgatagagt 60
tattttacca ctccctatca gtgatagaga aaagtgaaat gaatagttcg acaaaaatct 120
agattagaaa ggaggattta aatatgagac agattactga tatttcacag ctgaaagaag 180
ccataaaaca ataccattca gagggcaagt caatcggatt tgttccgacg atggggttc 240
tgcatgaggg gcatttaacc ttagcagaca aagcaagaca agaaaacgac gccgttatta 300
tgagtattt tgtgaatcct gcacaattcg gccctaatga agattttgaa gcatatccgc 360
gcgatattga gcgggatgca gctcttgcag aaaacgccgg agtcgatatt ctttttacgc 420
cagatgctca tgatatgtat cccggtgaaa agaatgtcac gattcatgta gaaagacgca 480
cagacgtgtt atgcgggcgc tcaagagaag gacattttga cggggtcgcg atcgtactga 540
cgaagctttt caatctagtc aagccgactc gtgcctattt cggtttaaaa gatgcgcagc 600

<210> .80

<211> 4450

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
pAN443 plasmid

aggtagetgt tgttgatggg ttaatcageg acttetteat ggatattgaa ttggtteetg 660 tcgatacggt cagagaggaa gacggcttag ccaaaagctc tcgcaatgta tacttaacag 720 ctgaggaaag aaaagaagcg cctaagctgt atcgggccct tcaaacaagt gcggaacttg 780 tccaagccgg tgaaagagat cctgaagcgg tgataaaagc tgcaaaagat atcattgaaa 840 cgactagcgg aaccatagac tatgtagagc tttattccta tccggaactc gagcctgtga 900 atgaaattgc tggaaagatg attctcgctg ttgcagttgc tttttcaaaa gcgcgtttaa 960 tagataatat cattattgat attcgtagaa aggaggtgaa ttaatatgta tcgtacgatg 1020 atgageggea aaetteacag ggeaaetgtt aeggaageaa aeetgaaeta tgtgggaage 1080 attacaattg atgaagatct cattgatgct gtgggaatgc ttcctaatga aaaagtacaa 1140 attgtgaata ataataatgg agcacgtett gaaaegtata ttatteetgg taaaegggga 1200ageggegtea tatgettaaa eggtgeagee geaegeettg tgeaggaagg agataaggte 1260 attattattt cctacaaaat gatgtctgat caagaagcgg caagccatga gccgaaagtg 1320 gctgttctga atgatcaaaa caaaattgaa caaatgctgg ggaacgaacc agcccgtaca 1380 attttgtaaa ggatccccg gggatccctc gaggtcgacc tgcaggggga ccatggtctc 1440 agegettgga gecaceegea gttegaaaaa taataagett gaeetgtgaa gtgaaaaatg 1500 gegeacattg tgegaeattt tttttgtetg eegtttaeeg etaetgegte aeggatetee 1560 acgcgccctg tagcggcgca ttaagcgcgg cgggtgtggt ggttacgcgc agcgtgaccg 1620 ctacacttgc cagogocota gogocogoto ctttogottt cttcccttcc tttctcgcca 1680 cgttcgccgg ctttccccgt caagctctaa atcgggggct ccctttaggg ttccgattta 1740 gtgetttacg gcacctcgac cccaaaaaac ttgattaggg tgatggttca cgtagtgggc 1800 categeeetg atagaeggtt tttegeeett tgaegttgga gteeaegtte tttaatagtg 1860 gactettgtt ceaaactgga acaacactca accetatete ggtetattet tttgatttat 1920 aagggatttt geegattteg geetattggt taaaaaatga getgatttaa caaaaattta 1980 acgegaattt taacaaaata ttaacgetta caattteagg tggcaetttt eggggaaatg 2040 tgcgcggaac ccctatttgt ttatttttct aaatacattc aaatatgtat ccgctcatga 2100 gacaataacc ctgataaatg cttcaataat attgaaaaag gaagagtatg agtattcaac 2160 atttccgtgt cgcccttatt cccttttttg cggcattttg ccttcctgtt tttgctcacc 2220 cagaaacgct ggtgaaagta aaagatgctg aagatcagtt gggtgcacga gtgggttaca 2280 tegaactgga teteaacage ggtaagatee ttgagagttt tegeceegaa gaacgtttte 2340

caatgatgag cacttttaaa gttctgctat gtggcgcggt attatcccgt attgacgccg 2400 ggcaagagca actcggtcgc cgcatacact attctcagaa tgacttggtt gagtactcac 2460 cagtcacaga aaagcatctt acggatggca tgacagtaag agaattatgc agtgctgcca 2520 taaccatqaq tgataacact gcggccaact tacttctgac aacgatcgga ggaccgaagg 2580 agctaaccgc ttttttgcac aacatggggg atcatgtaac tcgccttgat cgttgggaac 2640 cggagctgaa tgaagccata ccaaacgacg agcgtgacac cacgatgcct gtagcaatgg 2700 caacaacgtt gcgcaaacta ttaactggcg aactacttac tctagcttcc cggcaacaat 2760 tgatagactg gatggaggcg gataaagttg caggaccact tctgcgctcg gcccttccgg 2820 ctggctggtt tattgctgat aaatctggag ccggtgagcg tggctctcgc ggtatcattg 2880 cagcactggg gccagatggt aagccctccc gtatcgtagt tatctacacg acggggagtc 2940 aggcaactat ggatgaacga aatagacaga tegetgagat aggtgeetea etgattaage 3000 attggtagga attaatgatg tctcgtttag ataaaagtaa agtgattaac agcgcattag 3060. agotgottaa tgaggtogga atogaaggtt taacaacoog taaactogoo cagaagotag 3120 🕢 gtqtagagca gcctacattg tattggcatg taaaaaataa gcgggctttg ctcgacgcct 3180 . tagccattga gatgttagat aggcaccata ctcacttttg ccctttagaa ggggaaagct 3240 ggcaagattt tttacgtaat aacgctaaaa gttttagatg tgctttacta agtcatcgcg 3300 . 🐇 atggagcaaa agtacattta ggtacacggc ctacagaaaa acagtatgaa actctcgaaa 3360 atcaattagc ctttttatgc caacaaggtt tttcactaga gaatgcatta tatgcactca 3420 gcgcagtggg gcattttact ttaggttgcg tattggaaga tcaagagcat caagtcgcta 3480 aagaagaaag ggaaacacct actactgata gtatgccgcc attattacga caagctatcg 3540 aattatttga tcaccaaggt gcagagccag ccttcttatt cggccttgaa ttgatcatat 3600 gcggattaga aaaacaactt aaatgtgaaa gtgggtetta aaageageat aacettttte 3660 cgtgatggta acttcactag tttaaaagga tctaggtgaa gatccttttt gataatctca 3720 tgaccaaaat cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga 3780 tcaaaggatc ttcttgagat ccttttttc tgcgcgtaat ctgctgcttg caaacaaaaa 3840 aaccaccgct accageggtg gtttgtttgc eggatcaaga getaccaact ettttteega 3900 aggtaactgg cttcagcaga gcgcagatac caaatactgt ccttctagtg tagccgtagt 3960 taggccacca cttcaagaac tctgtagcac cgcctacata cctcgctctg ctaatcctgt 4020 taccagtggc tgctgccagt ggcgataagt cgtgtcttac cgggttggac tcaagacgat 4080

PCT/US00/25993

agttacegga taaggegeag eggtegget gaacgggggg ttegtgeaca eageceaget 4140 tggagegaac gacetacace gaactgagat acetacageg tgagetatga gaaagegeea 4200 egetteeega agggagaaag geggacaggt ateeggtaag eggeagggte ggaacaggag 4260 agegeacgag ggagetteea gggggaaacg eetggtatet ttatagteet gtegggtte 4320 gecacetetg acttgagegt egatttttgt gatgetegte aggggggggg ageetatgga 4380 aaaacgeeag caacgeggee tttttaeggt teetggeett ttgetggeet tttgeteaca 4440 tgaecegaca

<210> 81

<211> 10212

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant
pAN251 plasmid

<400> 81 geggeegeta aaaagagett gaggatttge ggagtgaaaa teagacattg eggaateage 60 🐇 tagagatgac agaagaggat tacaaggcac tgatcgatat catggatcgg gccagaaaaa 120. tggttgtttc gaaggaagac ggaagaatga aaaaagcggc tcaagaaacg taaagaaacg 180: cctgaaatga accggcccta tagtaagaat aggccggttg ttttgatttc tatgcagact 240'. ctcccggtgt catttcgcga tccatatcag gatgccagat gagcgggtct tcccctttgt 300. cccgcqccat atcatactta acagttttaa agttcatttt gttccaaaat tccgctgatt 360 tcattctcgg atttgtccgg atcggcattt tgaatgattt tgcaaattca accaaggctc 420 toccgtatec cetgttetgg tageceggaa gaaceteaag ettecacage tecaaataat 480 cctqqcggtt qtcaaaataq ggattcgatt tgccgttaac ttgatacaga ctcattcgtg 540 ctacaagttt atcqccaaaa taaatcccgt aaaaaggcga ggtgctgtca ttttcaataa 600 tattatectq aaqttettea ageattgaaa geteetgaat geegtattet ttgaatttet 660 tgaattette caqeqtttta tagttgataa geagaegtte taeetttgte aaacaaatet 720 ccccctttgt tgtttctaca tatattgtaa acgctttatt taaaaaaatcc aaatatttaa 780 actttaattt taagcacatg ggatctttga gaagtaattt cttcttactt ctgctatgat 840 aatacqtaaa tqcqtcqacc qaaacagcag ttataaggca tgaagctgtc cggtttttgc 900 aaaaqtggct qtgactgtaa aaagaaatcg aaaaagaccg ttttgtgtga aaacggtctt 960

tttgtttcct tttaaccaac tgccataact cgaggcctac ctagcttcca agaaagatat 1020 cctaacagca caagagcgga aagatgtttt gttctacatc cagaacaacc tctgctaaaa 1080 ttcctgaaaa attttgcaaa aagttgttga ctttatctac aaggtgtggt ataataatct 1140 taacaacagc aggacgctct agaggaggag acatcatgaa aattggaatt atcggcggag 1200 gctccgttgg tcttttatgc gcctattatt tgtcacttta tcacgacgtg actgttgtga 1260 cgaggcggca agaacaggct gcggccattc agtctgaagg aatccggctt tataaaggcg 1320 gggaggaatt cagggctgat tgcagtgcgg acacgagtat caattcggac tttgacctgc 1380 ttgtcgtgac agtgaagcag catcagcttc aatctgtttt ttcgtcgctt gaacgaatcg 1440 ggaagacgaa tatattattt ttgcaaaacg gcatggggca tatccacgac ctaaaagact 1500 ggcacgttgg ccattccatt tatgttggaa tcgttgagca cggagctgta agaaaatcgg 1560 atacagetgt tgatcataca ggcctaggtg cgataaaatg gagcgcgttc gacgatgctg 1620 aaccagaccg gctgaacatc ttgtttcagc ataaccattc ggattttccg atttattatg 1680 agacggattg gtaccgtctg ctgacgggca agctgattgt aaatgcgtgt attaatcctt 1740 taactgcgtt attgcaagtg aaaaatggag aactgctgac aacgccagct tatctggctt 1800 ttatgaaget ggtattteag gaggeatgee geattttaaa aettgaaaat gaagaaaagg 1860 cttgggageg ggttcaggec gtttgtgggc aaacgaaaga gaategttca tcaatgetgg 1920 ttgacgtcat tggaggccgg cagacggaag ctgacgccat tatcggatac ttattgaagg 1980 aagcaagtet teaaggtett gatgeegtee acetagagtt titatatgge agcateaaag 2040 cattggagcg aaatacaaac aaagtetttt gagettttte ggtaacatge tatactcatt 2100 toggatoact aactatttat tggagaaagg aagttotaga agatgoagot aactgaactt 2160 tccatcaaaa atcagaatgt gtttgtacag cactatatag atggcaaaga agaaatgtct 2220 tetttttttg attacagtat teateataag gaeatgtgge gegaaagaet ggaagaetta 2280 tcttcccggt ttttcgcaag agaggaattg gcggcgtact taacctctta ccataataaa 2340 ttcggttcaa gtgcgatgca gtctgctatt gagaagctga aggacccgtc aagtgccgct 2400 gtagtcggcg gacagcaggc aggactttta acaggaccgc tttacaccat acataaaatc 2460 atttcaatca ttgttttagc aaagcaacaa gaaaaggaac tgcaagtgcc tgtcatacca 2520 atcttctggg tggctggaga agaccacgat ttggatgaga ttaattttgt tcacacatct 2580 gaagagaatg ggcctgtgaa aaaaaagctg cctcagtctt attggaagaa atcatcagca 2640 gcgagtacat cgcttgatca ggaaaagtgt gccgcgtgga tagatgatgt ttttgccgct 2700

. . --

tttgaagaaa cagaccatac gaatacactt ctcgacaatg tgaaacgatg tttaagggaå 2760 tctgttacgt ttactgactt ctttgaactg ctgatcgcgg atttgttcca agaagagggc 2820 ttagttttat taaattctgg ggatcctgtt ttggcggatg agagaagatt ttcagcctga 2880 tacagattaa atcagaacgc agaagcggtc tgataaaaca gaatttgcct ggcggcagta 2940 gegeggtggt eccaeetgae eccatgeega acteagaagt gaaaegeegt agegeegatg 3000 gtagtgtggg gtctccccat gcgagagtag ggaactgcca ggcatcaaat aaaacgaaag 3060 gctcagtcga aagactgggc ctttcgtttt atctgttgtt tgtcggtgaa cgctctcctg 3120 agtaggacaa atccgccggg agcggatttg aacgttgcga agcaacggcc cggagggtgg 3180 cgggcaggac gcccgccata aactgccagg catcaaatta agcagaaggc catcctgacg 3240 gatggccttt ttgcgtttct acaaactctt ggtacccaga aaaagcggca aaagcggctg 3300 ttaaaaaagc gaaatcgaag aagctgtctg ccgctaagac ggaatatcaa aagcgttctg 3360 ctgttgtgtc atctttaaaa gtcacagccg atgaatccca gcaagatgtc ctaaaatact 3420. tgaacaccca gaaagataaa ggaaatgcag accaaattca ttcttattat gtggtgaacg 3480. ggattgctgt tcatgcctca aaagaggtta tggaaaaagt ggtgcagttt cccgaagtgg 3540 aaaaggtgct tootaatgag aaacggcage tttttaagtc atootcocca tttaatatga 3600 🖖 aaaaagcaca gaaagctatt aaagcaactg acggtgtgga atggaatgta gaccaaatcg 3660 atgccccaaa agcttgggca cttggatatg atggaactgg cacggttgtt gcgtccattg 3720 ataccggggt ggaatggaat catccggcat taaaagagaa atatcgcgga tataatccgg 3780 aaaateetaa tgageetgaa aatgaaatga aetggtatga tgeegtagea ggegaggeaa 3840 gcccttatga tgatttggct catggaaccc acgtgacagg cacgatggtg ggctctgaac 3900 ctgatggaac aaatcaaatc ggtgtagcac ctggcgcaaa atggattgct gttaaagcgt 3960 tetetgaaga tggeggeact gatgetgaca ttttggaage tggtgaatgg gttttageac 4020 caaaggacgc ggaaggaaat ccccacccgg aaatggctcc tgatgttgtc aataactcat 4080 ggggaggggg ctctggactt gatgaatggt acagagacat ggtcaatgcc tggcgttcgg 4140 ccgatatttt ccctgagttt tcagcgggga atacggatct ctttattccc ggcgggcctg 4200 gttctatcgc aaatccggca aactatccag aatcgtttgc aactggagcg actgagaatt 4260 ccaattcccc atggagagaa aagaaaatcg ctaatgttga ttactttgaa cttctgcata 4320 ttettgaatt taaaaagget gaaagagtaa aagattgtge tgaaatatta gagtataaae 4380 aaaatcgtga aacaggcgaa agaaagttgt atcgagtgtg gttttgtaaa tccaggcttt 4440

gtccaatgtg caactggagg agagcaatga aacatggcat tcagtcacaa aaggttgttg 4500 ctgaagttat taaacaaaag ccaacagttc gttggttgtt tctcacatta acagttaaaa 4560 atgtttatga tggcgaagaa ttaaataaga gtttgtcaga tatggctcaa ggatttcgcc 4620 gaatgatgca atataaaaaa attaataaaa atcttgttgg ttttatgcgt gcaacggaag 4680 tgacaataaa taataaagat aattottata atcagcacat gcatgtattg gtatgtgtgg 4740 aaccaactta ttttaagaat acagaaaact acgtgaatca aaaacaatgg attcaatttt 4800 ggaaaaaggc aatgaaatta gactatgatc caaatgtaaa agttcaaatg attcgaccga 4860 aaaataaata taaatcggat atacaatcgg caattgacga aactgcaaaa tatcctgtaa 4920 aggatacgga ttttatgacc gatgatgaag aaaagaattt gaaacgtttg tctgatttgg 4980 aggaaggttt acaccgtaaa aggttaatct cctatggtgg tttgttaaaa gaaatacata 5040 aaaaattaaa ccttgatgac acagaagaag gcgatttgat tcatacagat gatgacgaaa 5100 aagccgatga agatggattt tctattattg caatgtggaa ttgggaacgg aaaaattatt 5160 ttattaaaga gtagttcaac aaacgggcca tattgttgta taagtgatga aatactgaat 5220 ttaaaactta gtitatatgt ggtaaaatgt tttaatcaag tttaggagga attaattatg 5280 aagtgtaatg aatgtaacag ggttcaatta aaagagggaa gcgtatcatt aaccctataa 5340 actacgtctg ccctcattat tggagggtga aatgtgaata catcctattc acaatcgaat 5400 ttacgacaca accaaatttt aatttggctt tgcattttat ctttttttag cgtattaaat 5460 gaaatggttt tgaacgtctc attacctgat attgcaaatg attttaataa accacctgcg 5520 agtacaaact gggtgaacac ageetttatg ttaacetttt ccattggaac agetgtatat 5580 ggaaagctat ctgatcaatt aggcatcaaa aggttactcc tatttggaat tataataaat 5640 tgtttcgggt cggtaattgg gtttgttggc cattctttct tttccttact tattatggct 5700 cgttttattc aaggggctgg tgcagctgca tttccagcac tcgtaatggt tgtagttgcg 5760 cgctatattc caaaggaaaa taggggtaaa gcatttggtc ttattggatc gatagtagcc 5820 atgggagaag gagtcggtcc agcgattggt ggaatgatag cccattatat tcattggtcc 5880 tatcttctac tcattcctat gataacaatt atcactgttc cgtttcttat gaaattatta 5940 aagaaagaag taaggataaa aggtcatttt gatatcaaag gaattatact aatgtctgta 6000 gtgctgtcat tcctgatatt tgtaaaacat atcaggaaag taacagatcc ttttgttgat 6120 cccggattag ggaaaaatat accttttatg attggagttc tttgtggggg aattatattt 6180

ggaacagtag cagggtttgt ctctatggtt ccttatatga tgaaagatgt tcaccagcta 6240 agtactgccg aaatcggaag tgtaattatt ttccctggaa caatgagtgt cattattttc 6300 ggctacattg gtgggatact tgttgataga agaggtcctt tatacgtgtt aaacatcgga 6360 gttacatttc tttctgttag ctttttaact gcttcctttc ttttagaaac aacatcatqq 6420 ttcatgacaa ttataatcgt atttgtttta ggtgggcttt cgttcaccaa' aacagttata 6480 tcaacaattg tttcaagtag cttgaaacag caggaagctg gtgctggaat gagtttgctt 6540 aactttacca gctttttatc agagggaaca ggtattgcaa ttgtaggtgg tttattatcc 6600 ataccettae ttgateaaag gttgttacet atggaagttg ateagteaac ttatetgtat 6660 agtaatttgt tattactttt ttcaggaatc attgtcatta gttggctggt taccttgaat 6720 gtatataaac attctcaaag ggatttctaa atcgttaagg gatcaacttt gggagagagt 6780 tcaaaattga tcctttttt ataacagttc gaagcggccg caattcttga agacgaaagg 6840 gcctcgtgat acgcctattt ttataggtta atgtcatgat aataatggtt tcttagacgt 6900 caggtggcac ttttcgggga aatgtgcgcg gaacccctat ttgtttattt ttctaaatac 6960 attcaaatat gtatccgctc atgagacaat aaccctgata aatgcttcaa taatattgaa 7020 aaaggaagag tatgagtatt caacatttcc gtgtcgccct tattcccttt tttgcggcat 7080 tttgccttcc tgtttttgct cacccagaaa cgctggtgaa agtaaaagat gctgaaqatc 7140. agttgggtgc acgagtgggt tacatcgaac tggatctcaa cagcggtaag atccttgaga 7200 . gttttcgccc cgaagaacgt tttccaatga tgagcacttt taaagttctg ctatgtggcg 7260 cggtattatc ccgtattgac gccgggcaag agcaactcgg tcgccgcata cactattctc 7320 agaatgactt ggttgagtac tcaccagtca cagaaaagca tcttacggat ggcatgacag 7380 taagagaatt atgcagtgct gccataacca tgagtgataa cactgcggcc aacttacttc 7440 tgacaacgat cggaggaccg aaggagctaa ccgctttttt gcacaacatg ggggatcatg 7500 taactcgcct tgatcgttgg gaaccggagc tgaatgaagc cataccaaac gacgagcgtg 7560 acaccacgat gcctgcagca atggcaacaa cgttgcgcaa actattaact ggcgaactac 7620 ttactctagc ttcccggcaa caattaatag actggatgga ggcggataaa gttgcaggac 7680 cacttetgeg eteggeeett eeggetget ggtttattge tgataaatet ggageeggtg 7740 agcgtgggtc tcgcggtatc attgcagcac tggggccaga tggtaagccc tcccgtatcg 7800 tagttatcta cacgacgggg agtcaggcaa ctatggatga acgaaataga cagatcgctg 7860 agataggtgc ctcactgatt aagcattggt aactgtcaga ccaagtttac tcatatatac 7920

t	ttagattga	tttaaaactt	catttttaat	ttaaaaggat	ctaggtgaag,	atcctttttg	7980
a	taatctcat	gaccaaaatc	ccttaacgtg	agttttcgtt	ccactgageg	tcagaccccg	8040
t	agaaaagat	caaaggatct	tcttgagatc	cttttttct	gcgcgtaatc	tgctgcttgc	61:00
a	aacaaaaa	accaccgcta	ccagcggtgg	tttgtttgcc	ggatcaag <u>a</u> g	ctaccaactc	81.60
t	ttttccgaa	ggtaactggc	ttcagcagag	cgcagatacc	aaatactgtc	cttctagtgt	8220
a	gccgtagtt	aggccaccac	ttcaagaact	ctgtagcacc	gcctacatac	ctcgctctgc	8280
t	aatcctgtt	accagtggct	gctgccagtg	gcgataagtc	gtgtcttacc	gggttggact	8340
C	aagacgata	gttaccggat	aaggcgcagc	ggtcgggctg	aacggggggt	tegtgcacac	8400
а	gcccagctt	ggagcgaacg	acctacaccg	aactgagata	cctacagcgt	gagctatgag	8460
a	aagcgccac	gcttcccgaa	gggagaaagg	cggacaggta	teeggtaage	ggcagggtcg	8520
ç	gaacaggaga	gcgcacgagg	gagettecag	ggggaaacgc	ctggtatctt	tatagtcctg	8580
t	cgggtttcg	ccacctctga	cttgagcgtc	gatttttgtg	atgctcgtca	ggggggcgga	8640
ç	gcctatggaa	aaacgccagc	aacgcggcct	ttttacggtt	cctggccttt	tgctggcctt	8700
t	tgctcacat	gttctttcct	gcgttatccc	ctgattctgt	ggataaccgt	attaccgcct	87.60
ŧ	tgagtgagc	tgataccgct	cgccgcagcc	gaacgaccga	gcgcag c gag	tcagtgagcg	8820
3	aggaagcgga	agagcgcctg	atgcggtatt	ttctccttac	gcatctgtgc	ggtatttcac	6889
ě	accgcatatg	gtgcactctc	agtacaatct	gctctgatgc	cgcatagtta	agccagtata	8940
•	cactccgcta	tegetacgtg	actgggtcat	ggctgcgccc	cgacacccgc	caacacccgc	9000
4	tgacgcgccc	tgacgggctt	gtctgctccc	ggcatecgct	tacagacaag	g ctgtgaccgt	9060
,	ctccgggagc	: tgcatgtgtc	agaggttttc	accgtcatca	ccgaaacgcg	-cgaggcagct	9120
	gcggtaaagc	: tcatcagcgt	ggtcgtgaag	cgattcacag	atgtctgcct	gttcatccgc	9180
	gtccagctcg	, ttgagtttct	ccagaagcgt	taatgtctgg	, cttctgataa	a agegggeeat	9.240
	gttaagggco	gtttttcct	gtttggtcac	: ttgatgcctc	cgtgtaagg	g ggaatttctg	9300
	ttcatggggg	g taatgatacc	: gatgaaac g a	gagaggatgo	tcacgatacq	g ggttactgat	9360
	gatgaacato	g ceeggttact	ggaacgttgt	gagggtaaac	aactggcgg(atggatgcgg	9420
	cgggaccaga	a gaaaaatcac	: tcagggtcaa	tgccagcgct	tcgttaata	c agatgtaggt	9480
	gttccacag	g gtagccagca	gcatectge	g atgcagated	ggaacataa	t ggtgcagggc	9540
	gctgacttco	c gcgtttccag	g actttacgaa	acacggaaad	c cgaagacca	t tcatgttgtt	9600
	acteaggtee	cagacgtttt	gcagcagcag	tegetteace	tteacteac	tatcggtgat	9660

WO 01/021772

tcattctgct aaccagtaag gcaaccccgc cagcctagcc gggtcctcaa cgacaggagc 9720
acgatcatgc gcacccgtgg ccaggaccca acgctgcccg agatgcgccg cgtgcggctg 9780
ctggagatgg cggacgcgat ggatatgttc tgccaagggt tggtttgcgc attcacagtt 9840
ctccgcaaga attgattggc tccaattctt ggagtggtga atccgttagc gaggtgccgc 9900
cggcttccat tcaggtcgag gtggcccggc tccatgcacc gcgacgcaac gcggggaggc 9960
agacaaggta tagggcggcg cctacaatcc atgccaaccc gttccatgtg ctcgccgagg 10020
cggcataaat cgccgtgacg atcagggtc cagtgatcga agttaggctg gtaagagccg 10080
cgagcgatcc ttgaagctgt ccctgatggt cgtcatctac ctgcctggac agcatggcct 10140
gcaacgcggg catcccgatg ccgccggaag cgagaagaat cataatggg aaggccatcc 10200
agcctcgcgt cg

<210> 82

<211> 10426

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN267 plasmid

aacaaaattc tccagtctc acatcggttt gaaaggagga agcggaagaa tgaagtaaga 60
gggatttttg actccgaagt aagtcttcaa aaaatcaaat aaggagtgtc aagaatgttt 120
gcaaaacgat tcaaaacctc tttactgccg ttattcgctg gattttatt gctgttcat 180
ttggttctgg caggaccggc ggctgcgagt gctgaaacgg cgaacaaatc gaatgagctt 240
acagcaccgt cgatcaaaag cggaaccatt cttcatgcat ggaattggtc gttcaatacg 300
ttaaaaccaa atatgaagga tattcatgat gcaggatata cagccattca gacatctccg 360
attaaccaag taaaggaagg gaatcaagga gataaaagca tgtcgaactg gtactggctg 420
tatcagccga catcgtatca aattggcaac cgttacttag gtactgaaca agaatttaaa 480
gaaatgtgtg cagccgctga agaatatggc ataaaggtca ttgttgacgc ggtcatcaat 540
cataccacca gtgattatgc cgcgattcc aatgaggta agagtatcc aaactggaca 600
catggaaaca cacaaattaa aaactggtct gatcgaaata gtacataatg gatttcctta 660
cgcgaaatac gggcagacat ggcctgcccg gttattatta tttttgacac cagaccaact 720
ggtaatggta gcgaccggcg ctcaggatcg tctcggtacc aagagtttgt agaaacgcaa 780
aaaggccatc cgtcaggatg gccttctgct taatttgatg cctggcagtt tatggcggge 840

gtcctgcccg ccaccctccg ggccgttgct tcgcaacgtt caaatccgct cccggcggat 900 ttgtcctact caggagageg ttcaccgaca aacaacagat aaaacgaaag goccagtctt 960 togactgago ctttcgtttt atttgatgcc tggcagttcc ctactctcgc atggggagac 1020 cccacactac catcggcgct acggcgtttc acttctgagt tcggcatggg gtcaggtggg 1080 accaccgcgc tactgccgcc aggcaaattc tgttttatca gaccgcttct gcgttctgat 1140 ttaatctgta tcaggctgaa aatcttctct catccgccaa aacaggatcc atcacgaagc 1200 gtcgtatcga aaaaattaat tttgcgcaac ggagaccacc gcttccttct tcttgccttg 1260 tttcacaaac ggcatcattt cacgaagett tetteecaet aettegattt gatgttegtt 1320 ctcgcttgca ttgatagcgt tgaaacgagg acggtttact tggtttttga cgatccactc 1380 ttttgcgaat gtaccgtttt ggatatettt taataettet ttcatagatt ettttaettt 1440 agegtecaca aegegaggge etgatacgaa atetececae tgtgetgtgt eagagattga 1500 atatctcatt cctgcaagtc cttcttcgta cataaggtct acgattaatt tcagctcatg 1560 aagacactcg aagtatgcaa gttcaggctg ataacctgct tcagttaagg tttcaaatcc 1620 ggetttgaca agegegetta ateegeegea aagaactget tgeteacega acaaatetgt 1680 ttctgtttct tctttaaatg tcgtttctaa tacgcccgct cttgcgccgc cgattccttt 1740 agcataagcg agggettigt etettgette tecagteaca tettgataga tigegaacaa 1800 tgcaggtacg ccagctcctt gctcatatgt tcttcttacc aagtgtcccg ggcctttagg 1860 ggccactaag aatacateta cateegeegg aggaacaatt tgatggaaat geacgttaaa 1920 tccatgagcg aatactaatg attttcctgc tgtcaattca tctttgattt cagcttcgta 1980 tactttttgc tgctgctcat ccggaagcag aaccatgatg atttcggctt gggctgccgc 2040 ttcttttact gaaaatactt tatgtccgtc ttcttgggct tgagtgaaag attttccttg 2100 tctaacaccg acgatcacgt ctactccgct ttctttaagg ttcagggcat gtgcgtggcc 2160 ttgcgaaccg tacccgataa ccgctactgt ttttccagcc aatacgttct ctttgatatc 2220 accgttataa tatactttta ccatttcaat ctctcccttg ttatgtttta tacaatagat 2280 attgttttat tggatgacgc cttttgctgg ttcccctcgc aaaagccgtt gtacctgttc 2340 togogattto tttaatgoca taaggtttta ataactcaat aagogottoa attttgttag 2400 attcacctgt cacctgaaca acgatgctgt ctctgctgac atcaacgaca gaggctctaa 2460 acggttctat gattccatta atctctgttc ttgttgaagg tgcggagaca accttgatta 2520 aggecagete cetetggaca ategattgat ttgtgatgte tgtgaettte ageacateaa 2580

WO 01/021772

tctgtttgtt gagctgtttc gttaactgtt caacatcatt ttcaccttca acatgaacga 2640 cgaaggtgat tctggaaacg ccggctgttt ctgtgtgtcc aactgtaatg ctttcaatgt 2700 tgtaatgeet tittgtgaat agaceggtga teeggtttaa eaceeeggag eggtteacea 2760 cagtcaatgt gataattett tteaaggttt caececeace attteatgea geeettteee 2820 cggagccacc atcgggaata ctttttcttc gctggcaacc cgcacgtcaa tgacaacagg 2880 ttetettgat gttaatgeet etteeagett tteetttget teegettetg atgaaattet 2940 gatgccttta atgccgtatg cttcggacaa tttgacgaag tcaggctgag aagcgaattt 3000 agattetgaa taaegttett catagaaaat tteetgeeae tgtetgaeea tteegagaea 3060 agogttatit aaaatcacta oottgacogg aagatttaat togogaataa catogagtto 3120 ttgaagcgtc atttggaatc cgccgtctcc gacaaccgcg acaacagtag catcttttc 3180 ggccagctgt gcgccgatcg ccgccggaag accgaatccc atcgttccaa gtccgcctga 3240 cgtgacccat ttatctgctt tttggaacgg ataaaattgc gctgaccaca tttgatgctg 3300° gcctacatcc gttgcgacaa tggcctctcc ttttgtaaat tgatgaatat attcaatcaa 3360 tttctgaggt ttaaaacctt cttcttcatt atctacatac cagagcggat actcttcttt 3420 ccattctgcg agctgttttt tccattcgct tgaatcgctt tgtttgccgt cttgtttgat 3480 cageteetge aggacaattt tgetgtetee gactacagga atetgtgttt teatgatttt 3540 teegatttea getggateaa tategatgtg ggetatettt gegtttetgg caaagtgttt .3600 caggittcct gigacacggi catcaaaacg ggcgccgata cigattaata gatcacatic 3660 atgaagggcc atattggctg tataagtacc gtgcattccc gccatcccta ggaaaagcgg 3720 atggtcagcc gggaagcctc cgagccccaa aagggtgtgt gccacaggga tttgctgctg 3780 ttcagcataa ttttttaatt cttctgacgc ttttccgtgc agtacgcccg cacccgccag 3840 gatcaceggt tttttegeac tgeteaegge ttecacaage ttgeggatet geaaataatt 3900 cggctctgtt gtcggctggt atcccgggag attcatctca tgatcgtagc tgaattctcc 3960 ttcaattgtt gctacatctt tcggaatatc aatcaataca ggtccgggtc ttccagttgt 4020 tgcaatatgg aacgcttett taatgatgeg eggeagatet teeggetgge gaacetggta 4080 gctgtgtttt gttactggca tcgtaatccc taaaatgtct gcttcctgaa atgcatcgct 4140 cccgattaca gaggttgcta cctgccctgt aaagacgact aacggcaatg aatcaatcat 4200 ggcatcagca aggcctgtaa caaggtttgt cgctcccggc ctgacgtggc aatgacgaca 4260 ccggtttccg gagaccettg cgtatccctc cgctgcatga attgctcctt gttcgtgacg 4320

ggaaggatat gtaccaacct gaatgtatag cttatcgtaa atcggaagca cagccccgcc 4380 cggataaccg aagatcattt ctactttctc tttttttaat gattcaatca gcattaatcg 4440 tecgeteate gtetgtgtae atteggeaga tgetgaatee acetgtaeat tagteeecat 4500 tttatctcct cctctagagc gtcctgctgt tgttaagatt attataccac accttgtaga 4560 taaagtcaac aacttttgc aaaatttttc aggaatttta gcagaggttg ttctggatgt 4620 agaacaaaac atctttccgc tcttgtgctg ttaggatatc tttcttggaa gctaggtagg 4680 cctcgagtta tggcagttgg ttaaaaggaa acaaaaagac cgttttcaca caaaacggtc 4740 tttttcgatt tctttttaca gtcacagcca cttttgcaaa aaccggacag cttcatgcct 4800 tataactgct gtttcggtcg acgaagcggc cgccgtttaa acgaattcct gcagctggcg 4860 aatggcgatt ttcgttcgtg aatacatgtt ataataacta taactaataa cgtaacgtga 4920 ctggcaagag atattttaa aacaatgaat aggtttacac ttactttagt tttatggaaa 4980 tgaaagatca tatcatatat aatctagaat aaaattaact aasataatta ttatctagat 5040 aaaaaattta gaagccaatg aaatctataa ataaactaaa ttaagtttat ttaattaaca 5100 . actatggata taaaataggt actaatcaaa atagtgagga ggatatattt gaatacatac 5160 gaacaaatta ataaagtgaa aaaaatactt cggaaacatt taaaaaataa ccttattggt 5220 acttacatgt ttggatcagg agttgagagt ggactaaaac caaatagtga tcttgacttt 5280: ttagtcgtcg tatctgaacc attgacagat caaagtaaag aaatacttat acaaaaaatt 5340, agacctattt caaaaaaaat aggagataaa agcaacttac gatatattga attaacaatt 5400 attattcage aagaaatggt accgtggaat catecteeca aacaagaatt tatttatgga 5460 gaatggttac aagagcttta tgaacaagga tacattcctc agaaggaatt aaattcagat 5520 ttaaccataa tgctttacca agcaaaacga aaaaataaaa gaatatacgg aaattatgac 5580 ttagaggaat tactacctga tattccattt tctgatgtga gaagagccat tatggattcg 5640 tcagaggaat taatagataa ttatcaggat gatgaaacca actctatatt aactttatgc 5700 cgtatgattt taactatgga cacgggtaaa atcataccaa aagatattgc gggaaatgca 5760 gtggctgaat cttctccatt agaacatagg gagagaattt tgttagcagt tcgtagttat 5820 cttggagaga atattgaatg gactaatgaa aatgtaaatt taactataaa ctatttaaat 5880 aacagattaa aaaaattata aaaaaattga aaaaatggtg gaaacacttt tttcaatttt 5940 tttgttttat tatttaatat ttgggaaata ttcattctaa ttggtaatca gattttagaa 6000 aacaataaac cettgcatag ggggatcgat atccgtttag getgggeggt gatagettet 6060

cgttcaggca gtacgcctct tttctttcc agacctgagg gaggcggaaa tggtgtgagg 6120 ttcccgggga aaagccaaat aggcgatcgc gggagtgctt tatttgaaga tcaggctatc 6180 actgeggtea atagatttea caatgtgatg getggaeage etgaggaact etegaaceeg 6240 aatggaaaca accagatatt tatgaatcag cgcggctcac atggcgttgt gctggcaaat 6300 gcaggttcat cctctgtctc tatcaatacg gcaacaaaat tgcctgatgg caggtatgac 6360 aataaagctg gagcgggttc atttcaagtg aacgatggta aactgacagg cacgatcaat 6420 gccaggtctg tagctgtgct ttatcctgat gatattgcaa aagcgcctca tgttttcctt 6480 gagaattaca aaacaggtgt aacacattct ttcaatgatc aactgacgat taccttgcgt 6540 gcagatgcga atacaacaaa agccgtttat caaatcaata atggaccaga cgacaggcgt 6600 ttaaggatgg agatcaattc acaatcggaa aaggagatcc aatttggcaa aacatacacc 6660 atcatgttaa aaggaacgaa cagtgatggt gtaacgagga ccgagaaata cagttttgtt 6720 aaaagagato cagogtoggo caaaaccato ggotatoaaa atoogaatoa ttggagocag 6780. gtaaatgctt atatctataa acatgatggg agccgagtaa ttgaattgac cggatcttgg 6840: cctggaaaac caatgactaa aaatgcagac ggaatttaca cgctgacgct gcctgcggac 6900 acggatacaa ccaacgcaaa agtgattttt aataatggca gcgcccaagt gcccggtcag 6960. aatcagcctg gctttgatta cgtgctaaat ggtttatata atgactcggg cttaagcggt 7020. tetetteece attgagggea aggetagaeg ggaettaceg aaagaaacea teaatgatgg 7080 tttctttttt gttcataaat cagacaaaac ttttctcttg caaaagtttg tgaagtgttg,7140 cacaatataa atgtgaaata cttcacaaac aaaaagacat caaagagaaa cataccctgc 7200 aaggatgetg atattgtetg catttgegee ggageaaace aaaaacetgg tgagacaege 7260 cttgaattag tagaaaagaa cttgaagatt ttcaaaggca tcgttagtga agtcatggcg 7320 ageggatttg aeggeatttt ettagteggg eggeaceteg etaaeggatt eaceaeteea 7380 agaattggag ccaatcaatt cttgcggaga actgtgaatg cgcaaaccaa cccttggcag 7440 aacatateca tegegteege cateteeage ageegeaege ggegeatete gggeagegtt 7500 gggtcctggc cacgggtgcg catgatcgtg ctcctgtcgt tgaggacccg gctaggctgg 7560 cggggttgcc ttactggtta gcagaatgaa tcaccgatac gcgagcgaac gtgaagcgac 7620 tgctgctgca aaacgtctgc gacctgagca acaacatgaa tggtcttcgg tttccgtgtt 7680 tegtaaagte tggaaaegeg gaagteageg eeetgeacea ttatgtteeg gatetgeate 7740 gcaggatgct gctggctacc ctgtggaaca cctacatctg tattaacgaa gcgctggcat 7800

tgaccctgag tgatttttct ctggtcccgc cgcatccata ccgccagttg tttaccctca 7860 caacgttcca gtaaccgggc atgttcatca tcagtaaccc gtatcgtgag catcctctct 7920 cgtttcatcg gtatcattac ccccatgaac agaaatcccc cttacacgga ggcatcagtg 7980 accaaacagg aaaaaaccgc ccttaacatg gcccgcttta tcagaagcca gacattaacg 8040 cttctggaga aactcaacga gctggacgcg gatgaacagg cagacatctg tgaatcgctt 8100 cacgaccacg ctgatgaget ttaccgcage tgcctcgcgc gtttcggtga tgacggtgaa 8160 aacetetgae acatgeaget eeeggagaeg gteaeagett gtetgtaage ggatgeeggg 8220 agcagacaag cccgtcaggg cgcgtcagcg ggtgttggcg ggtgtcgggg cgcagccatg 8280 acccagtcac gtagcgatag cggagtgtat actggcttaa ctatgcggca tcagagcaga 8340 ttgtactgag agtgcaccat atgcggtgtg aaataccgca cagatgcgta aggagaaaat 8400 accgcatcag gcgctcttcc gcttcctcgc tcactgactc gctgcgctcg gtcgttcggc 8460 tgcggcgagc ggtatcagct cactcaaagg cggtaatacg gttatccaca gaatcagggg 8520 ataacgcagg aaagaacatg tgagcaaaag gccagcaaaa ggccaggaac cgtaaaaagg 8580 ccgcgttgct ggcgtttttc cataggctcc gccccctga cgagcatcac aaaaatcgac 8640 gctcaagtca gaggtggcga aacccgacag gactataaag ataccaggcg tttccccctg 8700 gaageteest egtgegetet estgtteega eestgeeget taceggatae etgteegest 8760 ttctcccttc gggaagcgtg gcgctttctc atagctcacg ctgtaggtat ctcagttcgg 8820 tgtaggtcgt tcgctccaag ctgggctgtg tgcacgaacc ccccgttcag cccgaccgct 8880 gegeettate eggtaactat egtettgagt eeaaceeggt aagacaegae ttategeeae 8940 tggcagcagc cactggtaac aggattagca gagcgaggta tgtaggcggt gctacagagt 9000 tettgaagtg gtggeetaac tacggetaca ctagaaggac agtatttggt atctgegete 9060 tgctgaagcc agttaccttc ggaaaaagag ttggtagctc ttgatccggc aaacaaacca 9120 ccgctggtag cggtggtttt tttgtttgca agcagcagat tacgcgcaga aaaaaaggat 9180 ctcaagaaga teetttgate ttttetaegg ggtetgaege teagtggaae gaaaacteae 9240 gttaagggat tttggtcatg agattatcaa aaaggatctt cacctagatc cttttaaatt 9300 aaaaatgaag ttttaaatca atctaaagta tatatgagta aacttggtct gacagttacc 9360 aatgottaat cagtgaggoa cotatotoag ogatotgtot atttogttoa tocatagttg 9420 cctgactccc cgtcgtgtag ataactacga tacgggaggg cttaccatct ggccccagtg 9480 ctgcaatgat accgcgagac ccacgctcac cggctccaga tttatcagca ataaaccagc 9540 WO 01/021772 PCT/US00/25993

- 125 -

cagccggaag ggccgagegc agaagtggtc ctgcaacttt atccgcctcc atccagtcta 9600 ttaattgttg ccgggaagct agagtaagta gttcgccagt taatagtttg cgcaacgttg 9660 ttgccattgc tgcaggcatc gtggtgtcac gctcgtcgtt tggtatggct tcattcagct 9720 ccggttccca acgatcaagg cgagttacat gatcccccat gttgtgcaaa aaagcggtta 9780 gctccttcgg tcctccgatc gttgtcaqaa gtaagttggc cgcagtgtta tcactcatgg 9840 ttatggcagc actgcataat tctcttactg tcatgccatc cgtaagatgc ttttctgtga 9900 ctggtgagta ctcaaccaag tcattctgag aatagtgtat gcggcgaccg agttgctctt 9960 gcccggcgtc aacacgggat aataccgcgc cacatagcag aactttaaaa gtgctcatca 10020 ttggaaaacg ttcttcgggg cgaaaactct caaggatctt accgctgttg agatccagtt 10080 cgatgtaacc cactcgtgca cccaactgat cttcagcatc ttttactttc accagcgttt 10140 ctgggtgagc aaaaacagga aggcaaaatg ccgcaaaaaa gggadtaagg gcgacacgga 10200 aatgttgaat actcatactc ttcctttttc aatattattg aagcatttat cagggttatt 10260 gtctcatgag cggatacata tttgaatgta tttagaaaaa taaacaaata ggggttccgc 10320 gcacatttcc ccgaaaagtg ccaccigacg tctaagaaac cattattatc atgacattaa 10380 cctataaaaa taggcgtatc acgaggccct ttcgtcttca agaatt 10426

<220>

<400> 83

ccaatacgca aaccgcctct ccccgcgcgt tggccgattc attaatgcag cccggatccg 60
aaagaagagg atatcccggt tttacagaag gcattggatg atccaaaggt gtccatcaga 120
agacaggctg ttgtgtactt aggaatgatt gaaacacctg atgttcttcc tctattgtat 180
aaagcacttg aggacaaagc tgtatcagtc agaagaacgg ccggagactg cctgtctgat 240
atcggcgatc ctcaagccat tcctgctatg atcaagtcat taagcgactc cagcaagctt 300
gttcgctggc gtgccgccat gttcctgtac gaagtcggcg atgaaagtgc aattgaagct 360
ttgcgcgctg ccgaagatga ccccgaattt gaggtcagcc ttcaagtcaa aatggcgctt 420
gaacgtattg agcatggaga agaagcaaaa ggttctgttt ggaaacaaat gacggaaagc 480

<210> 83

<211> 4191

<212> DNA

<213> Artificial Sequence

<223> Description of Artificial Sequence: Recombinant
pAN263 plasmid

agaaaaaaag gcgaataaag ataaaaaagg tgcagatcat gcaccttttt tatgtgaatt 540 ggtcgaccga aacagcagtt ataaggcatg aagctgtccg gtttttgcaa aagtggctgt 600 gactgtaaaa agaaatcgaa aaagaccgtt ttgtgtgaaa acggtctttt tgtttccttt 660 taaccaactg ccataactcg aggectacct agettccaag aaagatatcc taacagcaca 720 agageggaaa gatgttttgt tetacateea gaacaaeete tgetaaaatt eetgaaaaat 780 tttgcaaaaa gttgttgact ttatctacaa ggtgtggtat aataatctta acaacagcag 840 gacgetetag aggaggagae accatggeag aattaegeag taatatgate acacaaggaa 900 tegatagage teegeacege agtttgette gtgeageagg ggtaaaagaa gaggattteg 960 gcaagccgtt tattgcggtg tgtaattcat acattgatat cgttcccggt catgttcact 1020 tgcaggagtt tgggaaaatc gtaaaagaag caatcagaga agcagggggc gttccgtttg 1080 aatttaatac cattggggta gatgatggca tcgcaatggg gcatatcggt atgagatatt 1140 cgctgccaag ccgtgaaatt atcgcagact ctgtggaaac ggttgtatcc gcacactggt 1200 ttgacggaat ggtctgtatt ccgaactgcg acaaaatcac accgggaatg cttatggcgg 1260 caatgegeat caacatteeg acgatttttg teageggegg acegatggeg geaggaagaa 1320 caagttacgg gcgaaaaatc teeettteet cagtattega aggggtagge geetaecaag 1380 cagggaaaat caacgaaaac gagcttcaag aactagagca gtteggatge ccaaegtgeg 1440 ggtcttgctc aggcatgttt acggcgaact caatgaactg tctgtcagaa gcacttggtc 1500 ttgctttgcc gggtaatgga accattctgg caacatctcc ggaacgcaaa gagtttgtga 1560 gaaaatcggc tgcgcaatta atggaaacga ttcgcaaaga tatcaaaccg cgtgatattg 1620 ttacagtaaa agcgattgat aacgcgtttg cactcgatat ggcgctcgga ggttctacaa 1680 ataccettct tcataccett gecettgeaa acgaageegg egttgaatae tetttagaae 1740 gcattaacga agtcgctgag cgcgtgccgc acttggctaa gctggcgcct gcatcggatg 1800 tgtttattga agatetteae gaagegggeg gegttteage ggetetgaat gagetttega 1860 agaaagaagg agcgcttcat ttagatgcgc tgactgttac aggaaaaact cttggagaaa 1920 ccattgccgg acatgaagta aaggattatg acgtcattca cccgctggat caaccattca 1980 ctgaaaaggg aggccttgct gttttattcg gtaatctagc teeggaegge gctateatta 2040 aaacaggcgg cgtacagaat gggattacaa gacacgaagg gccggctgtc gtattcgatt 2100 ctcaggacga ggcgcttgac ggcattatca accgaaaagt aaaagaaggc gacgttgtca 2160 tcatcagata cgaagggcca aaaggcggac ctggcatgcc ggaaatgctg gcgccaacat 2220 cccaaatcgt tggaatggga ctcgggccaa aagtggcatt gattacggac ggacgttttt 2280 ceggageete eegtggeete teaateggee aegtateaee tgaggeeget gagggeggge 2340 cgcttgcctt tgttgaaaac ggagaccata ttatcgttga tattgaaaaa cgcatcttgg 2400 atgtacaagt gccagaagaa gagtgggaaa aacgaaaagc gaactggaaa ggttttgaac 2460 cgaaagtgaa aaccggctac ctggcacgtt attctaaact tgtgacaagt gccaacaccg 2520 geggtattat gaaaatetag acceetggeg taatagegaa gaggeeegea eegategeee 2580 ttcccaacag ttgcgcagcc tgaatggcga atgagcttgc gccgtcccgt caagtcagcg 2640 taatgctctg ccagtgttac aaccaattaa ccaattctga ttagaaaaac tcatcgagca 2700 tcaaatgaaa ctgcaattta ttcatatcag gattatcaat accatatttt tgaaaaagcc 2760 gtttctgtaa tgaaggagaa aactcaccga ggcagttcca taggatggca agatcctggt 2820 ateggtetge gatteegact egteeaacat caatacaace tattaattte eectegteaa 2880 aaataaggtt atcaagtgag aaatcaccat gagtgacgac tgaatccggt gagaatggca 2940 aaaggttatg catttettte cagaettgtt caacaggeea geeattacge tegteateaa 3000 aatcactcgc atcaaccaaa ccgttattca ttcgtgattg cgcctgagcg agacgaaata 3060 cgcgatcgct gttaaaagga caattacaaa caggaatcga atgcaaccgg cgcaggaaca 3120 ctgccagcgc atcaacaata ttttcacctg aatcaggata ttcttctaat acctggaatg 3180 ctyttttccc agggatcgca gtggtgagta accatgcatc atcaggagta cggataaaat 3240 gettgatggt eggaagagge ataaatteeg teagecagtt tagtetgace ateteatetg 3300 taacatcatt ggcaacgcta cetttgccat gtttcagaaa caactctggc gcatcgggct 3360 ' tcccatacaa tcaatagatt gtcgcacctg attgcccgac attatcgcga gcccatttat 3420 acceatataa atcageatee atgttggaat ttaategegg eetegaegag eaagaegttt 3480 cccgttgaat atggctcata acaccccttg tattactgtt tatgtaagca gacagtttta 3540 ttgttcatga tgatatattt ttatcttgtg caatgtaaca tcagagattt tgagacactc 3600 gacaagatga tettettgag ategttttgg tetgegegta atetettget etgaaaaega 3660 aaaaaccgcc ttgcagggcg gtttttcgaa ggttctctga gctaccaact ctttgaaccg 3720 aggtaactgg cttggaggag cgcagtcacc aaaacttgtc ctttcagttt agccttaacc 3780 ggcgcatgac ttcaagacta actcctctaa atcaattacc agtggctgct gccagtggtg 3840 cttttgcatg tctttccggg ttggactcaa gacgatagtt accggataag gcgcagcqqt 3900 eqqactgaac ggggggttcg tgcatacagt ccagcttgga gcgaactgcc tacccggaac 3960

	tgagt	gtc	ag g	cgtg	gaat	g ag	acaa	acgc	ggc	cata	aca	gegg	aatg	ac a	ccgg	taaac	4020
	cgaaa	aggc	ag g	aaca	ggag	a gc	gcac	gagg	gag	ccgc	cag	ggga	aacg	cc t	ggta	tcttt	4080
	'atagi	tcct	gt c	gggt	ttcg	c ca	ccac	tgat	ttg	agcg	tca	gatt	tegt	ga t	gctt	gtcag	4140
	gggg	gcgg	ag c	ctat	ggaa	a aa	cggc	tttg	ccg	cggc	cct	ctca	cttc	cc ț			4191
		. 04									•						
	<210: <211: <212: <213:	> 70 > DN	2 A	us s	ubti	lis											
•	<220 <221 <222	> CD		699)													
	<400																
	ttg Met 1	tta Leu	ctg Leu	gtt Val	atc Ile 5	gat Asp	gtg Val	Gly ggg	aac Asn	acc Thr 10	aat Asn	act Thr	gta Val	ctt Leu	ggt Gly 15	gta Val	48
	tat , Tyr	cat His	gat Asp	gga Gly 20	aaa Lys	tta Leu	gaa Glu	tat Tyr	His	tgg Trp	cyt Arg	ata Ile	gaa Glu	aca Thr 30	agc Ser	agg Arg	96 ·
	cat His	aaa Lys	aca Thr 35	gaa Glu	gat Asp	gag Glu	ttt Phe	ggg Gly 40	atg Met	att Ile	ttg Leu	cgc Arg	tcc Ser 45	tta Leu	ttt Phe	gat As <u>p</u>	144.
	cac His	tcc Ser 50	ggg Gly	ctt Leu	atg Met	ttt Phe	gaa Glu 55	cag Gln	ata Ile	gat Asp	ggc Gly	att Ile 60	att Ile	att Ile	tcg Ser	tca Ser	192
	gta Val 65	gtg Val	ccg Pro	cca Pro	atc Ile	atg Met 70	ttt Phe	gcg Ala	tta Leu	gaa Glu	aga Arg 75	atg Met	tgc Cys	aca Thr	aaa Lys	tac Tyr 80	240
	ttt Phe	cat His	atc Ile	gag Glu	cct Pro 85	caa Gln	att Ile	gtt Val	ggt Gly	cca Pro 90	ggt Gly	atg Met	aaa Lys	acc Thr	ggt Gly 95	tta Leu	288
	aat Asn	ata Ile	aaa Lys	tat Tyr 100	gac Asp	aat Asn	ccg Pro	aaa Lys	gaa Glu 105	Val	ggg Gly	gca Ala	gac Asp	aga Arg 110	Ile	gta Val	336
	aat Asn	gct Ala	gtc Val 115	gct Ala	gcg Ala	ata Ile	cac His	ttg Leu 120	Tyr	ggc	aat Asr	cca Pro	tta Leu 125	att Ile	gtt Val	gtc Val	384
	gat Asp	ttc Phe 130	Gly	acc Thr	gcc Ala	aca Thr	acg Thr 135	tac Tyr	tgc Cys	tat Tyr	att	gat Asp 140	Glu	aac Asn	aaa Lys	-caa Gln	432
	tac Tyr 145	Met	ggc	Gly	gcg Ala	att Ile 150	Ala	cct Pro	G1Å ååå	att Ile	aca Tha 155	att Ile	tcg Ser	aca Thr	gag •Glu	gcg Ala 160	480

- 129 -

CELL TYP Ser Arg Ala Ala Lays Leu Pro Arg Ile Glu Ile Thr Arg Pro 165 165 165 165 165 165 165 165 165 165																	•
Asp Asn Ile Ile Gly Lys Asn Thr Val Ser Ala Met Gln Ser Gly Ile 180 tta ttt ggc tat gtc ggc caa gga gga agg agg atc gar agg caa ag caa aa Leu Phe Gly Tyr Val Gly Gln Val Glu Gly Ile Val Lys Arg Met Lys 200 tgg cag gca aaa cag gac cca agg tca ttg cga cag gag gcc tgg cgc Trp Gln Ala Lys Gln Asp Pro Arg Ser Leu Arg Gln Glu Ala Trp Arg 210 cgc tca ttg cga acg aat cag att gta tag Arg Ser Leu Arg Gln Glu Ala Trp Arg 225 cgc tca ttg cga acg aat cag att gta tag Arg Ser Leu Arg Ser Leu Arg Gln Glu Ala Trp Arg 225 c210					Ala					Arg					Arg		528
Leu Phe Gly Tyr Val Gly Gln Val Glu Gly Ile Val Lys Arg Met Lys 205 tgg cag gca aaa cag gac caa agg tca ttg cga cag gag gcc tgg cgc 772 Trp Gln Ala Lys Gln Asp Pro Arg Ser Leu Arg Gln Glu Ala Trp Arg 210 cgc tca ttg cga acg aat cag att gta tag 220 cgc tca ttg cga acg aat cag att gta tag 220 c210> 85 c211> 233 c212> PRT c213> Bacillus subtilis c400> 85 Met Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val 1 1 5 Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg 20 His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp 40 Tis Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser 50 Val Val Pro Pro Ile Met Phe Ala Leu Glu Apr Met Cys Thr Lys Tyr 65 Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Asn Pro Leu Ile Val Val 100 Asn Ala Val Ala Ala Ala Ile His Leu Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala				Ile					Val					Ser			576
Trp Gln Ala Lys Gln Asp Pro Arg Ser Leu Arg Gln Glu Ala Trp Arg 210 cgc tca ttg cga acg aat cag att gta tag Arg Ser Leu Arg Ser Leu Arg Thr Asn Gln Ile Val 220 702 225 <210	tta Leu	ttt Phe	Gly	tat Tyr	gtc Val	ggc Gly	caa Gln	Val	gaa Glu	gga Gly	atc Ile	gtt Val	Lys	cga Arg	atg Met	aaa Lys	624
Arg Ser Leu Arg Thr Asn Gln Ile Val 225 <pre> 2210 85 2211 233 2212 PRT 2213 Bacillus subtilis <400 85 Met Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val</pre>	tgg Trp	Gln	gca Ala	aaa Lys	cag Gln	gac Asp	Pro	agg Arg	tca Ser	ttg Leu	cga Arg	Gln	gag Glu	gcc Ala	tgg Trp	cgc Arg	672
<pre><211> 233 <212> PRT <213> Bacillus subtilis </pre> <pre><400> 85 Met Leu Leu Val Ile Asp Val Gly Asn Thr Asn Thr Val Leu Gly Val</pre>	Arg					Asn				tag			,				702
Met Leu Leu Val Ile Asp Val Gly Asp Thr Asp Thr Val Leu Gly Val Tyr His Asp Gly Lys Leu Glu Tyr His Trp Arg Ile Glu Thr Ser Arg His Lys Gly Leu Met Phe Gly Met Ile Leu Arg Ser Leu Phe Asp His Ser Gly Leu Met Phe Glu Ile Asp Gly Ile Ile Ser Ser Val Val Pro Pro Ile Met Phe Ala Leu Gly Pro Gly Thr Lys Tyr Re Cys Thr Lys Tyr Re Lys Tyr Re Lys Tyr Re Lys Tyr Re Lys Tyr <td< td=""><td><211 <212</td><td>l> 23 2> PE</td><td>33 RT</td><td>lus :</td><td>subt:</td><td>ilis</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></td<>	<211 <212	l> 23 2> PE	33 RT	lus :	subt:	ilis											
His Lys Thr Glu Asp Glu Phe Gly Met Ile Leu Arg Ser Leu Phe Asp 40 His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ser Ser 50 Gly Leu Met Phe Asp 70 Pro Ile Met 70 Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr 80 Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu 95 Phe His Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val 110 Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 115 Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala	Met			Val		Asp	Val	Gly	Asn		Asn	Thr	Val	Leu	_	, Val	
His Ser Gly Leu Met Phe Glu Gln Ile Asp Gly Ile Ile Ile Ser Ser Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr 65	Tyr	His	Asp		Lys	Leu	Glu	Tyr		Trp	Arg	Ile	Glu		Ser	Arg	;
Val Val Pro Pro Ile Met Phe Ala Leu Glu Arg Met Cys Thr Lys Tyr 75 Met Ble Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu 95 Man Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val 100 Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 115 Met Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala	His	Lys		Glu	Asp	Glu	Phe	_	Met	Ile	Leu	Arg		Leu	Phe	Asp	
Phe His Ile Glu Pro Gln Ile Val Gly Pro Gly Met Lys Thr Gly Leu 95 Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val 100 Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 115 Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala	His		Gly	Ĺеu	Met	Phe		Gln	Ile	Asp	Gly		Ile	Ile	Ser	Ser	
Asn Ile Lys Tyr Asp Asn Pro Lys Glu Val Gly Ala Asp Arg Ile Val 110 Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 115 Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala		Val	Pro	Pro	Ile		Phe	Ala	Leu	Glu	_	Met	Cys	Thr	Lys	-	
Asn Ala Val Ala Ala Ile His Leu Tyr Gly Asn Pro Leu Ile Val Val 115 Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala	Phe	His	Ile	Glu		Gln	Ile	Val	Gly		Gly	Met	Lys	Thr		Leu	
Asp Phe Gly Thr Ala Thr Thr Tyr Cys Tyr Ile Asp Glu Asn Lys Gln 130 Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala	Asn	Ile	Lys		Asp	Asn	Pro	Lys		Val	Gly	Ala	Asp		Ile	Val	
130 135 140 Tyr Met Gly Gly Ala Ile Ala Pro Gly Ile Thr Ile Ser Thr Glu Ala	Asn	Ala			Ala	Ile	His			Gly	Asn	Pro			Val	Val	
484	Asp		Gly	Thr	Ala	Thr		Tyr	Cys	Tyr	Ile	_		Asn	Lys	Gln	
			Gly	Gly	Ala		Ala	Pro	Gly	Ile		Ile	Ser	Thr	Glu		

O

Leu	Tyr	Ser	Arg	Ala 165	Ala	Lys	Leu	Pro	Arg 170	Ile	Glu	Ile	Thr	Arg 175	Pro	
Asp	Asn	Ile	Ile 180	Gly	Lys	Asn	Thr	Val 185	Ser	Ala	Met	Gln	Ser 190	Gly	Ile	
Leu	Phe	Gly 195	Tyr	Val	Gly	Gln	Val 200	·Glu	Gly	Ile	Val	Lys 205	Arg	Met	Lys	,
Trp	Gln 210	Ala	Lys	Gln	Asp	Pro 215	Arg	Ser	Leu	Arg	.Gln 220	Glu	Ala	Trp	Arg	•
Arg 225	Ser	Leu	Arg	Thr	Asn 230	Gln	Ile	Val				•				•
<21 <21	0> 8: 1> 1 2> D: 3> B:	623	lus	subt	ilis									1		
	1> C	DS 1)	(162	0)								ı				
atq	Tyr	ttg	gca Ala	ttc Phe 5	Gln	gtg Val	caa Gln	aaa Lys	ttg Leu 10	Met	.cgg Arg	tat Tyr	ttg Leu	acg Thr 15	Leu	48
tac Tyr	aag Lys	ata Ile	aag Lys 20	Asp	ctg Leu	aaa Lys	tta Leu	tcg Ser 25	Leu	ecc Pro	Gly	acg Thr	aac Asn 30	Lys	acg Thr	96
caç Glr	caa Gln	ttc Phe 35	Met	gcc Ala	caa Gln	gca Ala	gtc Val 40	. Gly	cgt Arg	tta Lev	act Thr	gga Gly 45	Lys	eco Pro	gga Gly	.144
gto Val	gtg Val	Leu	gto Val	aca Thr	tca Ser	gga Gly 55	Pro	ggt Gly	gco Ala	tct Sei	aac Asr	Leu	gca Ala	a aca	ggc Gly	192
cto Let	ı Let	g aca ı Thr	gcç Ala	g aac a Asr	act Thr	Glu	gga Gly	a gad / Asi	e cct	t gto Val	L Val	: gcg L Ala	cti Lei	gci 1 Ala	gga Gly 80	240
aa: As:	c gto n Val	g ato l Ile	cgt Ar	g Ala 85	туг	cgt Arg	tta Jei	a aaa ı Ly:	a cgo s Aro	g Th	a cat r Hi:	caa s Glr	tci Se:	t tte r Lei 9	g gat u Asp 5	288
aa As:	t gcg n Ala	g gcg a Ala	g cta Le 10	ı Phe	c cag e Glr	rcco Pro	att o Ile	t ace e Th	r Ly	a ta s Ty:	c ag	t gta r Val	ga: Gl:	u Va	t caa l Gln	336
ga As	t gta	a aaa l Lys 115	s As	t ata n Ile	a cco	g gaa Glu	a gct 1 Ala 12	a Va	t ac	a aa r As:	t gca	a tti a Phe 12	e Ar	g at	a gcg e Ala	384
tc	a gc	a gg	g ca	g gc	t ·ggç	g gco	c gc	t tt	t gt	g ag	c tt	t co	g ca	a ga	t gtt	432

Ser	Ala 130	Gly	Gln	Ala	Gly	Ala 135	Ala	Phe	Val	Ser	Phe 140	Pro	Gln	Asp	Val	
gtg Val 145	aat Asn	gaa Glu	gtc Val	aca Thr	aat Asn 150	acg Thr	aaa Lys	aac Asn	gtg Val	cgt Arg 155	gct Ala	gtt Val	gca Ala	gcg Ala	cca Pro 160	480
aaa Lys	ctc Leu	ggt Gly	cct Pro	gca Ala 165	gca Ala	gat Asp	gat Asp	gca Ala	atc Ile 170	agt Ser	gcg Ala	gcc Ala	ata 'Ile	gca Ala 175	aaa Lys	528
atc Ile	caa Gln	aca Thr	gca Ala 180	aaa Lys	ctt Leu	cct Pro	gtc Val	gtt Val 185	ttg Leu	gtc Val	ggc Gly	atg Met	aaa Lys 190	ggc Gly	gga Gly	576
aga Arg	ccg Pro	gaa Glu 195	gca Ala	att Ile	aaa Lys	gcg Ala	gtt Val 200	cgc Arg	aag Lys	ctt Leu	ttg Leu	aaa Lys 205	aag Lys	gtt Val	cag Gln	624
ctt Leu	cca Pro 210	ttt Phe	gtt Val	gaa Glu	aca Thr	tat Tyr 215	caa Gln	gct Ala	gcc Ala	ggt Gly	acc Thr 220	ctt Leu	tct Ser	aga Arg	gat Asp	672
tta Leu 225	gag Glu	gat Asp	caa Gln	tat Tyr	ttt Phe 230	ggc Gly	cgt Arg	atc Ile	ggt Gly	ttg Leu 235	ttc Phe	cgc Arg	aac Asn	cag Gln	cct Pro 240	720
Gly ggc	gat Asp	tta Leu	ctg Leu	cta Leu 245	gag Glu	cag Gln	gca Ala	gat Asp	gtt Val 250	gtt Val	ctg Leu	acg Thr	atc Ile	ggc Gly 255	tat Tyr	768 :
gac Asp	ccg Pro	att Ile	gaa Glu 260	tat Tyr	gat Asp	ccg Pro	aaa Lys	ttc Phe 265	tgg Trp	aat Asn	atc Ile	aat Asn	gga Gly 270	gac Asp	cgg Arg	816
aca Thr	att Ile	atc Ile 275	cat His	tta Leu	gac Asp	gag Glu	att Ile 280	atc Ile	gct Ala	gac Asp	att Ile	gat Asp 285	cat His	gct Ala	tac Tyr	864
cag Gln	cct Pro 290	gat Asp	ctt Leu	gaa Glu	ttg Leu	atc Ile 295	ggt Gly	ga'c Asp	att Ile	ccg Pro	tcc Ser 300	acg Thr	atc Ile	aat Asn	cat His	912
atc Ile 305	gaa Glu	cac His	gat Asp	gct Ala	gtg Val 310	aaa Lys	gtg Val	gaa Glu	ttt Phe	gca Ala 315	gag Glu	cgt Arg	gag Glu	cag Gln	aaa Lys 320	960
atc Ile	ctt Leu	tct Ser	gat Asp	tta Leu 325	aaa Lys	caa Gln	tat Tyr	atg Met	cat His 330	gaa Glu	ggt Gly	gag Glu	cag Gln	gtg Val 335	cct Pro	1008
gca Ala	gat Asp	tgg Trp	aaa Lys 340	tca Ser	gac Asp	aga Arg	gcg Ala	cac His 345	cct Pro	ctt Leu	gaa Glu	atc Ile	gtt Val 350	aaa Lys	gag Glu	1056
ttg Leu	cgt Arg	aat Asn 355	gca Ala	gtc Val	gat Asp	gat Asp	cat His 360	gtt Val	aca Thr	gta Val	act Thr	tgc Cys 365	gat Asp	atc Ile	ggt Gly	1104

tcg Ser	cac His 370	tcc Ser	att Ile	tgg Trp	atg Met	tca Ser 375	cgt Arg	tat Tyr	ttc Phe	ege Arg	agc Ser 380	tac Tyr	gag Glu	ccg Pro	tta Leu	1152
aca Thr 385	tta Leu	atg Met	atc Ile	agt Ser	aac Asn 390	ggt Gly	atg Met	caa Gln	aca Thr	ctc Leu 395	ggc Gly	gtt Val	gcg Ala	ctt Leu	cct Pro 400	1200
tgg Trp	gca Ala	atc Ile	ggc Gly	gct Ala 405	tca Ser	ttg Leu	gtg Val	aaa Lys	ccg Pro 410	gga Gly	gaa Glu	aaa Lys	gtg Val	gtt Val 415	tct Ser	1248
gtc Val	tct Ser	ggt Gly	gac Asp 420	ggc Gly	ggt Gly	ttc Phe	tta Leu	ttc Phe 425	tca Ser	gca Ala	atg Met	gaa Glu	tta Leu 430	gag Glu	aca Thr	1296
gca Ala	gtt Val	cga Arg 435	Leu	aaa Lys	gca Ala	cca Pro	att Ile 440	gta Val	cac His	att Ile	gta Val	tgg Trp 445	aac Asn	gac Asp	agc Ser	1344
aca Thr	tat Tyr 450	Asp	atg Met	gtg Val	cat His	ttc Phe 455	cag Gln	caa Gln	ttg Leu	aaa Lys	aaa Lys 460	tat Tyr	aac Asn	cgt Arg	aca Thr	1392
tct Ser 465	Ala	gtc Val	gat Asp	ttc Phe	gga Gly 470	Asn	atc Ile	, gat Asp	atc Ile	gtg Val 475	Lys	tat Tyr	gcg Ala	gaa Glu	agc Ser 480	1440
tto Phe	gga Gly	gca Ala	act Thr	gcg Ala 485	Leu	cgc Arg	gta Val	gaa 'Glu	tca Ser 490	Pro	gac Asp	cag Gln	ctg Leu	gca Ala 495	Asp	1488
gtt Val	cto Lev	g cgt 1 Arg	caa Glr 500	Gly	atg Met	aac Asn	gct	gaa Glu 505	ı Gly	cct Pro	gto Val	ato Ile	ato 1le 510	Asp	gtc Val	.·1536
cc; Pro	g gtt o Val	gad L Asi 51	э Туг	agt Ser	gat Asp	aac Asn	att 11e 520	Asr	tta Lei	a gca a Ala	a agt a Sei	gac Asp 525) Lys	ctt Lev	ccg Pro	1584
aaa Ly:	a gaa s Gli 530	ı Ph	c ggg	g gaa y Glu	a cto 1 Leo	ato 1 Met 539	Lys	a acq	g aaa c Ly:	a gct	t cte a Lei 540		3			1623
<2	10>	87														

<211> 540

<212> PRT

<213> Bacillus subtilis

<400> 87

Met Tyr Leu Ala Phe Gln Val Gln Lys Leu Met Arg Tyr Leu Thr Leu

Tyr Lys Ile Lys Asp Leu Lys Leu Ser Leu Pro Gly Thr Asn Lys Thr 20 25

Gln	Gln	Phe 35	Met	Ala	Gln	Ala	Val 40	Gly	Arg	Leu	Thr	Gly 45	Lys	Pro	Gly
Val	Val 50	Leu	Val	Thr	Ser	Gly 55	Pro	Gly	Ala	Ser	Asn 60	Leu	Ala	Thr	Gly
Leu 65	Leu	Thr	Ala	Asn	Thr 70	Glu	Gly	Asp	Pro	Val 75	Val	Ala	Leu	Ala	Gly 80
Asn	Val	Ile	Arg	Ala 85	Tyr	Arg	Leu	Lys	Arg 90	Thr	His	Gln	Ser	Leu .95	Asp
Asn	Ala	Ala	Leu 100	Phe	Gln	Pro	Ile	Thr 105	Lys	Tyr	Ser	Val	Glu 110	Val	Gln
Asp	Val	Lys 115	Asn	Ile	Pro	Glu	Ala 120	Val	Thr	Asn	Ala	Phe 125	Arg	Ile	Ala
Ser	Ala 130	Gly	Gln	Ala	Gly	Ala 135	Ala	Phe	Val	Ser	Phe 140	Pro	Gln	Asp	Val
145					150					155			Ala		160
				165	,				170				Ile	175	
			180					185					Lys 190	_	-
		195					200					205	Lys		•
	210					215	•				220		Ser		-
225					230					235			Asn		240
				245					250				Ile	255	-
			260					265					Gly 270		
		275					280					285	His		
	290					295					300		Ile		
305					310					315			Glu		320
				325					330				Gln	335	
Ala	Asp	Trp	Lys	Ser	Asp	Arg	Ala	His	Pro	Leu	Glu	Ile	Val	Lys	Glu

PCT/US00/25993

			340					345			•		350		
Leu	Arg	Asn 355	Ala	Val	Asp	Asp	His 360	Val	Thr	Val	Thr	Cys 3.65	Asp	Ile	Gly
Ser	His 370	Ser	Ile	Trp	Met	Ser 375	Arg	Tyr	Phe	Arg	Ser 380	Tyr	Glu	Pro	Leu
Thr 385	Leu	Met	Ile	Ser	Asn 390	Gly	Met	Gln	Thr	Leu 395	Gly	Val	Ala	Leu	Pro 400
Trp	Ala	Ile	Gly	Ala 405	Ser	Leu	Val	Lys	Pro 410		"Glu	Lys	-Val	Val 415	Ser
Val	Ser	Gly	Asp 420	Gly	Gly	Phe	Leu	Phe 425	Ser	Ala	Met	Glu	Leu 430	Glu	Thr
Ala	Val	Arg 435	Leu	Lys	Ala	Pro	Ile 440	Val	His	Ile	Val	Trp 445	Asn	Asp	Ser
Thr	Tyr 450	Asp	Met	Val	His	Phe 455	Gln	Gln	Łeu	Lys	Lys 460	Tyr	Asn	Arg	Thr
Ser 465	Ala	Val	Asp	Phe	Gly 470	Asn	Ile	Asp	Ile	Val 475	Lys	Tyr	Ala	·Glu	Ser 480
Phe	GŢĀ	Ala	Thr	Ala 485	Leu	Arg	Val.	-Glu	Ser 490		Asp	Gln	Leu	Ala 495	
Val	Leu	Arg	Gln 500		Met	Asn	Ala	-Glu 505		Pro	Val	Ile	1le 510		Val
Pro	Val	Asp 515		Ser	Asp	Asn	11e 520		Leu	Ala	·Ser	Asp 525	Lys	Leu	Pro
Lys	-Glu 530		Gly	Glu	Leu	Met 535		Thr	Lys	a Ala	Leu 540				•.
	.0> 8 .1> 2														

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ribosome
 binding site

<220×

<223> All occurrences of n indicate any nucleotide

<400> 88

agaaaggagg tgannnnnnn atg

23

<210> 89

<211> 7

<212> PRT

```
<213> Artificial Sequence
<223> Description of Artificial Sequence: PanC
      C terminus
<400> 89
Ile Arg Glu Met Glu Arg Ile
                  5
<210> 90
<211> 5
<212> PRT
<213> Artificial Sequence
<223> Description of Artificial Sequence: PanC
      C terminus
<400> 90
Ile Arg Glu Arg Arg
 1
<210> 91
<211> 7
<212> PRT
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: PanC
      C terminus
<400> 91
Ile Arg Arg Lys Glu Val Asn
  1
<210> 92
<211> 6688
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: Recombinant
      pAN336 plasmid
<400> 92
tgcgccgcta cagggcgcgt ccattcgcca ttcaggctgc gcaactqttq qqaaqqcqa 60
tcggtgcggg cctcttcgct attacgccag ctggcgaaag ggggatgtgc tgcaaggcga 120
ttaagttggg taacgccagg gttttcccag tcacgacgtt gtaaaacgac ggccagtgaa 180
ttgtaatacg actcactata gggcgaattg ggcccgacgt cgcatgcacc aggcttctca 240
ggcgctgact tagaaaacct cttgaatgaa gctgcgcttg tagcggctcq tcaaaacaaq 300
```

aaaaaaatcg atgcgcgtga tattgacgaa gcgacggacc gtgtaattgc cggacccgct 360 aagaagagcc gcgttatctc caagaaagaa cgcaatatcg tggcttatca cgaaggcgga 420 cacaccgtta tcggtctcgt tttagatgag gcagatatgg ttcataaagt aacgattgtt 480 cctcggggcc aggctggcgg ttatgctgtt atgctgccaa gagaagaccg ttatttccaa 540 acaaagccgg agctgcttga taaaattgtc ggcctcttgg gcggacgtgt tgctgaagag 600 attatcttcg gtgaagtcag cacaggggcg cacaatgact tecagegtge gaegaatatt 660 gcaagacgaa tggttacaga attcggtatg tcagaaaaac tgggaccgtt gcaatttgga 720 cagtctcagg gcggtcaggt attcttaggc cgtgatttca acaacgaaca gaactacagt 780 gatcaaatcg cttacgaaat tgatcaggaa attcagcgca tcatcaaaga atgttatgag 840 cgtgcgaaac aaatcctgac tgaaaatcgt gacaagcttg aattgattgc ccaaacgctt 900 ctgaaagttg aaacgcttga cgctgaacaa atcaaacacc ttatcgatca tggaacatta 960 cctgagcgta atttctcaga tgatgaaaag aacgatgatg tgaaagtaaa cattctgaca 1020 aaaacagaag aaaagaaaga cgatacgaaa gagtaattcg ctttctttct aaaaaaactg 1080 coggetgacg etggcagttt ttttatgtaa atgattgget cagetgegge ttttacaate 1140 atccaattct ggtatcgatt tgtttacaaa tgagccgctg atcgtytatg gtattgtaga 1200 atgtttgtaa aaagtaaagt agagaaacta ttcaaaagtg gtgatagagg ttgttactgg. 1260 ttatcgatgt ggggaacacc ctgcagctcg agtgaaatac cgcacagatg cgtaaggaga 1320 aaataccgca tcaggcgata aacccagcga accatttgag gtgataggta agattatacc: 1380 gaggtatgaa aacgagaatt ggacctttac agaattactc tatgaagcgc catatttaaa 1440 aagctaccaa gacgaagagg atgaagagga tgaggaggca gattgccttg aatatattga 1500 caatactgat aagataatat atcttttata tagaagatat cgccgtatgt aaggatttca 1560 gggggcaagg cataggcagc gcgcttatca atatatctat agaatgggca aagcataaaa 1620 acttgcatgg actaatgctt gaaacccagg acaataacct tatagcttgt aaattctatc 1680 ataattgtgg tttcaaaatc ggctccgtcg atactatgtt atacgccaac tttcaaaaca 1740 actttgaaaa agctgttttc tggtatttaa ggttttagaa tgcaaggaac agtgaattgg 1800 agttcgtctt gttataatta gcttcttggg gtatctttaa atactgtaga aaagaggaag 1860 gaaataataa atggctaaaa tgagaatatc accggaattg aaaaaactga tcgaaaaata 1920 ccgctgcgta aaagatacgg aaggaatgtc tcctgctaag gtatataagc tggtgggaga 1980 aaatgaaaac ctatatttaa aaatgacgga cagccggtat aaagggacca cctatgatgt 2040

ggaacgggaa aaggacatga tgctatggct ggaaggaaag ctgcctgttc caaaggtcct 2100 gcactttgaa cggcatgatg gctggagcaa tctgctcatg agtgaggccg atggcgtcct 2160 ttgctcggaa gagtatgaag atgaacaaag ccctgaaaag attatcgagc tgtatgcgga 2220 gtgcatcagg ctctttcact ccatcgacat atcggattgt ccctatacga atagcttaga 2280 · cagccgctta gccgaattgg attacttact gaataacgat ctggccgatg tggattgcga 2340 aaactgggaa gaagacactc catttaaaga teegegegag etgtatgatt ttttaaagae 2400 ggaaaagccc gaagaggaac ttgtcttttc ccacggcgac ctgggagaca gcaacatctt 2460 tgtgaaagat ggcaaagtaa gtggctttat tgatcttggg agaagcggca gggcggacaa 2520 gtggtatgac attgccttct gcgtccggtc gatcagggag gatatcgggg aagaacagta 2580 tgtcgagcta ttttttgact tactggggat caagcctgat tgggagaaaa taaaatatta 2640 tattttactg gatgaattgt tttagtacct agatttagat gtctaaaaag ctttaactac 2700 aagettttta gacatetaat ettttetgaa gtacateege aactgteeat actetgatgt. 2760 tttatatett ttetaaaagt tegetagata ggggteeega gegeetaega ggaatttgta 2820 tegecatteg ceatteagge tgegeaactg ttgggaaggg egateggtge ggtegaetgg:2880 caggcaaaac aggacccaag gtcattgcga caggaggcct ggcgccgctc attgcgaacg 2940 aatcagattg tatagacatc gttgatccat tcttaaccct aaaagggctg gaattgattt 3000 atgaaagaaa ccgcgtagga agtgtatagg aggtttagta atggattatt tagtaaaagc 3060 acttgcgtat gacggaaaag ttcgggctta tgcagcgaga acgactgata tggtaaatga 3120 ggggcagaga cgccatggta cgtggccgac agcatccgct gcactaggcc gtacaatgac 3180 agetteactt atgeteggeg etatgetgaa gggegatgat aagetgaeeg tgaaaatega 3240 gggcggaggt ccgatcggag ctattgtagc tgatgccaat gccaaaggag aagtcagagc 3300 ctatgtctct aaccogcaag ttcattttga tttaaatgaa caaggtaagc ttgatgtcag 3360 acgtgcggtt ggaacaaacg gaacgttaag tgtcgtaaaa gatttaggtt tgcgcgagtt 3420 cttcacagga caagtagaaa tcgtttcagg agaattagga gatgatttta cttactatct 3480 tgtgtcatct gagcaggttc cttcatcagt gggcgtaggt gtgctcgtaa atcctqacaa 3540 taccattett geggeagggg getttattat teagetgatg cegggaacag atgatqaaac 3600 aatcacaaaa attgaacagc gtctatctca agtagagccg atttctaagc tcatccaaaa 3660 agggctgaca ccagaagaaa ttttagaaga agtcctaggc gagaaacctg agattttgga 3720 aacgatgeet gteagattee attgeeettg tteaaaagaa eggttegaaa eageeatttt 3780

aggactaggc aaaaaagaaa ttcaagatat gatagaagaa gatggacaag ccgaagcagt 3840 atgccatttt tgtaatgaaa agtacttatt tacaaaagaa gagctggaag ggcttcgtga 3900 ccaaactacc cgctaagctc tttagcgggt ttttaatttg agaaaagggg ctgaaagcag 3960 gtttgaaatc aagaacaatc tggacgcgtt ggatgcatag cttgagtatt ctatagtgtc 4020 acctaaatag cttggcgtaa tcatggtcat agctgtttcc tgtgtgaaat tgttatccgc 4080 tcacaattcc acacaacata cgagccggaa gcataaagtg taaagcctgg ggtgcctaat 4140 gagtgageta acteacatta attgegttge geteactgee egettteeag tegggaaace 4200 tgtcgtgcca gctgcattaa tgaatcggcc aacgcgcggg gagaggcggt ttgcgtattg 4260 ggcgctcttc cgcttcctcg ctcactgact cgctgcgctc ggtcgttcgg ctgcggcgag 4320 cggtatcagc tcactcaaag gcggtaatac ggttatccac agaatcaggg gataacgcag 4380 gaaagaacat gtgagcaaaa ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc 4440 tggcgttttt cgataggctc cgccccctg acgagcatca caaaaatcga cgctcaagtc 4500 agagytggcg aaacccgaca ggactataaa gataccaggc gtttccccct ggaagctccc 4560 togtgogoto tootgttoog accotgoogo tracoggata cotgtoogoo titotoocti: 4620 cgggaagcgt ggcgctttct catagctcac gctgtaggta tctcagttcg gtgtaggtcg 4680 ttegetecaa getgggetgt gtgcacgaac ceceegttea geeegaeege tgegeettat 4740 ccggtaacta tcgtcttgag tccaacccgg taagacacga cttatcgcca ctggcagcag 4800 ccactggtaa caggattagc agagcgaggt atgtaggcgg tgctacagag ttcttgaagt 4860 ggtggcctaa ctacggctac actagaagga cagtatttgg tatctgcgct ctgctgaagc 4920 cagttacctt cggaaaaaga gtt.ggtagct cttgatccgg caaacaaacc accgctggta 4980 gcggtggttt ttttgtttgc aagcagcaga ttacgcgcag aaaaaaagga tctcaagaag 5040 atcetttgat ettttetaeg gggtetgaeg eteagtggaa egaaaaetea egttaaggga 5100 ttttggtcat gagattatca aaaaggatct tcacctagat ccttttaaat taaaaatgaa 5160 gttttaaatc aatctaaagt atatatgagt aaacttggtc tgacagttac caatgcttaa 5220 tcagtgaggc acctatctca gcgatctgtc tatttcgttc atccatagtt gcctgactcc 5280 ccgtcgtgta gataactacg atacgggagg gcttaccatc tggccccagt gctgcaatga 5340 taccgcgaga cccacgctca ccggctccag atttatcagc aataaaccag ccagccggaa 5400 gggccgagcg cagaagtggt cctgcaactt tatccgcctc catccagtct attaattgtt 5460 gccgggaagc tagagtaagt agttcgccag ttaatagttt gcgcaacgtt gttggcattg 5520

ctacaggcat cgtggtgtca cgctcgtcgt ttggtatggc ttcattcagc tccggttccc 5580 aacgatcaag gcgagttaca tgatccccca tgttgtgcaa aaaagcggtt agctccttcg 5640 gtcctccgat cgttgtcaga agtaagttgg ccgcagtgtt atcactcatg gttatgqcag 5700 cactgcataa ttctcttact gtcatgccat ccgtaagatg cttttctgtg actggtgagt 5760 actcaaccaa gtcattctga gaataccgcg cccggcgacc gagttgctct tgcccggcgt 5820 caatacggga taatagtgta tgacatagca gaactttaaa agtgctcatc attggaaaac 5880 gttcttcggg gcgaaaactc tcaaggatct taccgctgtt gagatccagt tcgatgtaac 5940 ccactcgtgc acccaactga tettcagcat ettttaettt caccagegtt tetgggtgag 6000 caaaaacagg aaggcaaaat gccgcaaaaa agggaataag ggcgacacgg aaatgttgaa 6060 tactcatact cttccttttt caatattatt gaagcattta tcagggttat tgtctcatga 6120 gcggatacat atttgaatgt atttagaaaa ataaacaaat aggggttccg cgcacatttc 6180 cccgaaaagt gccacctgta tgcggtgtga aataccgcac agatgcgtaa ggagaaaata 6240 ccqcatcagg cgaaattgta aacgttaata ttttgttaaa attcgcgtta aatatttgtt 6300 aaatcagete atttttaac caataggeeg aaateggeaa aatecettat aaatcaaaag:6360 aatagaccga gatagggttg agtgttgttc cagtttggaa caagagtcca ctattaaaga 6420 acgtggactc caacgtcaaa gggcgaaaaa ccgtctatca gggcgatggc ccactacgtg 6480 aaccatcacc caaatcaagt tttttgcggt cgaggtgccg taaagctcta aatcggaacc 6540 staaagggag cccccgattt agagcttgac ggggaaagcc ggcgaacgtg gcgagaaagg: 6600 aagggaagaa agcgaaagga gcgggcgcta gggcgctggc aagtgtagcg gtcacgctgc 6660 gcgtaaccac cacacccgcc gcgcttaa 6688

<210> 93

<211> 8503

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Recombinant pAN004 plasmid

<400> 93

gaattttgeg geegettega aagetgtaat ataaaaacet tetteaacta aeggggeagg 60 ttagtgacat tagaaaaceg aetgtaaaaa gtacagtegg cattatetea tattataaaa 120 geeagteatt aggeetatet gacaatteet gaatagagtt cataaacaat eetgeatgat 180 aaceateaca aacagaatga tgtacetgta aagatagegg taaatatatt gaattacett 240

tattaatgaa ttttcctgct gtaataatgg gtagaaggta attactatta ttattgatat 300 ttaagttaaa cccagtaaat gaagtccatg gaataataga aagagaaaaa gcattttcag 360 gtataggtgt tttgggaaac aatttccccg aaccattata tttctctaca tcagaaaggt 420 ataaatcata aaactctttg aagtcattct ttacaggagt ccaaatacca gagaatgttt 480 tagatacacc atcaaaaatt gtataaagtg getetaaett atcecaataa cetaactete 540 cqtcqctatt qtaaccaqtt ctaaaaqctq tatttqaqtt tatcaccctt qtcactaaga 600 aaataaatgc agggtaaaat ttatatcctt cttgttttat gtttcggtat aaaacactaa 660 tatcaatttc tgtggttata ctaaaagtcg tttgttggtt caaataatga ttaaatatct 720 cttttctctt ccaattgtct aaatcaattt tattaaagtt catttgatat gcctcctaaa 780 tttttatcta aagtgaattt aggaggetta ettgtetget ttetteatta gaatcaatee 840 tttttaaaa gtcaatatta ctgtaacata aatatatatt ttaaaaatat cccactttat 900 ccaattttcg tttgttgaac taatgggtgc tttagttgaa gaataaagac cacattaaaa 960 aatgtggtct tttgtgtttt tttaaaggat ttgagcgtag cgaaaaatcc ttttcttct 1020 tatcttgata ataagggtaa ctattgaatt cggtaccaag agtttgtaga aacgcaaaaa 1080 ggccatccgt caggatggcc ttctgcttaa tttgatgcct ggcagtttat ggcgggcgtc 1140 ctgcccgcca ccctccgggc cgttgcttcg caacgttcaa atccgctccc ggcggatttg 1200 tectaeteag gagagegtte acegacaaac aacagataaa aogaaaggee cagtetteeg 1260 actgagcctt tcgttttatt tgatgcctgg cagttcccta ctctcgcatg gggagacccc 1320 acactaccat eggegetacg gegttteact tetgagtteg geatggggte aggtgggace 1380 accgcgctac tgccgccagg caaattctgt tttatcagac cgcttctgcg ttctgattta 1440 atctgtatca ggctgaaaat cttctctcat ccgccaaaac aggatcctac ggaaatggag 1500 cggcaaaacc gttttactct caaaatctta aaagaaaacc cccgataaag ggggcttttc 1560 ttctacaaaa ttgtacgggc tggttcgttc cccagcattt gttcaatttt gttttgatca 1620 ttcagaacag ccactttcgg ctcatggctt gccgcttctt gatcagacat cattttgtag 1680 gaaataataa tgaccttatc teetteetge acaaggegtg eggetgeace gtttaageat 1740 atgacgccgc ttccccgttt accaggaata atatacgttt caagacgtgc tccattatta 1800 ttattcacaa tttgtacttt ttcattagga agcattccca cagcatcaat gagatcttca 1860 tcaattgtaa tgcttcccac atagttcagg tttgcttccg taacagttgc cctgtgaagt 1920 ttgccgctca tcattgttcg atacatatta tattctctcc atttctcgaa tatcaataat 1980

WO 01/021772

gatattatct attaaacgcg cttttgaaaa agcaactgca acagcgagaa tcatctttcc 2040 agcaatttca ttcacagget cgagtteegg ataggaataa agetetacat agtetatggt 2100 tecgetagte gttteaatga tatettttge agettttate acegetteag gatetettte 2160 accggcttgg acaagttccg cacttgtttg aagggcccga tacagcttag gcgcttcttt 2220 tettteetea getgttaagt atacattgeg agagettttg getaageegt etteetet 2280 gaccgtatcg acaggaacca attcaatatc catgaagaag tcgctgatta acccatcaac 2340 aacagctacc tgctgcgcat cttttaaacc gaaataggca cgagtcggct tgactagatt 2400 gaaaagcttc gtcagtacga tcgcgacccc gtcaaaatgt ccttctcttg agcgcccgca 2460 taacacgtct gtgcgtcttt ctacatgaat cgtgacattc ttttcaccgg gatacatatc 2520 atgagcatct ggcgtaaaaa gaatatcgac tccggcgttt tctgcaagag ctgcatcccg 2580 ctcaatatcg cgcggatatg cttcaaaatc ttcattaggg ccgaattgtg caggattcac 2640 aaaaatactc ataataacgg cgtcgttttc ttgtcttgct ttgtctgcta aggttaaatg 2700 cccctcatgc agaaacccca tcgtcggaac aaatccgatt gacttgccct ctgaatggta. 2760 tigittiatg gettettea getgigaaat ateagtaate tgteteatet tattiteece 2820 cgtacaagcc gtcaagcact gtctggttca tttgaaagga atgcttttgt tcagggaaag 2880 cacgatgtct tacatcctga acatatccgc tgattgctgt ttcgatggtt tcatcaatgc 2940 gcgtatattg ctttacaaat ttaggtgttc tctcaacacc gtggccgata atatcatgat: 3000 aaacgagaac ttgtccgtcc gctttcacac cagccccgat tccaatgacc ggtatgctta 3060 gcgtctcggc aattttggct gtgagttctg ccggcacaca ttccagcaca agcatcatag 3120 ctcctgcttc ttcgcatttt atactgtctt ctattaattt tttggcgctt tgttcgtctt 3180 tgccctgtac tttatagccg cccagtacgc cgactgactg cggtgtcaaa cctaagtgac 3240 tgactactgg aatgeeteea agegteaatg egegaatgga tteaaacaeg eetteteege 3300 cctcaagett cagtgcgtca gctccgcttt cctgaacgat agccgctgca tttttcagcg 3360 tatcttcctt agacaggtga taagacataa acggcatatc tgtcacaata aaggtattcg 3420 gcgcacccct tttaacggct tttgtatgat ggatcatgtc cgcaactgtc acaccgacag 3480 ttgaatcaag gccgaggacg accattccaa gtgaatcacc gactaaaatc atgtcaactc 3540 ccgcttgttc agcaagttta gctgccggat aatcataagc ggtcagcatg acaatcggtt 3600 gagcgtcctg ctgttgttaa gattattata ccacaccttg tagataaagt caacaacttt 3720

ttgcaaaatt tttcaggaat tttagcagag gttgttctgg atgtagaaca aaacatcttt 3780 ccgctcttgt gctgttagga tatctttctt ggaagctagg taggcctcga gttatggcag 3840 ttggttaaaa ggaaacaaaa agaccgtttt cacacaaaac ggtctttttc gatttctttt 3900 tacagteaca gecaettttg caaaaacegg acagetteat geettataae tgetgttteg 3960 gtcgacaagc ttcgcgaagc ggccgcaaaa ttcactggcc gtcgttttac aacgtcgtga 4020 ctgggaaaac cctggcgtta cccaacttaa tcgccttgca gcacatcccc ctttcgccag 4080 ctggcgtaat agcgaagagg cccgcaccga tcgcccttcc caacagttgc gcagcctgaa 4140 tggcgaatgg cgcctgatgc ggtattttct ccttacgcat ctgtgcggta tttcacaccg 4200 catatggtgc actctcagta caatctgctc tgatgccgca tagttaagcc agccccgaca 4260 cccgccaaca cccgctgact atgcttgtaa accgttttgt gaaaaaattt ttaaaataaa 4320 aaaggggacc tctagggtcc ccaattaatt agtaatataa tctattaaag gtcattcaaa 4380 aggtcateca eeggateage ttagtaaage eetegetaga ttttaatgeg gatgttgega: 4440 ttacttcgcc aactattgcg ataacaagaa aaagccagcc tttcatgata tatctcccaa 4500 tttgtgtagg gcttattatg cacgcttaaa aataataaaa gcagacttga cctgatagtt 4560 rggctgtgag caattatgtg cttagtgcat ctaacgcttg agttaagccg cgccgcgaag 4620 cggcgtcgyc ttgaacgaat tgttagacat tatttgccga ctaccttggt gatctcgcct 4680 ttcacgtagt ggacaaattc ttccaactga tctgcgcgcg aggccaagcg atcttcttct(4740 tgtccaagat aagcctgtct agcttcaagt atgacgggct gatactgggc cggcaggcgc 4800 tocattgecc agteggeage gacateette ggegegattt tgeeggttae tgegetgtae 4860 caaatgeggg acaacgtaag cactacattt egeteatege cageccagte gggeggegag 4920 ttccatagcg ttaaggtttc atttagcgcc tcaaatagat cctgttcagg aaccggatca 4980 aagagtteet cegeegetgg acetaceaag geaaegetat gttetettge ttttgteage 5040 aagatageca gatcaatgte gatcgtgget ggetegaaga tacetgeaag aatgteattg 5100 cgctgccatt ctccaaattg cagttcgcgc ttagctggat aacgccacgg aatgatgtcg 5160 tegtgcacaa caatggtgac ttctacageg eggagaatet egetetetee aggggaagee 5220 quagtiticca amaggicgtt gatcamaget egeogegitg titeatemag cettaeggie 5280 accetaacca gcaaatcaat atcactetet gettcaggc cecatccac tecegeagcce 5340 tacaaatgta cggccagcaa cgtcggttcg agatggcgct cgatgacgcc aactacctct 5400 gatagttgag tegataette ggegateace getteeetea tgatgtttaa etttgtttta 5460

gggcgactgc cctgctgcgt aacatcgttg ctgctccata acatcaaaca tcgacccacg 5520 gcgtaacgcg cttgctgctt ggatgcccga ggcatagact gtaccccaaa aaaacagtca 5580 taacaagcca tgaaaaccgc cactgcgccg ttaccaccgc tgcgttcggt caaggttctg 5640 gaccagttgc gtgagcgcat acgctacttg cattacagct tacgaaccga acaggcttat 5700 gtccactggg ttcgtgcctt catccgtttc cacggtgtgc gtcacccggc aaccttgggc 5760 agcagegaag tegaggeatt tetgteetgg etggegaaeg agegeaaggt tteggtetee 5820 acgeategte aggeattgge ggcettgetg ttettetacg geaaggtget gtgeaeggat 5880 etgecetgge tteaggagat eggaagaeet eggeegtege ggegettgee ggtggtgetg 5940 accoeggatg aagtggtteg cateeteggt tttetggaag gegageateg tttgttegee 6000 cagcttctgt atggaacggg catgcggatc agtgagggtt tgcaactgcg ggtcaaggat 6060 ctggatttcg atcacggcac gatcatcgtg cgggagggca agggctccaa ggatcgggcc 6120 ttgatgttac cegagagett ggcacccage etgegegage aggggaattg atceggtgga. 6180 tgaccttttg aatgaccttt aatagattat attactaatt aattggggac cctagaggtc 6240 cccttttta ttttaaaaat tttttcacaa aacggtttac aagcataacg ggttttgctg 6300 cccgcaaacg ggctgttctg gtgttgctag tttgttatca gaatcgcaga tccggcttca 6360 ggtttgccgg ctgaaagcgc tatttcttcc agaattgcca tgattttttc cccacgggag 6420 gcgtcactgg ctcccgtgtt gtcggcagct ttgattcgat aagcagcatc gcctgtttca, 6480 ggctgtctat gtgtgactgt tgagctgtaa caagttgtct caggtgttca atttcatgtt 6540 ctagttgctt tgttttactg gtttcacctg ttctattagg tgttacatgc tgttcatctg 6600 ttacattgtc gatctgttca tggtgaacag ctttaaatgc accaaaaact cgtaaaagct 6660 ctgatgtatc tatcttttt acaccgtttt catctgtgca tatggacagt tttccctttg 6720 atatctaacg gtgaacagtt gttctacttt tgtttgttag tcttgatgct tcactgatag 6780 atacaagage cataagaace teagateett eegtatttag eeagtatgtt etetagtgtg 6840 gttcgttgtt tttgcgtgag ccatgagaac gaaccattga gatcatgctt actttgcatg 6900 tcactcaaaa attttgcctc aaaactggtg agctgaattt ttgcagttaa agcatcgtgt 6960 agtgtttttc ttagtccgtt acgtaggtag gaatctgatg taatggttgt tggtattttg 7020 teaceattea tittiatetg gitgitetea agiteggita egagateeat tigietatet 7080 agttcaactt ggaaaatcaa cgtatcagtc gggcggcctc gcttatcaac caccaatttc 7140 atattgctgt aagtgtttaa atctttactt attggtttca aaacccattg gttaagcctt 7200 ttaaactcat ggtagttatt ttcaagcatt aacatgaact taaattcatc aaggctaatc 7260 tctatatttg ccttgtgagt tttcttttgt gttagttctt ttaataacca ctcataaatc 7320 ctcatagagt atttgttttc aaaagactta acatgttcca gattatattt tatgaatttt 7380 tttaactgga aaagataagg caatatctct tcactaaaaa ctaattctaa tttttcgctt 7440 gagaacttgg catagtttgt ccactggaaa atctcaaagc ctttaaccaa aggattcctg 7500 atttccacag ttctcgtcat cagetetetg gttgctttag ctaatacace ataagcattt 7560 tecetactga tgtteateat etgagegtat tggttataag tgaaegatae egteegttet 7620 ttccttgtag ggttttcaat cgtggggttg agtagtgcca cacagcataa aattagcttg 7680 gtttcatgct ccgttaagtc atagcgacta atcgctagtt catttgcttt gaaaacaact 7740 aattcagaca tacatctcaa ttggtctagg tgattttaat cactatacca attgagatgg 7800 gctagtcaat gataattact agtccttttc ctttgagttg tgggtatctg taaattctgc 7860 tagacetttg etggaaaact tgtaaattet getagaceet etgtaaatte egetagacet 7920 ttgtgtgttt tttttgttta tattcaagtg gttataattt atagaataaa gaaagaataa 7980 aaaaagataa aaagaataga toocagooot gtgtataact cactacttta ytcagttoog 8040 cagtattaca aaaggatgte gcaaacgetg tttgeteete tacaaaacag acettaaaac 8100 cctaaagget taagtageae ectegeaage tegggeaaat egetgaatat teettttgte 8160 teegaccate aggeacetga gtegetgtet tittegtgae atteagtteg etgegeteae 8220 ggctctggca gtgaatgggg gtaaatggca ctacaggcgc cttttatgga ttcatgcaag 8280 gaaactaccc ataatacaag aaaagcccgt cacgggcttc tcagggcgtt ttatggcggg 8340 tetgetatgt ggtgetatet gaetttttge tgtteageag tteetgeeet etgattttee 8400 agtotgacca ottoggatta tocogtgaca ggtoattoag actggotaat gcaccoagta 8460 aggeageggt ateateaaca ggettaeeeg tettaetgte aac 8503

<210> 94

<211> 7381

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:recombinant
pAN006 plasmid

<400> 94

ttgcggccgc ttcgaaagct gtaatataaa aaccttcttc aactaacggg gcaggttagt 60

gacattagaa aaccgactgt aaaaagtaca gtcggcatta tctcatatta taaaagccag 120 tcattaggcc tatctgacaa ttcctgaata gagttcataa acaatcctgc atgataacca 180 tcacaaacag aatgatgtac ctgtaaagat agcggtaaat atattgaatt acctttatta 240 atgaattttc ctgctgtaat aatgggtaga aggtaattac tattattatt gatatttaag 300 ttaaacccag taaatgaagt ccatggaata atagaaagag aaaaagcatt ttcaggtata 360 ggtgttttgg gaaacaattt ccccgaacca ttatatttct ctacatcaga aaggtataaa 420 tcataaaact ctttgaagtc attctttaca ggagtccaaa taccagagaa tgttttagat 480 acaccatcaa aaattgtata aagtggctct aacttatccc aataacctaa ctctccgtcg 540 ctattgtaac cagttctaaa agctgtattt gagtttatca cccttgtcac taagaaaata 600 aatgcagggt aaaatttata teettettgt tttatgttte ggtataaaae actaatatea 660 atttctgtgg ttatactaaa agtcgtttgt tggttcaaat aatgattaaa tatctctttt 720 ctcttccaat tgtctaaatc aattttatta aagttcattt gatatgcctc ctaaattttt 780 atctaaagtg aatttaggag gettaettgt etgetttett eattagaate aateettett 840 taaaagtcaa tattactgta acataaatat atattttaaa aatatcccac tttatccaat 900 tttcgtttgt tgaactaatg ggtgctttag ttgaaqaata aagaccacat taaaaaatgt 960 ggtcttttgt gtttttttaa aggatttgag cgtagcgaaa aatccttttc tttcttatct 1020 tgataataag ggtaactatt gaatteggta ecaagagttt gtagaaaege aaaaaggeea 1080 teegteagga tggeettetg ettaatttga tgeetggeag tttatggegg gegteetgee 1140 cgccaccete egggeegttg ettegeaacg tteaaateeg eteceggegg atttgteeta 1200 ctcaggagag cgttcaccga caaacaacag ataaaacgaa aggcccagtc tttcgactga 1260 geetttegtt ttatttgatg cetggeagtt ceetactete geatggggag acceeacaet 1320 accateggeg ctaeggegtt teacttetga gtteggeatg gggteaggtg ggaecaeege 1380 gctactgccg ccaggcaaat tctgttttat cagaccgctt ctgcgttctg atttaatctg 1440 tatcaggctg aaaatcttct ctcatccgcc aaaacaggat cctacggaaa tggagcggca 1500 aaaccgtttt actctcaaaa tcttaaaaga aaacccccga taaagggggc ttttcttcta 1560 caaaattgta cgggctggtt cgttccccag catttgttca attttgtttt gatcattcag 1620 aacagccact ttcggctcat ggcttgccgc ttcttgatca gacatcattt tgtaggaaat 1680 aataatgacc ttatctcctt cctgcacaag gcgtgcggct gcaccgttta agcatatgac 1740 gccgcttccc cgtttaccag gaataatata cgtttcaaga cgtgctccat tattattatt 1800

ca	caatttgt	actttttcat	taggaagcat	teccaeagea	tcaatgagat	Clicalcaat	1000
tg	taatgctt	cccacatagt	tcaggtttgc	ttccgtaaca	gttgccctgt	gaagtttgcc	1920
gc	tcatcatt	gttcgataca	tattatattc	tctccatttc	tcgaatatca	ataatgatat	1980
ta	itctattaa	acgcgctttt	gaaaaagcaa	ctgcaacagc	gagaatcatc	tttccagcaa	2040
tt	tcattcac	aggctcgagt	tccggatagg	aataaagctc	tacatagtct	atggttccgc	2100
ta	gtcgtttc	aatgatatct	tttgcagctt	ttatcaccgc	ttcaggatct	ctttcaccgg	2160
ct	tggacaag	ttccgcactt	gtttgaaggg	cccgatacag	cttaggcgct	tetttettt	2220
c	ctcagctgt	taagtataca	ttgcgagagc	ttttggctaa	gccgtcttcc	tctctgaccg	2280
ta	atcgacagg	aaccaattca	atatccatga	agaagteget	gattaaccca	tcaacaacag	2340
ct	tacctgctg	cgcatctttt	aaaccgaaat	aggcacgagt	cggcttgact	agattgaaaa	2400
g	cttcgtcag	tacgatcgcg	accccgtcaa	aatgtccttc	tcttgagcgc	ccgcataaca	2460
C	gtctgtgcg	tctttctaca	tgaatcgtga	cattettte	accgggatac	atatcatgag	2520
C	atctggcgt	aaaaagaata	tcgactccgg	cgttttctgc	aagagctgca	tecegeteaa	2580
t	ategegege	atatgcttca	aaatcttcat	tagggccgaa	ttgtgcagga	ttcacaaaaa	2640
t	actcataat	aacggcgtcg	tttcttgtc	: ttgctttgtc	: tgctaaggtd	: aaatgcccct	2700
С	atgcagaaa	cccatcgtc	ggaacaaato	cgattgactt	gecetetgaa	tggtattgtt	: 2760
t	tatggctto	tttcagctgt	gaaatatcag	g taatctgtct	: catcttattt	teccegtae	2820
а	agccgtcaa	gcactgtctg	gttcatttga	a aaggaatgct	: tttgttcag	g gaaagcacga	2880
t	gtcttaca	t cctgaacata	tccgctgatt	gctgtttcga	a tggtttcato	c aatgcgcgta	2940
t	attgcttt	a caaatttagg	tgttctctca	a acaccgtgg	c cgataatate	c atgataaacq	3000
a	gaacttgt	c cgtccgcttt	cacaccagco	c ccgattccaa	a tgaccggta	t gcttagcgto	30.60
t	cggcaatt	t tggctgtgag	g ttctgccgg	c acacattcc	a gcacaagca	t catagetee	3120
ç	gcttcttcg	c attttatact	gtettetati	t aatttttg	g egetttgtt	c gtctttgcc	3180
ŧ	tgtacttta	t agccgccca	g tacgccgac	t gactgcggt	g tcaaaccta	a gtgactgac	t 3240
ě	actggaatg	c ctccaagegt	caatgcgcg	a atggattca	a acacgcctt	c teegeeete	a 3300
ä	agcttcagt	g cgtcagctco	c gctttcctg	a acgatagec	g ctgcatttt	t cagcgtatc	t 3360
1	tccttagac	a ggtgataaga	a cataaacgg	c atatctgtc	a caataaagg	t attcggcgc	a 3420
,	ccccttta	a cggcttttg	t atgatggat	c atgtccgca	a ctgtcacac	c gacagttga	a 3480
			t topaantna	a tracroset	a aaatcatot	c aact-ceege	t 3540

tgttcagcaa gtttagctgc cggataatca taagcggtca gcatgacaat cggttcttca 3600 gacteettea tittagaaa ateeagtitt gitteatgt titeteetee tetagagegt 3660 cctgctgttg ttaagattat tataccacac cttgtagata aagtcaacaa ctttttgcaa 3720 aatttttcag gaattttagc agaggttgtt ctggatgtag aacaaaacat ctttccgctc 3780 ttgtgctgtt aggatatett tettggaage taggtaggee tegagttatg geagttggtt 3840 aaaaggaaac aaaaagaccg ttttcacaca aaacggtctt tttcgatttc tttttacagt 3900 cacagecaet titgeaaaaa eeggaeaget teatgeetta taaetgetgt tieggiegae 3960 ctgcaggcat gcaagcttcg cgaagcggcc gccgacgcga ggctggatgg ccttccccat 4020 tatgatteti etegetteeg geggeategg gatgeeegeg ttgeaggeea tgetgteeag 4080 gcaygtagat gacgaccatc agggacaget tcaaggateg etegeggete ttaccageet 4140 aacttegate actggacege tgategteae ggegatttat geegeetegg egageaeatg 4200 gaacgggttg gcatggattg taggcgccgc cctatacctt gtctgcctcc ccgcgttgcg 4260 tegeggtgca tggageeggg ccacetegae etgaatggaa geeggeggea cetegetaac 4320 gyattcacca ctccaagaat tggagccaat caattcttgc ggagaactgt gaatgcgcaa 4380 accaaccett ggcagaacat atccategeg teegecatet ecageageeg caegeggege 4440 atetegggea gegttgggte etggeeaegg gtgegeatga tegtgeteet gtegttgagg, 4500 acceggetag getggegggg ttgeettaet ggttageaga atgaateace gataegegag 4560 cgaacgtgaa gcgactgctg ctgcaaaacg tctgcgacct gagcaacaac atgaatggtc 4620 tteggtttee gtgtttegta aagtetggaa aegeggaagt cagegeeetg caecattatg 4680 ttccggatct gcatcgcagg atgctgctgg ctaccctgtg gaacacctac atctgtatta 4740 acgaageget ggcattgace etgagtgatt tttetetggt eeegeegeat eeatacegee 4800 agttgtttac cctcacaacg ttccagtaac cgggcatgtt catcatcagt aacccgtatc 4860 gtgagcatcc tctctcgttt catcggtatc attaccccca tgaacagaaa ttccccctta 4920 cacggaggca tcaagtgacc aaacaggaaa aaaccgccct taacatggcc cgctttatca 4980 gaagccagac attaacgctt ctggagaaac tcaacgagct ggacgcggat gaacaggcag 5040 acatctgtga atcgcttcac gaccacgctg atgagcttta ccgcagctgc ctcgcgcgtt 5100 teggtgatga eggtgaaaac etetgacaca tgeageteee ggagaeggte acagettgte 5160 tgtaagcgga tgccgggagc agacaagccc gtcagggcgc gtcagcgggt gttggcgggt 5220 gtcggggcgc agccatgacc cagtcacgta gcgatagcgg agtgtatact ggcttaacta 5280

tgcggcatca gagcagattg tactgagagt gcaccatatg cggtgtgaaa taccgcacag 5340 atgegtaagg agaaaatace geateaggeg etetteeget teetegetea etgaeteget 5400 gegeteggte gtteggetge ggegageggt ateageteae teaaaggegg taataeggtt 5460 atccacagaa tcaggggata acgcaggaaa gaacatgtga gcaaaaggcc agcaaaaggc 5520 caggaaccgt aaaaaggccg cgttgctggc gtttttccat aggctccgcc cccctgacga 5580' gcatcacaaa aatcgacgct caagtcagag gtggcgaaac ccgacaggac tataaagata 5640 ccaggegttt ccccctggaa gctccctcgt gcgctctcct gttccgaccc tgccgcttac 5700 eggatacetg teegeettte teeetteggg aagegtggeg ettteteata geteaegetg 5760 taggtatete agtteggtgt aggtegtteg etceaagetg ggetgtgtge acgaaeceee 5820 cgttcagccc gaccgctgcg ccttatccgg taactatcgt cttgagtcca acccggtaag 5880 acacgaetta tegecaetgg cageagecae tggtaacagg attageagag egaggtatgt 5940 aggeggtget acagagttet tgaagtggtg geetaactae ggetacacta gaaggacagt 6000 atttggtate tgegetetge tgaagecagt tacettegga aaaagagttg gtagetettg 6060 atccggcaaa caaaccaccg ctggtagcgg tggttttttt gtttgcaagc agcagattac 6120 gegeagaaaa aaaggatete aagaagatee tttgatettt tetaeggggt etgaegetea 6180 gtggaacgaa aactcacgtt aagggatttt ggtcatgaga ttatcaaaaa ggatcttcac 6240 ctagateett ttaaattaaa aatgaagttt taaateaate taaagtatat atgagtaaae 6300 ttggtctgac agttaccaat gcttaatcag tgaggcacct atctcagcga tctgtctatt 6360 tegtteatee atagttgeet gacteeegt egtgtagata actaegatae gggagggett 6420 accatetgge eccagtgetg caatgatace gegagaeeea egeteacegg etccagattt 6480 atcagcaata aaccagccag ccggaagggc cgagcgcaga agtggtcctg caactttatc 6540 cgcctccatc cagtctatta attgttgccg ggaagctaga gtaagtagtt cgccagttaa :6600 tagtttgcgc aacgttgttg ccattgctgc aggcatcgtg gtgtcacgct cgtcgtttgg 6660 tatggettea tteageteeg gtteecaaeg ateaaggega gttacatgat eeeecatgtt 6720 gtgcaaaaaa gcggttagct ccttcggtcc tccgatcgtt gtcagaagta agttggccgc 6780 agtgttatca ctcatggtta tggcagcact gcataattct cttactgtca tgccatccgt 6840 aagatgettt tetgtgaetg gtgagtaete aaccaagtea ttetgagaat agtgtatgeg 6900 gegacegagt tgetettgee eggegteaat aegggataat aeegegeeac atageagaac 6960 tttaaaagtg ctcatcattg gaaaacgttc ttcggggcga aaactctcaa ggatcttacc 7020 WO 01/021772 PCT/US00/25993

- 149 -

gctgttgaga	tccagttcga	tgtaacccac	tcgtgcaccc	aactgatctt	cagcatcttt'	7080
tactttcacc	agcgtttctg	gģtgagcaaa	aacaggaagg	caaaatgccg	caaaaaaggg	7140
aataagggcg	acacggaaat	gttgaatact	catactcttc	ctttttcaat	attattgaag	7200
catttatcag	ggttattgtc	tcatgagcgg	atacatattt	gaatgtattt	agaaaaataa	7260
acaaataggg	gttccgcgca	catttccccg	aaaagtgcca	cctgacgtct	aagaaaccat	7320
tattatcatg	acattaacct	ațaaaaatag	gcgtatcacg	aggccctttc	gtcttcaaga	7380
a						7381

onal Application No PLI/US 100/25993

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C12N15/52 C12N15/53 C12N15/60 C12N15/75 C12N15/54 C12N9/12 C12N9/10 C12N9/88 C12N9/02 C12N9/00 C12P13/06 C12P13/02 C12P7/42 According to International Patent Classification (IPC) or to both national classification and IPC **B. FIELDS SEARCHED** Minimum documentation searched (classification system followed by classification symbols) IPC 7 , C12N C12P Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EMBL, EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, EMBASE C. DOCUMENTS CONSIDERED TO BE RELEVANT Relevant to claim No. Citation of document, with indication, where appropriate, of the relevant passages Category * EP 0 590 857 A (TAKEDA CHEMICAL INDUSTRIES 12,13, Х LTD) 6 April 1994 (1994-04-06) 24,26, cited in the application 27,48, 51,55, 59,60, 71,76 the whole document 1-6. Υ 33-35. 54, 56-58, 62-64. 78-82 page 14, line 1-3 Further documents are listed in the continuation of box C. Patent family members are listed in annex. 'Special categories of cited documents: T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance invention "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family . Date of the actual completion of the international search Date of mailing of the international search report 10 July 2001 22.10. m

Form PCT/ISA/210 (second sheet) (July 1992)

Fax: (+31-70) 340-3016

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31-651 epo nl, Authorized officer

van de Kamp, M

ional Application No

C.(Continua	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	SAHM H ET AL.: "D-Pantothenate synthesis in Corynebacterium glutamicum and use of panBC and genes encoding L-valine synthesis for D-pantothenate overproduction" APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 65, no. 5, May 1999 (1999-05), pages 1973-1979, XP002169517 cited in the application the whole document	12,13, 24,26, 27,48, 51,54, 56,59, 60,71, 76,78,79
		1-6, 33-35, 54, 56-58, 62-64, 78-82
X	SOROKIN A ET AL.: "Sequence analysis of the Bacillus subtilis chromosome region between the serA and kdg loci cloned in a yeast artificial chromosome" MICROBIOLOGY, vol. 142, no. 8, 1996, pages 2005-2016, XP000910121 ISSN: 1350-0872	83-86, 92,93, 95, 97-100, 102,103
Y	page 2011, right-hand column, line 14-20:	1-6, 33-35, 54, 56-58, 62-64, 78-82
x ·	DATABASE EM PRO [Online] EMBL; ID BSYPIA, AC L47709, 23 January 1996 (1996-01-23) HENNER D ET AL.: "Bacillus subtilis (clone YAC15-6B) ypiABF genes, qcrABC genes, ypjABCDEFGHI genes, birA gene, panBCD genes, dinG gene, ypmB gene, aspB gene, asnS gene, dnaD gene, nth gene and ypoC gene, complete cds." XP002171539 page 5, line 43-60 page 10	83-86, 98-100, 102,103
A	BAIGORI M ET AL.: "Isolation and characterization of Bacillus subtilis mutants blocked in the syntehsis of pantothenic acid" JOURNAL OF BACTERIOLOGY, vol. 173, no. 13, July 1991 (1991-07), pages 4240-4242, XP001002216 abstract	99,100, 102,103

PCT/US 90/25993

A W0 98 18954 A (MONSANTO CO) 7 May 1998 (1998-05-07) the whole document A EP 0 224 294 A (GIST BROCADES NV) 3 June 1987 (1987-06-03) the whole document P,X FP 1 096 192 A (DEGUSSA) 7 June 2000 (2000-06-07) EP 1 096 189 A (DEGUSSA) 7 June 2000 (2000-06-07)	C.(Continual	tion) DOCUMENTS CONSIDERED TO BE RELEVANT		4 Rolement to obaim No.
A	Category •	-Citation of document, with indication, where appropriate, of the relevant passages	·	Relevant to claim No.
3 June 1987 (1987-06-03) the whole document P,X EP 1 006 192 A (DEGUSSA) 7 June 2000 (2000-06-07) EP 1 006 192 A (DEGUSSA) 24,26, 27,48, 51, 54-56, 59,60, 71,76, 78,79 examples 2-5 P,X EP 1 006 189 A (DEGUSSA ;KERNFORSCHUNGSANLAGE JUELICH (DE)) 7 June 2000 (2000-06-07) 12,13, 24,26, 27,48, 51,54, 56,59, 60,71, 76,78,79	A	7 May 1998 (1998-05-07)	,	
7 June 2000 (2000-06-07) 24,26, 27,48, 51, 54-56, 59,60, 71,76, 78,79 examples 2-5 P,X EP 1 006 189 A (DEGUSSA; KERNFORSCHUNGSANLAGE JUELICH (DE)) 7 June 2000 (2000-06-07) 12,13, 24,26, 27,48, 51,54, 56,59, 60,71, 76,78,79	A	EP 0 224 294 A (GIST BROCADES NV) 3 June 1987 (1987-06-03) the whole document		
P,X EP 1 006 189 A (DEGUSSA; KERNFORSCHUNGSANLAGE JUELICH (DE)) 7 June 2000 (2000-06-07) 21,13, 24,26, 27,48, 51,54, 56,59, 60,71, 76,78,79	P,X	7 June 2000 (2000-06-07)		24,26, 27,48, 51, 54-56, 59,60, 71,76,
;KERNFORSCHUNGSANLAGE JUELICH (DE)) 7 June 2000 (2000-06-07) 24,26, 27,48, 51,54, 56,59, 60,71, 76,78,79		examples 2-5		
60,71, 76,78,79	P,X	EP 1 006 189 A (DEGUSSA ;KERNFORSCHUNGSANLAGE JUELICH (DE)) 7 June 2000 (2000-06-07)		24,26, 27,48, 51,54, 56.59.
	·	examples 1,7,9		60,71, 76,78,79
	. 1			. ,
	•			·
				-

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

national application No. PCT/US 00/25993

Box 1 Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows:
see additional sheet
As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-6,12,13,24,26-28,33-35,48,51,54-64,71,76,78-86,92,93,95,97100,102,103 (all Partially)
Remark on Protest The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-6,12,13,24,26-28,33-35,48,51,54-64,71,76,78-86, 92,93,95,97-100,102,103 (all partially)

A method of producing pantothenate $\{e.g., 2 \text{ g/L up to } 40 \text{ g/L}\}$ at least), pantoate, or ketopantoate, e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses a ketopantoate hydroxymethyltransferase-encoding gene, e.g. the panB gene, e.g., from Bacillus, under conditions such that said panto-compound is produced, and possibly further recovering the compound. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) which overexpresses a Bacillus (subtilis) ketopantoate hydroxymethyltransferase-encoding gene. A recombinant vector encoding a Bacillus (subtilis) ketopantoate hydroxymethyltransferase-encoding gene operably linked to regulatory sequences, e.g., comprising a nucleic acid sequence according to SEQ ID NO:23 or part of SEQ ID NO:59. An isolated nucleic acid molecule encoding a Bacillus (subtilis) ketopantoate hydroxymethyltransferase, and said isolated ketopantoate hydroxymethyltransferase polypeptide.

2. Claims: 1-6,12,13,24,26-28,33-35,48,51,54-64,71,76,78-86, 92,93,95,97-100,102,103 (all partially)

A method of producing panthotenate (e.g., 2 g/L up to 40 g/L at least), pantoate, or ketopantoate, e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses a pantothenate synthetase-encoding gene, e.g. the panC gene, e.g., from Bacillus, under conditions such that said panto-compound is produced, and possibly further recovering the compound. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) which overexpresses a Bacillus (subtilis) pantothenate synthetase-encoding gene. A recombinant vector encoding a Bacillus (subtilis) pantothenate synthetase-encoding gene operably linked to regulatory sequences, e.g., comprising a nucleic acid sequence according to SEQ ID NO:25 or part of SEQ ID NO:59. An isolated nucleic acid molecule encoding a Bacillus (subtilis) pantothenate synthetase, and said isolated pantothenate synthetase polypeptide.

3. Claims: 1-6,12-14,24,26-28,33-35,48,49,51,54-64,66,71,76, 78-86,92,93,95,97-100,102,103 (all partially); 15, 17,19,23,32,106,107 (both completely)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), pantoate, or ketopantoate, e.g., independent of aspartate or beta-alanine feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses an aspartate-alpha-decarboxylase-encoding gene, e.g., from Bacillus, e.g., the aspartate-alpha-decarboxylase-encoding panD gene from Bacillus subtilis, under conditions such that said panto-compound is produced, and possibly further recovering the compound. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) which overexpresses a Bacillus (subtilis) aspartate-alpha-decarboxylase-encoding gene. A recombinant vector encoding a Bacillus (subtilis) aspartate-alpha-decarboxylase-encoding gene operably linked to regulatory sequences, e.g., comprising a nucleic acid sequence according to SEQ ID NO:27 or part of SEQ ID NO:59. An isolated nucleic acid molecule encoding a Bacillus (subtilis) aspartate-alpha-decarboxylase, and said isolated aspartate-alpha-decarboxylase polypeptide.

4. Claims: 1-6,24,26-28,33-35,48,49,51,54-64,71,76,78-87,92, 93,95,97-100,102,103 (all partially); 7-11,65,101, 104,105 (all completely)

> A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), pantoate, or ketopantoate, e.g., independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses a ketopantoate reductase-encoding gene, e.g., from Bacillus, e.g., the ketopantoate reductase-encoding panE1 gene from Bacillus subtilis, under conditions such that said panto-compound is produced, and possibly further recovering the compound. A recombinant microorganism (e.g., Gram-positive (e.g. belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) which overexpresses a Bacillus (subtilis) ketopantoate reductase-encoding gene. A recombinant vector encoding a Bacillus (subtilis) ketopantoate reductase-encoding gene operably linked to regulatory sequences, e.g., comprising a nucleic acid sequence according to SEQ ID NO:29. An isolated nucleic acid molecule encoding a Bacillus (subtilis)

ketopantoate reductase, and said isolated ketopantoate reductase polypeptide.

5. Claims: 14,16,18,28,48,54-61,66,77-82, 97 (all partially); 20,29 (completely)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of valine or alpha-ketoisovalerate feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a deregulated isoleucine-valine (ilv) pathway, wherein the microorganism overexpresses acetohydroxyacid synthase or is transformed with a vector comprising an ilvBN nucleic acid sequence or an alsS sequence, e.g., from Bacillus, under conditions such that pantothenate is produced, and possibly further recovering the pantothenate. Said microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)), and said vector.

6. Claims: 14,16,18,28,48,54-61,66,77-82, 97 (all partially); 21,30 (completely)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of valine or alpha-ketoisovalerate feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a deregulated isoleucine-valine (ilv) pathway, wherein the microorganism overexpresses acetohydroxyacid isomeroreductase or is transformed with a vector comprising an ilvC nucleic acid sequence, e.g., from Bacillus, under conditions such that pantothenate is produced, and possibly further recovering the pantothenate. Said microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)), and said vector.

7. Claims: 14,16,18,28,48,54-61,66,77-82,97 (partially); 22, 31 (completely)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of valine or alpha-ketoisovalerate feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci

or Streptomyces) or Gram-negative) having a deregulated isoleucine-valine (ilv) pathway, wherein the microorganism overexpresses dihydroxyacid dehydratase or is transformed with a vector comprising an ilvD nucleic acid sequence, e.g., from Bacillus, under conditions such that pantothenate is produced, and possibly further recovering the pantothenate. Said microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)), and said vector.

8. Claims: 25,28,50,54-61 (all partially)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant avtA gene under conditions such that pantothenate is produced, and possibly further recovering the pantothenate.

9. Claims: 25,28,50,54-61 (all partially)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant ilvE gene under conditions such that pantothenate is produced, and possibly further recovering the pantothenate.

10. Claims: 25,28,50,54-61 (all partially)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant ansB gene under conditions such that pantothenate is produced, and possibly further recovering the pantothenate.

11. Claims: 25,28,50,54-61 (all partially)

A method of producing pantothenate (e.g., 2 g/L up to 40 g/L at least), e.g., in a manner independent of precursor feed, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative)

having a mutant alsO gene under conditions such that pantothenate is produced, and possibly further recovering the pantothenate.

12. Claims: 36,37,54-59,61 (all partially)

A method of producing beta-alanine, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses an aspartate-alpha-decarboxylase-encoding gene, under conditions such that beta-alanine is produced, wherein the asparte-alpha-decarboxylase-overexpressing microorganism has a mutation in a nucleic acid sequence encoding ketopantoate hydroxymethyltransferase (panB), and possibly further recovering the compound.

13. Claims: 36,37,54-59,61 (all partially)

A method of producing beta-alanine, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses an aspartate-alpha-decarboxylase-encoding gene, under conditions such that beta-alanine is produced, wherein the asparte-alpha-decarboxylase-overexpressing microorganism has a mutation in a nucleic acid sequence encoding ketopantoate reductase (panE), and possibly further recovering the compound.

14. Claims: 36,37,54-59,61 (all partially)

A method of producing beta-alanine, comprising culturing a microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) which overexpresses an aspartate-alpha-decarboxylase-encoding gene, under conditions such that beta-alanine is produced, wherein the asparte-alpha-decarboxylase-overexpressing microorganism has a mutation in a nucleic acid sequence encoding pantothenate synthetase (panC), and possibly further recovering the compound.

15. Claim: 38 (completely)

A method of producing beta-alanine comprising contacting a composition comprising aspartate with an isolated Bacillus aspartate-alpha-decarboxylase enzyme under conditions such that beta-alanine is produced.

16. Claims: 41,44-47,51,53,54-61,69,71,72,75,78-81, 97 (all partially); 39,43,52,67,70,74,88-91, 108-110 (all completely)

A method for producing or for enhancing production of ketopantoate, pantoate, or pantothenate (e.g., 10, 20 or 40 g/L at least), comprising culturing a mutant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant pantothenate kinase-encoding coaX gene, under conditions such that said panto-compound is produced or that production is enhanced, and possibly further recovering the compound. A method for identifying compounds which modulate pantothenate kinase activity comprising contacting a recombinant cell expressing the coaX gene, possibly further comprising a mutant coaA gene encoding a pantothenate kinase with reduced activity, with a test compound and determining the ability of the test compound to modulate pantothenate kinase activity in said cell. A recombinant microorganism having a mutant coaX gene encoding a pantothenate kinase with reduced activity. A vector comprising a mutant coaX gene encoding a pantothenate kinase with reduced activity, possibly further comprising regulatory sequences. A recombinant microorganism comprising a vector comprising an isolated coaX gene (e.g., from Bacillus (subtilis)), and said vector, possibly further comprising regulatory sequences, e.g., a constitutively active promoter. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) that overproduces a panto-compound having a mutation in a coaX gene that results in a reduced level of pantothenate kinase activity, resulting in a decrease in the capacity of the microorganism to synthesize coenzyme A. An isolated nucleic acid molecule comprising a (mutant) coaX gene, and an isolated pantothenate kinase protein encoded by a coaX gene.

17. Claims: 41,44-47,51,53-61,69,71,72,75,78-81, 97 (all partially); 40,42,68,73 (all completely)

A method for producing or for enhancing production of a panto-compound, e.g., ketopantoate, pantoate or pantothenate (e.g., 10, 20 or 40 g/L at least), comprising culturing a mutant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus, Corynebacterium, Lactobacillus, Lactococci or Streptomyces) or Gram-negative) having a mutant pantothenate kinase-encoding coaA gene, under conditions such that the panto-compound is produced or that production is enhanced, and possibly further recovering the panto-compound. A recombinant microorganism having a mutant coaA gene encoding a pantothenate kinase with reduced

activity. A recombinant microorganism (e.g., Gram-positive (e.g., belonging to the genus Bacillus (e.g., Bacillus subtilis), Corynebacterium, Lactobacillus, Lactococci or Streptomyces)) that overproduces a panto-compound having a mutation in a coaA gene that results in a reduced level of pantothenate kinase activity, resulting in a decrease in the capacity of the microorganism to synthesize coenzyme A. A vector containing a (mutated) coaA gene.

18. Claim: 94 (partially)

A vector containing regulatory sequences comprising the constitutively active promoter Pveg (SEQ ID NO:41).

19. Claim: 94 (partially)

A vector containing regulatory sequences comprising the constitutively active promoter P15 (SEQ ID NO:39).

20. Claim: 94 (partially)

A vector containing regulatory sequences comprising the constitutively active promoter P26 (SEQ ID NO:40).

21. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:49.

22. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:50.

23. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:51.

24. Claim : 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:52.

25. Claim: 96 (partially)

A vector containing regulatory sequences comprising an

artificial RBS according to SEQ ID NO:53.

26. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID N0:54.

27. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID N0:55.

28. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID NO:56.

29. Claim: 96 (partially)

A vector containing regulatory sequences comprising an artificial RBS according to SEQ ID N0:57.

In... Application No
PCT/US 00/25993

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0590857	A	06-04-1994	CN 1095105 A,B CN 1225394 A CN 1225388 A CN 1225392 A CN 1225358 A JP 6261772 A US 5518906 A	16-11-1994 11-08-1999 11-08-1999 11-08-1999 11-08-1999 20-09-1994 21-05-1996
WO 9818954	Α	07-05-1998	US 5834259 A EP 0934426 A JP 2001503979 T	10-11-1998 11-08-1999 27-03-2001
EP 0224294	A	03-06-1987	AT 56989 T DE 3674548 D DK 533286 A ES 2031818 T FI 864547 A ' JP 62181789 A PT 83714 A,B US 5024943 A	15-10-1990 31-10-1990 09-05-1987 01-01-1993 09-05-1987 10-08-1987 01-12-1986 18-06-1991
EP 1006192	A	07-06-2000	DE 19855313 A BR 9905776 A CN 1256314 A HU 9904447 A JP 2000228990 A SK 163399 A US 6184007 B	08-06-2000 24-04-2001 14-06-2000 28-11-2000 22-08-2000 11-07-2000 06-02-2001
EP 1006189	Α	07-06-2000	DE 19855312 A BR 9905783 A CN 1256313 A HU 9904448 A JP 2000166580 A SK 164099 A US 6177264 B	08-06-2000 24-04-2001 14-06-2000 28-11-2000 20-06-2000 11-07-2000 23-01-2001

THIS PAGE BLANK (USPTO)